arXiv:hep-th/0408228v2 6 Sep 2004

Aspects of particle mixing in Quantum Field Theory

Antonio Capolupo

Dipartimento di Fisica dell'Universita' di Salerno
and Istituto Nazionale di Fisica Nucleare
I-84100 Salerno, Italy

Abstract

The results obtained on the particle mixing in Quantum Field Theory are reviewed.

The Quantum Field Theoretical formulation of fermion and boson mixed fields is analyzed in detail and new oscillation formulas exhibiting corrections with respect to the usual quantum mechanical ones are presented. It is proved that the space for the mixed field states is unitary inequivalent to the state space where the unmixed field operators are defined. The structure of the currents and charges for the charged mixed fields is studied. Oscillation formulas for neutral fields are also derived. Moreover the study some aspects of three flavor neutrino mixing is presented, particular emphasis is given to the related algebraic structures and their deformation in the presence of CP violation. The non-perturbative vacuum structure associated with neutrino mixing is shown to lead to a non-zero contribution to the value of the cosmological constant. Finally, phenomenological aspects of the non-perturbative effects are analyzed. The systems where this phenomena could be detected are the $\eta - \eta'$ and $\phi - \omega$ mesons.

Acknowledgments

I would like to thank Prof.Giuseppe Vitiello, my advisor, for introducing me to the Quantum Field Theory of particle mixing. He motivated me to work at my best while allowing me the freedom to pursue those areas that interested me the most. I further owe thanks to Dr.Massimo Blasone for his advice, help and numerous discussions. I am grateful to have had the opportunity to work with the nuclear and particle theory group at North Carolina State University under the leadership of Prof.Chueng-Ryong Ji and I thank Yuriy Mishchenko of for his collaboration during my permanence in USA. This work is dedicated to my family and to Ada.

Contents

Introduction **1**

PART 1. FERMION MIXING **6**

1 Neutrino oscillations in Quantum Mechanics **8**
1.1 Introduction 8
1.2 Leptonic numbers 9
1.3 Majorana neutrinos 10
1.4 Dirac neutrinos 14
1.5 Neutrino oscillations 16

2 Quantum field theory of fermion mixing **20**
2.1 Introduction 20
2.2 The vacuum structure for fermion mixing 21
2.3 Generalization of mixing transformations 29
2.4 The current structure for field mixing 32
2.5 The exact formula for neutrino mixing 35
2.6 Discussion 37

3 Quantum Field Theory of three flavor neutrino mixing and oscillations with CP violati
3.1 Introduction 38
3.2 Three flavor fermion mixing 39
3.3 The parameterizations of the three flavor mixing matrix 44
3.4 Currents and charges for three flavor fermion mixing 46
3.5 Neutrino Oscillations 49
3.6 CP and T violations in neutrino oscillations 54
3.7 Conclusions 56

4 Group theoretical aspects of neutrino mixing in Quantum Field Theory **58**
4.1 Introduction 58
4.2 SU(3) deformed algebra 58
4.3 Group representations and the oscillation formula 61

4.3.1 Two flavors 61
4.3.2 Three flavors 63
4.4 Conclusions 66

5 Mixing and oscillations of Majorana fermions 67
5.1 Introduction 67
5.2 Majorana fermions 68
5.3 Discussion and Conclusions 72

6 Neutrino mixing and cosmological constant 74
6.1 Introduction 74
6.2 Neutrino mixing contribution to the cosmological constant 75
6.3 Conclusions 79

PART 2. BOSON MIXING 82

7 Boson mixing in Quantum Mechanics 84
7.1 Usual representation of boson oscillations: the Gell-Mann Pais model 84

8 Quantum field theory of boson mixing 88
8.1 Introduction 88
8.2 Mixing of boson fields in QFT 89
8.2.1 Arbitrary mass parameterization 93
8.3 The currents for mixed boson fields 94
8.4 The oscillation formula for mixed bosons 97
8.4.1 The oscillation formula of Binger and Ji 98
8.4.2 The exact oscillation formula 98
8.5 Mixing of neutral particles 103
8.6 Conclusions 109

9 Phenomenology of flavor oscillations with nonperturbative Quantum Field Theoretic e
9.1 Introduction 110
9.2 Phenomenological application of the nonperturbative oscillation formula111
9.2.1 $K^0 - \bar{K}^0$, $B^0 - \bar{B}^0$ and $D^0 - \bar{D}^0$ mixing 111
9.2.2 The effect of the particle decay 112
9.2.3 The nonperturbative effect in the boson sector 112
The magnitude of the nonperturbative term 113
9.2.4 The nonperturbative effect in fermion sector 114
9.3 Conclusions 116

A Double beta decay 117

B Useful formulas for the generation of the flavor vacuum 118

C Wave function and $V_{\mathbf{k}}$ 119

D The flavor vacuum 120

E Orthogonality of flavor vacuum states and flavor states at different times 121

F Useful relations for three flavors fermion mixing 123

G Anti-neutrino oscillation formulas 126

H Useful formulas for the generation of the mixing matrix 128

I Arbitrary mass parameterization and physical quantities 129

J Temporal component of spinorial derivative 131

K Orthogonality between mass and flavor vacua in boson mixing 133

Bibliography 135

Introduction

The particle mixing and oscillation [1, 2, 3, 4] are one of the most important topics in modern Particle Physics: these phenomena are observed when a source create a particle that is a mixture of two or more mass eigenstate and different mixing are observed in a detector.

The oscillations are observed experimentally in the meson sector, for kaons, B^0, D^0, and the system $\eta - \eta'$, and the evidence of the neutrino oscillations seems now certain [5, 6, 7, 8, 9, 10, 11, 12, 13]. The Standard Model incorporates the mixing of fermion fields through the Kobayashi - Maskawa [14] mixing of 3 quark flavors, i.e. a generalization of the original Cabibbo [15] mixing matrix between the d and s quarks. Recently, the mixing phenomenon helped to provide a vital insight into the puzzle of solar neutrinos [11] and neutrino masses [13]. In the boson sector, the mixing of K^0 and $\bar{K}^0$ mesons via weak currents provided the first evidence of CP violation [2] and $B^0 - \bar{B}^0$ mixing plays an important role in determining the precise profile of a CKM [14, 15] unitarity triangle [16] in Wolfenstein parameter space [17]. In this light, beyond any doubt, the mixing of flavors at this date is the most promising phenomenon of the physics beyond the Standard Model.

Regarding the vanishing magnitudes of the expected violation effects (such as neutrino mass differences or unitarity violation in CKM matrix), it is imperative that the theoretical aspects of the quantum mixing were very well understood.

There are still many unanswered questions about the physics of the mixing and the oscillations in particular from a theoretical point of view.

Let us remind some basic facts about particle mixing and oscillations.

In the Standard Model, the neutrinos appear among the fundamental constituents, together with the corresponding charged leptons and the quarks. The idea of the neutrino was first introduced by W.Pauli in 1930 in order to save the energy and momentum conservation laws in beta-decay process of atomic nuclei and was first observed in 1956 by Reines and Cowen in the process $\bar{\nu_e} + p \rightarrow n + e^+$.

Traditionally, it has been supposed that neutrinos are massless fermions, and therefore are described by two component Weyl spinors, but according to recent experimental data [5, 6, 7, 8, 9, 10, 11, 12, 13], neutrinos may have a mass. The fact that they are electrically neutral, makes possible the existence of two different types of massive neutrinos, namely Dirac or Majorana neutrinos. In the first case, the massive neutrino would be described by a four component Dirac spinor, similar to the one describing the electron. In the case of Majorana neutrino, the spinor has only two components, since neutrino and antineutrino are identified.

To have oscillations it is necessary that neutrino masses are not zero and mixing is present, i.e. that the neutrinos belonging to different generations do have a mixed mass term. Then the time evolution of a neutrino mixed state would lead to flavor oscillations, i.e. to a conversion of a neutrino of one flavor into one of another flavor.

Several questions are still open. For example, the nature of the neutrino mass (Dirac or Majorana) is not understood; it is not justified the smallness of the neutrino masses with respect to those of the other leptons; it is not clear how the mixing arises and also it is difficult to understand the large mixing angles necessary to fit the latest experimental data [5, 6, 7, 8, 9, 10, 11, 12, 13].

On the other hand, from a theoretical point of view, mixing is extremely interesting, since it appear to be a fundamental problem in particle physics.

From a mathematical point of view, the problem in defining properly the Hilbert space for mixed particles was resolved only recently [18, 19, 20, 21].

It is indeed, a problem of unitarily inequivalent representations: the choice of a proper Hilbert space, is involved when mixed fields are considered. This is due to the peculiar mathematical structure of Quantum Field Theory (QFT), where many inequivalent representations (many different Hilbert space) are allowed for a given dynamics [22, 23].

This situation contrast the one of Quantum Mechanics, where, due to the von Neumann theorem, only one Hilbert space is admitted due to the finiteness of the number of the degrees of freedom of the system under consideration.

It was noted, that in QFT the use of the usual perturbation approach to mixing, where the flavor quantum states are defined in the Fock space of the mass-eigenstate fields, leads to certain difficulty [20]. That difficulty originates from the unitary inequivalence between the quantum states of flavor and the quantum states of energy [20]. Thus, it was suggested that the problem may be fixed [20, 24] by defining the flavor states in a ("flavor") space "different", i.e. unitarily inequivalent from the mass state space.

The non trivial nature of the mixing transformations manifests itself also in the case of the mixing of boson fields. The mesons that oscillates are: $K^0 - \bar{K}^0$, $B^0 - \bar{B}^0$, $D^0 - \bar{D}^0$, and the system $\eta - \eta'$.

Extensive study of the mixing phenomenon with the new definition of the flavor states both for neutrinos and mesons has been carried out and the general formulation of such theory for bosons and fermions has been also suggested [20, 21, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33]. Explicit form of the flavor vacuum has been found and it has been shown that indeed the quantum states of flavor are always unitarily inequivalent to those of energy.

In the framework of QFT, exact oscillation formulae has been calculated for neutrinos [20, 30] and bosons [26] case and it turned out to have an additional oscillating term and energy dependent amplitudes, in contrast with the usual quantum mechanical formulas, (Pontecorvo formula [1] in the neutrino case, Gell-Mann Pais formula [3] in the boson case), which are however recovered in the relativistic limit.

Moreover, the non-perturbative vacuum structure associated with neutrino mixing leads to a non zero contribution to the value of the cosmological constant [34].

The thesis is organized in the following way:

Part 1 is devoted to the fermion mixing; in particular, in Chapter 1 we present the neutrino oscillations in Quantum Mechanics and the Bilenky and Pontecorvo oscillation formulae [1].

The Quantum Field Theoretical formulation of fermion mixing is analyzed in Chapter 2, where the unitary inequivalence of the flavor and mass representations is proved, and the currents and charges for mixed fields are introduced and used to derive the new oscillation formulae, exhibiting corrections with respect to the usual quantum mechanical ones [18, 19, 20, 21, 24, 25, 28, 29].

The extension of the results to the case of three flavor neutrino mixing and a study of the algebraic structures and their deformations in the presence of CP violation [30, 31, 35] are given in Chapters 3 and 4, where by use of the generators of the mixing transformations, are recovered all the known parametrizations of the three flavor mixing matrix [14, 15, 17, 36, 37, 38, 39, 40] and a number of new ones are found [30]. In Chapter 5 it is discussed the mixing and the oscillations of Majorana fermions in Quantum Field Theory [41]. In Chapter 6 we show that the neutrino mixing leads to a non-zero contribution to the value of the cosmological constant [34].

In Part 2 the boson mixing is analyzed.

The usual representation of boson oscillations in Quantum Mechanics, the Gell-Mann Pais model [3], is presented in Chapter 7.

In Chapter 8 we treat the quantum field theory of boson mixing [26, 42, 43] and we derive the exact oscillation formulae. In Chapter 9 we analyze phenomenological aspects of the non-perturbative effect. We argue that the system where the new oscillation formulae could be best checked are $\eta - \eta'$ and $\phi - \omega$ mesons [43].

Finally, details of the mathematical formalism are confined to the Appendices A-K.

The corrections introduced by the present formalism to the usual Quantum Mechanics formulae are in principle experimentally testable. The fact that these corrections may be quantitatively below the experimental accuracy reachable at the present state of the art in the detection of the neutrino and boson oscillations, does not justify neglecting them in the analysis of the particle mixing and oscillation mechanism. The exact oscillation formulae derived in QFT are the result of a mathematically consistent analysis which cannot be ignored in a correct treatment of the field mixing phenomenon. The QFT formalism accounts for all the known parameterizations of the mixing matrix and explains their origin and their reciprocal relations, thus unifying the phenomenological proposals scattered in the literature where such parameterizations have been presented. Moreover, the QFT formalism clearly points to the truly non-perturbative character of the particle mixing phenomenon. A lot of Physics must be there waiting to be discovered.

PART 1. FERMION MIXING

Chapter 1

Neutrino oscillations in Quantum Mechanics

In this chapter we present the theoretical model adopted to describe the neutrino mixing and oscillations in Quantum Mechanics. In particular, the neutrino oscillation will be treated by using the formalism of Bilenky and Pontecorvo [1].

1.1 Introduction

In the weak interaction it is assumed that the leptonic numbers are strictly conserved. Then, the neutrino oscillations, that is the mutation of the neutrino flavor, may arise if, in addition to the usual weak interaction, a super-weak interaction which does not conserve leptonic numbers is also taking place.

In such a case, the neutrino masses are assumed to be different from zero, and the state vectors of the ordinary electron, muon and tau neutrinos ν_e, ν_μ, ν_τ (flavor states), are superpositions of the state vectors of neutrinos ν_1, ν_2, ν_3 (mass states), which propagate with different frequencies due to different masses m_1, m_2, m_3. Then, if a beam, of muon neutrinos, for example, is produced in a weak process, at a certain distance from the production place, the beam will be a coherent superposition of ν_e, ν_μ and ν_τ, that is, there arise oscillations $\nu_e \leftrightarrows \nu_\mu \leftrightarrows \nu_\tau$.

The first theories of neutrino oscillations are formulated with Majorana neutrinos and Dirac neutrinos and consider only two type of neutrinos, ν_e and ν_μ. In both theories, possible oscillations $\nu_e \leftrightarrows \nu_\mu$, $\bar{\nu}_e \leftrightarrows \bar{\nu}_\mu$ are described by identical expressions, in which two parameters are present, the mixing angle θ and the difference $\Delta m^2 \equiv |m_1^2 - m_2^2|$ of neutrino masses squared.

We will consider the theories in which the neutrino oscillations arise [1]. They are all based on the assumption that, in addition to the usual weak interaction, there exist also an interaction which does not conserve the leptonic numbers. In agreement with experimental data, it is assumed that this additional interaction is weaker (super-weak) than the usual weak interaction.

Section 1.2 is devoted to describe the usual weak interaction, in Sections 1.3 and 1.4, the theories with mixing of Majorana neutrinos and Dirac neutrinos, respectively, are exposed and in Section 1.5 the Pontecorvo oscillation formula is presented.

1.2 Leptonic numbers

The usual weak interaction hamiltonian is

$$H_w = H_w^c + H_w^0 \tag{1.1}$$

with

$$H_w^c = \frac{G}{\sqrt{2}} J^\alpha \bar{J}_\alpha \tag{1.2}$$

where

$$J_\alpha = (\bar{\nu}_{eL}\gamma_\alpha e_L) + (\bar{\nu}_{\mu L}\gamma_\alpha \mu_L) + J_\alpha^h \tag{1.3}$$

is the weak charged current, J_α^h is the hadronic current and

$$\Psi_L = \frac{1}{2}\left(1+\gamma^5\right)\Psi \tag{1.4}$$

is the left-handed component of the field operator Ψ. $G \simeq 10^{-5} M_p^{-2}$ is the weak interaction constant, with M_p proton mass.

The second term of Eq.(1.1), H_w^0 is the neutral current contribution, the structure of which is of no relevance in this contest.

Let us remember that the neutral current for neutrinos is

$$J_{(0)}^\mu\left(\nu\right) = \frac{1}{2}\left[\bar{\nu}\ \gamma^\mu \left(1-\gamma^5\right)\nu\right] \tag{1.5}$$

and for electrons

$$J_{(0)}^\mu\left(e\right) = \frac{1}{2}\left[\bar{e}\ \gamma^\mu \left(g_V - g_A\,\gamma^5\right) e\right] \tag{1.6}$$

where

$$g_V = -1 + 4\sin^2\theta_W \tag{1.7}$$

with $\sin^2\theta_W = (0.23 \pm 0.02)$, θ_W is Weinberg angle and $g_A = -1$.

The γ_j are the Dirac matrices defined as

$$\gamma_j = \begin{pmatrix} 0 & -i\sigma_j \\ i\sigma_j & 0 \end{pmatrix} \tag{1.8}$$

with $j = 1, 2, 3$, σ_j Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \qquad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \tag{1.9}$$

moreover

$$\gamma_4 = \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \qquad \text{with} \qquad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \tag{1.10}$$

and

$$\gamma_5 = \frac{i}{4!}\varepsilon_{\mu\nu\lambda\eta}\gamma^\mu\gamma^\nu\gamma^\lambda\gamma^\eta, \tag{1.11}$$

where $\varepsilon_{\mu\nu\lambda\eta}$ is the totally antisymmetric Levi-Civita tensor defined as

$\varepsilon_{\alpha\beta\gamma\delta} = 0$ for any two equal indices

$\varepsilon_{0123} = 1$

$\varepsilon_{\alpha\beta\gamma\delta}$ changes sign under interchange of two consecutive indices.

The interaction given by the Hamiltonian H_w^c, does conserve separately the sum of the electron L_e, and muon L_μ lepton numbers

$$\sum_i L_e^{(i)} = constant, \tag{1.12}$$

$$\sum_i L_\mu^{(i)} = constant. \tag{1.13}$$

The experiments in which the neutrino oscillations are searched for will allow one to test the hypothesis on the existence of an interaction non-conserving the lepton numbers.

1.3 Majorana neutrinos

The first theory of neutrino oscillations was based on two component neutrino theory [1].

According to the theory of the Majorana neutrinos oscillations, only left-handed components of the neutrinos fields

$$\nu_{eL} = \frac{1+\gamma_5}{2}\nu_e, \qquad \nu_{\mu L} = \frac{1+\gamma_5}{2}\nu_\mu \tag{1.14}$$

and right-handed components of the antineutrinos fields

$$\nu^C_{eR}=\frac{1-\gamma_5}{2}\nu^C_e=\left(\nu_{eL}\right)^C, \qquad \nu^C_{\mu R}=\frac{1-\gamma_5}{2}\nu^C_\mu=\left(\nu_{\mu L}\right)^C \tag{1.15}$$

can appear in the hamiltonian.

Here

$$\nu^C_{e,\mu}=C\bar{\nu}_{e,\mu} \tag{1.16}$$

is the charge conjugated spinor. The matrix C satisfies the following relations:

$$\begin{array}{rcl} C^\dagger C &=& 1 \\ C\gamma^T_\alpha C^{-1} &=& -\gamma_\alpha \\ C^T &=& -C \end{array} \tag{1.17}$$

The hamiltonian of the interaction which does not conserve the lepton numbers, quadratic in the neutrino fields, has the following general form

$$H=m_{\bar{e}e}\bar{\nu}^C_{eR}\nu_{eL}+m_{\bar{\mu}\mu}\bar{\nu}^C_{\mu R}\nu_{\mu L}+m_{\bar{\mu}e}\left(\bar{\nu}^C_{\mu R}\nu_{eL}+\bar{\nu}^C_{eR}\nu_{\mu L}\right)+h.c. \tag{1.18}$$

where the parameters $m_{\bar{e}e}$, $m_{\bar{\mu}\mu}$, $m_{\bar{\mu}e}$ have the dimensions of a mass.

The hamiltonian Eq.(1.18) can be written more compactly

$$H=\bar{\nu}^C_R M\nu_L+\bar{\nu}_L M^\dagger\nu^C_R, \tag{1.19}$$

where

$$\nu_L=\begin{pmatrix}\nu_{eL}\\ \nu_{\mu L}\end{pmatrix}, \qquad \nu^C_R=\begin{pmatrix}\nu^C_{eR}\\ \nu^C_{\mu R}\end{pmatrix}, \qquad M=\begin{pmatrix}m_{\bar{e}e} & m_{\bar{\mu}e}\\ m_{\bar{\mu}e} & m_{\bar{\mu}\mu}\end{pmatrix}. \tag{1.20}$$

If the interaction Eq.(1.18) is invariant under the CP-transformation, the parameters $m_{\bar{e}e}$, $m_{\bar{\mu}\mu}$, $m_{\bar{\mu}e}$ are real and

$$M^\dagger=M, \qquad M^T=M\Longrightarrow M^*=M. \tag{1.21}$$

In such a case the interaction hamiltonian can be expressed as follows:

$$H=\bar{\nu}^C_R M\left(\nu_L+\nu^C_R\right)+\bar{\nu}_L M\left(\nu^C_R+\nu_L\right) \tag{1.22}$$

To demonstrate Eq.(1.22) we note that the following equality holds

$$H=\bar{\nu}^C_R M\nu_L+\bar{\nu}_L M\nu^C_R=\bar{\nu}^C_R M\left(\nu_L+\nu^C_R\right)+\bar{\nu}_L M\left(\nu^C_R+\nu_L\right). \tag{1.23}$$

Indeed, being

$$\nu_L=\frac{1}{2}\left(1+\gamma_5\right)\nu, \qquad \bar{\nu}_L=\frac{1}{2}\bar{\nu}\left(1-\gamma_5\right), \tag{1.24}$$

we have

$$\bar{\nu}_L \nu_L = \frac{\bar{\nu}}{4} \left(1-\gamma_5\right)\left(1+\gamma_5\right)\nu = 0. \tag{1.25}$$

Likewise

$$\bar{\nu}_R^C \nu_R^C = 0. \tag{1.26}$$

In fact

$$\left(1-\gamma_5\right)\left(1+\gamma_5\right) = 0, \tag{1.27}$$

since

$$\gamma_5 = \left(\begin{array}{cccc} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{array}\right) \Longrightarrow \gamma_5\ \gamma_5 = \left(\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right). \tag{1.28}$$

Let we put

$$\chi = \nu_L + \nu_R^C = \left(\begin{array}{c} \nu_{eL} + \nu_{eR}^C \\ \nu_{\mu L} + \nu_{\mu R}^C \end{array}\right) = \left(\begin{array}{c} \chi_e \\ \chi_\mu \end{array}\right), \tag{1.29}$$

with χ_e, χ_μ fields of Majorana neutrinos, the Hamiltonian Eq.(1.22), then can be written as

$$H = \bar{\chi} M \chi, \tag{1.30}$$

moreover we have

$$\chi^C = C\bar{\chi} = \chi. \tag{1.31}$$

The matrix M in the Eq.(1.30) can be diagonalized by using the orthogonal matrix U $(U^T U = 1)$:

$$U = \left(\begin{array}{cc} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{array}\right). \tag{1.32}$$

We have

$$M = U M_0 U^{-1} \tag{1.33}$$

with

$$M_0 = \left(\begin{array}{cc} m_1 & 0 \\ 0 & m_2 \end{array} \right). \tag{1.34}$$

The expressions relating the masses m_1, m_2 and the mixing angle θ to the values $m_{\bar{e}e}$, $m_{\bar{\mu}\mu}$, $m_{\bar{\mu}e}$, are obtained by Eq.(1.33):

$$\left(\begin{array}{cc} m_{\bar{e}e} & m_{\bar{\mu}e} \\ m_{\bar{\mu}e} & m_{\bar{\mu}\mu} \end{array} \right) = \left(\begin{array}{cc} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{array} \right) \left(\begin{array}{cc} m_1 & 0 \\ 0 & m_2 \end{array} \right) \left(\begin{array}{cc} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{array} \right) \tag{1.35}$$

and

$$\begin{aligned} m_{\bar{e}e} &= m_1 \cos^2\theta + m_2 \; \sin^2\theta & (1.36) \\ m_{\bar{\mu}\mu} &= m_1 \sin^2\theta + m_2 \cos^2\theta & (1.37) \\ m_{\bar{\mu}e} &= (-m_1 + m_2) \sin\theta \cos\theta. & (1.38) \end{aligned}$$

Then

$$\begin{aligned} tg2\theta &= \frac{2m_{\bar{\mu}e}}{m_{\bar{\mu}\mu} - m_{\bar{e}e}} & (1.39) \\ m_{1,2} &= \frac{1}{2}\left[m_{\bar{e}e} + m_{\bar{\mu}\mu} \pm \sqrt{(m_{\bar{e}e} - m_{\bar{\mu}\mu})^2 + 4m_{\bar{\mu}e}^2} \right]. & (1.40) \end{aligned}$$

The oscillations take place if both $\theta \neq 0$ and $m_1 \neq m_2$.

Moreover for $\theta = \frac{\pi}{4}$ we have the maximum mixing and then $m_{\bar{e}e} = m_{\bar{\mu}\mu}$ with $m_{\bar{\mu}e} \neq 0$.

Letting

$$\Phi = U^T \chi, \tag{1.41}$$

the Hamiltonian

$$H = \bar{\chi} M \chi \tag{1.42}$$

becomes

$$H = \bar{\Phi} M_0 \Phi = \sum_{\sigma=1,2} m_\sigma \bar{\Phi}_\sigma \Phi_\sigma. \tag{1.43}$$

Then Φ_1 and Φ_2 are the fields of Majorana neutrinos with masses m_1, m_2 respectively.

Being

$$\Phi = U^T \chi \Longrightarrow \chi = U\Phi, \tag{1.44}$$

then the fields $\nu_{eL}, \nu_{\mu L}$ which are present in the ordinary weak interaction hamiltonian, are connected with the fields of Majorana neutrinos by the relations

$$\nu_{eL} = \sum_{\sigma=1,2} U_{1\sigma}\Phi_{\sigma L}, \qquad \nu_{\mu L} = \sum_{\sigma=1,2} U_{2\sigma}\Phi_{\sigma L}. \tag{1.45}$$

Thus, in the usual weak interaction hamiltonian there appear orthogonal superpositions of the fields of Majorana neutrinos, the mass of which m_1, m_2 are not equal to zero and in such a case there arise the neutrino oscillations. Making use of the expressions

$$\Phi = U^T\chi \qquad \text{and} \qquad U = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}, \tag{1.46}$$

we have

$$\begin{aligned} \Phi_1 &= \chi_1\cos\theta - \chi_2\sin\theta, \\ \Phi_2 &= \chi_1\sin\theta + \chi_2\cos\theta, \end{aligned} \tag{1.47}$$

and

$$\begin{aligned} \nu_{eL} &= \Phi_{1L}\cos\theta + \Phi_{2L}\sin\theta, \\ \nu_{\mu L} &= -\Phi_{1L}\sin\theta + \Phi_{2L}\cos\theta. \end{aligned} \tag{1.48}$$

From these expressions it is clear that the angle θ characterizes the degree of mixing of the Majorana fields Φ_1 and Φ_2.

1.4 Dirac neutrinos

This theory is based on the analogy between leptons and quarks [1].

The total charged hadronic current is

$$\left(J^h_\alpha\right) = \left(J^h_\alpha\right)_C + \left(J^h_\alpha\right)_{GIM}, \tag{1.49}$$

where $\left(J^h_\alpha\right)_C$ is the hadron Cabibbo current and $\left(J^h_\alpha\right)_{GIM}$ is the hadron current of the G.I.M. model.

In particular: $\left(J^h_\alpha\right)_C$ is given by the expression

$$\left(J^h_\alpha\right)_C = \bar{u}_L\gamma_\alpha d'_L \tag{1.50}$$

where

$$d' = d\cos\theta_c + s\sin\theta_c, \tag{1.51}$$

with θ_c Cabibbo angle, u, d, s field operators of the u-quarks $\big(Q = \frac{2}{3},\ T_3 = \frac{1}{2},\ S = 0\big)$, d-quarks $\big(Q = -\frac{1}{3},\ T_3 = -\frac{1}{2},\ S = 0\big)$, s-quarks $\big(Q = -\frac{1}{3},\ T = 0,\ S = -1\big)$ respectively. Moreover, the hadron current of the G.I.M. model $\big(J^h_\alpha\big)_{GIM}$ is

$$\big(J^h_\alpha\big)_{GIM} = \bar{c}_L \gamma_\alpha s'_L \tag{1.52}$$

where

$$s' = -d \sin\theta_c + s \cos\theta_c, \tag{1.53}$$

c field operator of the c quark $\big(Q = \frac{2}{3},\ T = 0,\ S = 0,\ C = 1\big)$.

The leptonic current has the same structure as the charged hadron current:

$$J_\alpha = (\bar{\nu}_{eL} \gamma_\alpha e_L) + (\bar{\nu}_{\mu L} \gamma_\alpha \mu_L)\,. \tag{1.54}$$

Only the left-handed field components are present in both currents.

Comparing the hadron current with the lepton current Eq.(1.54), we see an important difference between them: whereas orthogonal superpositions of the d and s quark fields are present in the hadronic current, in the lepton current the electron neutrino and muon neutrino fields appear unmixed, (the charged hadron current does not conserve strangeness, while the lepton current is conserving the lepton numbers).

In order to remove this difference, let us assume that there exist two neutrinos (ν_1, ν_2) with finite masses (m_1, m_2) and that the operators ν_e, ν_μ are orthogonal combinations of ν_1, ν_2

$$\begin{aligned} \nu_e &= \nu_1 \cos\theta + \nu_2 \sin\theta \\ \nu_\mu &= -\nu_1 \sin\theta + \nu_2 \cos\theta \end{aligned} \tag{1.55}$$

where ν_1, ν_2 are the fields of Dirac neutrinos with masses m_1 and m_2, θ is the mixing angle. Let us note that θ is different to the Cabibbo angle θ_c.

Two values of the mixing angle are of special significance, $\theta = 0$ and $\theta = \frac{\pi}{4}$. The case $\theta = 0$ (no mixing) corresponds to the theory with strict conservation of the electron and muon lepton number. The case $\theta = \frac{\pi}{4}$ corresponds to maximum amplitude of oscillations. Thus in the Dirac theory there is a full analogy between lepton and quarks weak currents.

In this scheme there is no lepton number distinguishing the two types of neutrinos ν_1 and ν_2; the neutrinos ν_1 and ν_2 differ only in their mass values (just as the d and s quarks do).

The mass term of the hamiltonian in the Dirac theory has the form

$$H_1 = m_1 \bar{\nu}_1 \nu_1 + m_2 \bar{\nu}_2 \nu_2. \tag{1.56}$$

Let us express ν_1 and ν_2 through ν_e and ν_μ. We get

$$H_1 = m_{ee} \bar{\nu}_e \nu_e + m_{\mu\mu} \bar{\nu}_\mu \nu_\mu + m_{\mu e} (\bar{\nu}_\mu \nu_e + \bar{\nu}_e \nu_\mu) \tag{1.57}$$

with

$$\begin{aligned} m_{ee} &= m_1 \cos^2\theta + m_2 \sin^2\theta && (1.58)\\ m_{\mu\mu} &= m_1 \sin^2\theta + m_2 \cos^2\theta && (1.59)\\ m_{\mu e} &= (-m_1 + m_2)\sin\theta\cos\theta. && (1.60) \end{aligned}$$

Where m_{ee} and $m_{\mu\mu}$ are the bare masses of electron and muon neutrinos.

The term

$$H = m_{\mu e}(\bar{\nu}_\mu \nu_e + \bar{\nu}_e \nu_\mu), \qquad (1.61)$$

in H_1, does not conserve separately L_e and L_μ, but conserve $L_e + L_\mu$; then H is a super-weak interaction that does not conserve the leptonic number.

The masses m_1, m_2 and the mixing angle θ are related to the values m_{ee}, $m_{\mu\mu}$, $m_{\mu e}$, by

$$\begin{aligned} tg2\theta &= \frac{2m_{\mu e}}{m_{\mu\mu} - m_{ee}} && (1.62)\\ m_{1,2} &= \frac{1}{2}\left[m_{ee} + m_{\mu\mu} \pm \sqrt{(m_{ee} - m_{\mu\mu})^2 + 4m_{\mu e}^2}\right] && (1.63) \end{aligned}$$

These results are to be compared with the corresponding ones for the Majorana case in Section 1.3.

We have been considering the simplest theories with mixing of two neutrinos with finite masses.

In both the theories of Majorana and of Dirac, the neutrino masses are not equal to zero and the operators of neutrino fields are present in the hamiltonian in the form of orthogonal combinations.

The hamiltonian of the Majorana theory does not conserve the muon L_μ and the electron L_e lepton numbers. The hamiltonian of the Dirac theory is conserving the sum $L_e + L_\mu$. The neutrino oscillations are described by identical expressions in both theories. Contrarily to the Dirac theory, the Majorana theory allows in principle the existence of neutrino-less double beta decay and of other processes in which $L_e + L_\mu$ is not conserved (see Appendix A).

In the Majorana theory every type of neutrino is associated with two states; in the Dirac theory there are four states for every neutrino type. In this sense, in the Majorana theory there is no analogy between leptons and quarks.

1.5 Neutrino oscillations

The theories we have considered above lead to neutrino oscillations [1]. Let us denote by $|\nu_e\rangle$, and $|\nu_\mu\rangle$ the state vectors of the electron and muon neutrinos with momentum

$\mathbf{p}$ and helicity equal to -1. From

$$\begin{aligned} \nu_{eL} &= \sum_{\sigma=1,2} U_{1\sigma}\Phi_{\sigma L}, \qquad & \nu_{\mu L} = \sum_{\sigma=1,2} U_{2\sigma}\Phi_{\sigma L} \\ \nu_e &= \nu_1 \cos\theta + \nu_2 \sin\theta, \qquad & \nu_\mu = -\nu_1 \sin\theta + \nu_2 \cos\theta, \end{aligned}$$

it follows that

$$|\nu_l\rangle = \sum_{\sigma=1,2} U_{l\sigma}|\nu_\sigma\rangle \tag{1.64}$$

with $l = e,\ \mu$.

Here $|\nu_\sigma\rangle$, with $\sigma = 1, 2$, is the state vector of the neutrino with mass m_σ, momentum $\mathbf{p}$, helicity -1. The orthogonal matrix U has the form of the Eq.(1.32).

The vectors $|\nu_\sigma\rangle$ describe both Majorana neutrinos and Dirac neutrinos.

We have

$$H|\nu_\sigma\rangle = E|\nu_\sigma\rangle \tag{1.65}$$

where H is the total hamiltonian and $E_\sigma = \sqrt{m_\sigma^2 + p^2}$.

We also have

$$|\nu_\sigma\rangle = \sum_{l=e,\mu} U_{l\sigma}|\nu_l\rangle. \tag{1.66}$$

Let us consider the behavior of the beam of neutrinos, at initial time ($t = 0$) such a beam is described by the vector $|\nu_l\rangle$. At time t the state vector of the beam is given by the expression

$$|\nu_l(t)\rangle = e^{-iHt}|\nu_l\rangle = \sum_{\sigma=1,2} U_{l\sigma} e^{-iE_\sigma t}|\nu_\sigma\rangle. \tag{1.67}$$

Then the neutrino beam is not described by a stationary state, but by a superpositions of stationary states. This happens because $|\nu_l\rangle$ is not an eigenstate of the hamiltonian H.

If we denote with

$$a_{\nu_{l'};\nu_l}(t) = \sum_{\sigma=1,2} U_{l\sigma} e^{-iE_\sigma t} U_{l'\sigma} \tag{1.68}$$

the probability amplitude of finding $\nu_{l'}$ at a time t after the generation of ν_l, then we can expand the state vector Eq.(1.67) in terms of vectors $|\nu_{l'}\rangle$

$$|\nu_l(t)\rangle = \sum_{l'=e,\mu} a_{\nu_{l'};\nu_l}(t)|\nu_{l'}\rangle. \tag{1.69}$$

We have also

$$a_{\nu_{l'};\nu_l}(0) = \sum_{\sigma=1,2} U_{l\sigma} U_{l'\sigma} = \delta_{l'l}. \tag{1.70}$$

Clearly in the case $m_1 \neq m_2$ and $U_{l\sigma} \neq \delta_{l\sigma}$, we have $a_{\nu_e;\nu_\mu}(t) = a_{\nu_\mu;\nu e}(t) \neq 0$ that is there arise oscillations $\nu_e \rightleftarrows \nu_\mu$.

The probability of transitions $\nu_l \rightleftarrows \nu_{l'}$ is given by the expression

$$P_{\nu_{l'} \to \nu_l}(t) = P_{\nu_l \to \nu_{l'}}(t) = \sum_{\sigma,\sigma'} U_{l\sigma} U_{l'\sigma} U_{l\sigma'} U_{l'\sigma'} \cos\left(E_\sigma - E_{\sigma'}\right) t. \tag{1.71}$$

It easy to see that the probabilities $P_{\nu_{l'} \to \nu_l}(t)$ satisfy the relation

$$\sum_{l'=e,\mu} P_{\nu_{l'} \to \nu_l}(t) = 1. \tag{1.72}$$

In the case $p \gg m_1$, m_2 , which is of interest, we have

$$\begin{aligned} E_i &= \sqrt{p^2c^2 + m_i^2c^4} \approxeq pc\left(1 + \frac{m_i^2c^2}{2p^2}\right) = pc + \frac{m_i^2c^3}{2p}\ , \qquad \text{and setting} \quad c = 1 : \\ E_i &= p + \frac{m_i^2}{2p} \Longrightarrow E_1 - E_2 = \frac{m_1^2 - m_2^2}{2p}. \end{aligned} \tag{1.73}$$

Making use of expressions of U and $P_{\nu_{l'} \to \nu_l}(t)$, and indicating with

$$L = \frac{4\pi p}{|m_1^2 - m_2^2|} \tag{1.74}$$

the oscillation length, the probability of finding $\nu_{l'}$ at a distance R from a source of ν_l is

$$P_{\nu_e \to \nu e}(R) = P_{\nu_\mu \to \nu_\mu}(R) = 1 - \frac{1}{2}\sin^2 2\theta \left(1 - \cos\frac{2\pi R}{L}\right), \tag{1.75}$$

$$P_{\nu_e \to \nu_\mu}(R) = P_{\nu_\mu \to \nu_e}(R) = \frac{1}{2}\sin^2 2\theta \left(1 - \cos\frac{2\pi R}{L}\right). \tag{1.76}$$

Which can be expressed also as [1]

$$P_{\nu_e \to \nu e}(t) = P_{\nu_\mu \to \nu_\mu}(t) = 1 - \sin^2 2\theta\ \sin^2\left(\frac{\Delta\omega}{2}t\right), \tag{1.77}$$

$$P_{\nu_e \to \nu_\mu}(t) = P_{\nu_\mu \to \nu_e}(t) = \sin^2 2\theta\ \sin^2\left(\frac{\Delta\omega}{2}t\right), \tag{1.78}$$

where $\Delta\omega = \omega_1 - \omega_2$, is the energy difference of the mass components. The probability conservation is satisfied

$$P_{\nu_e \to \nu_e}(t) + P_{\nu_e \to \nu_\mu}(t) = 1. \tag{1.79}$$

The Eqs.(1.77), (1.78) are the Bilenky-Pontecorvo formulas for neutrino oscillations.

Chapter 2

Quantum field theory of fermion mixing

The fermion mixing transformations are studied in the quantum field theory framework. In particular neutrino mixing is considered and the Fock space of definite flavor states is shown to be unitarily inequivalent to the Fock space of definite mass states. We study the structure of the currents and charges for the mixed fermion fields and the flavor oscillation formula is computed for two flavors mixing, the oscillation amplitude is found to be momentum dependent. The flavor vacuum state exhibits the structure of $SU(2)$ generalized coherent state.

2.1 Introduction

The purpose of the present chapter is the study of the quantum field theory (QFT) framework of the fermion mixing transformations, thus focusing the attention on the theoretical structure of fermion mixing.

We will consider, in particular, neutrino mixing transformations and this analysis will lead to some modifications of the quantum mechanics neutrino oscillation formulas.

The chapter is organized as follows. In Section 2.2 we study the generator of the Pontecorvo neutrino mixing transformations (two flavors mixing for Dirac fields). We show that in the Lehmann-Symanzik-Zimmermann (LSZ) formalism of quantum field theory [22, 23] the Fock space of the flavor states is unitarily inequivalent to the Fock space of the mass eigenstates in the infinite volume limit [25]. The flavor states are obtained as condensate of massive neutrino pairs and exhibit the structure of $SU(2)$ coherent states [44]. In Section 2.3 we generalize the mixing transformations, indeed we expand the flavor fields in a basis with arbitrary masses [28, 29]. In Section 2.4

we analyze the structure of currents for mixed fermion fields [19] and in Section 2.5 we derive the neutrino flavor oscillation formulae [20, 21]. Finally, the Section 2.6 is devoted to the conclusions. We confine mathematical details to the Appendices B, C, D.

2.2 The vacuum structure for fermion mixing

We consider the Pontecorvo mixing relations [1], although the following discussion applies to any Dirac fields.

The mixing relations are

$$\begin{aligned} \nu_e(x) &= \nu_1(x)\cos\theta + \nu_2(x)\sin\theta \\ \nu_\mu(x) &= -\nu_1(x)\sin\theta + \nu_2(x)\cos\theta \end{aligned} \tag{2.1}$$

where $\nu_e(x)$ and $\nu_\mu(x)$ are the Dirac neutrino fields with definite flavors. $\nu_1(x)$ and $\nu_2(x)$ are the free neutrino fields with definite masses m_1 and m_2, respectively. The fields $\nu_1(x)$ and $\nu_2(x)$ are written as

$$\nu_i(x) = \frac{1}{\sqrt{V}}\sum_{\mathbf{k},r}\left[u^r_{\mathbf{k},i}(t)\alpha^r_{\mathbf{k},i}(t) + v^r_{-\mathbf{k},i}(t)\beta^{r\dagger}_{-\mathbf{k},i}(t)\right]e^{i\mathbf{k}\cdot\mathbf{x}}, \qquad i = 1,2 \tag{2.2}$$

with

$$u^r_{\mathbf{k},i}(t) = e^{-i\omega_{k,i}t}u^r_{\mathbf{k},i} \qquad \text{and} \qquad v^r_{\mathbf{k},i}(t) = e^{i\omega_{k,i}t}v^r_{\mathbf{k},i}, \tag{2.3}$$

$$u^1_{\mathbf{k},i} = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}}\begin{pmatrix} 1 \\ 0 \\ \frac{k_3}{\omega_{k,i}+m_i} \\ \frac{k_1+ik_2}{\omega_{k,i}+m_i} \end{pmatrix}; \qquad u^2_{\mathbf{k},i} = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}}\begin{pmatrix} 0 \\ 1 \\ \frac{k_1-ik_2}{\omega_{k,i}+m_i} \\ \frac{-k_3}{\omega_{k,i}+m_i} \end{pmatrix}$$

$$v^1_{-\mathbf{k},i} = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}}\begin{pmatrix} \frac{-k_3}{\omega_{k,i}+m_i} \\ \frac{-k_1-ik_2}{\omega_{k,i}+m_i} \\ 1 \\ 0 \end{pmatrix}; \qquad v^2_{-\mathbf{k},i} = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}}\begin{pmatrix} \frac{-k_1+ik_2}{\omega_{k,i}+m_i} \\ \frac{k_3}{\omega_{k,i}+m_i} \\ 0 \\ 1 \end{pmatrix}$$

and

$$\omega_{k,i} = \sqrt{\mathbf{k}^2 + m_i^2}. \tag{2.4}$$

The operator $\alpha^r_{\mathbf{k},i}$ and $\beta^r_{\mathbf{k},i}$, $i = 1,2$, $r = 1,2$ are the annihilator operators for the vacuum state $|0\rangle_{1,2} \equiv |0\rangle_1 \otimes |0\rangle_2$: $\alpha^r_{\mathbf{k},i}|0\rangle_{12} = \beta^r_{\mathbf{k},i}|0\rangle_{12} = 0$.

The anticommutation relations are:

$$\left\{\nu_i^{\alpha}(x), \nu_j^{\beta\dagger}(y)\right\}_{t=t'} = \delta^3(\mathbf{x}-\mathbf{y})\delta_{\alpha\beta}\delta_{ij}, \qquad \alpha, \beta = 1, ...4, \tag{2.5}$$

$$\left\{\alpha^r_{\mathbf{k},i}, \alpha^{s\dagger}_{\mathbf{q},j}\right\} = \delta_{\mathbf{kq}}\delta_{rs}\delta_{ij}; \qquad \left\{\beta^r_{\mathbf{k},i}, \beta^{s\dagger}_{\mathbf{q},j}\right\} = \delta_{\mathbf{kq}}\delta_{rs}\delta_{ij}, \qquad i, j = 1, 2. \tag{2.6}$$

All other anticommutators are zero. The orthonormality and completeness relations are

$$\begin{aligned} u^{r\dagger}_{\mathbf{k},i} u^s_{\mathbf{k},i} = v^{r\dagger}_{\mathbf{k},i} v^s_{\mathbf{k},i} &= \delta_{rs}, & (2.7)\\ u^{r\dagger}_{\mathbf{k},i} v^s_{-\mathbf{k},i} = v^{r\dagger}_{-\mathbf{k},i} u^s_{\mathbf{k},i} &= 0, & (2.8)\\ \sum_r (u^r_{\mathbf{k},i} u^{r\dagger}_{\mathbf{k},i} + v^r_{-\mathbf{k},i} v^{r\dagger}_{-\mathbf{k},i}) &= 1. & (2.9) \end{aligned}$$

The Eqs.(2.1) relate the respective hamiltonians $H_{1,2}$ (we consider only the mass terms) and $H_{e,\mu}$ [1]:

$$H_{1,2} = m_1 \, \nu_1^\dagger \nu_1 + m_2 \, \nu_2^\dagger \nu_2 \tag{2.10}$$

$$H_{e,\mu} = m_{ee} \, \nu_e^\dagger \nu_e + m_{\mu\mu} \, \nu_\mu^\dagger \nu_\mu + m_{e\mu} \left(\nu_e^\dagger \nu_\mu + \nu_\mu^\dagger \nu_e\right) \tag{2.11}$$

where

$$\begin{aligned} m_{ee} &= m_1 \cos^2\theta + m_2 \sin^2\theta, & (2.12)\\ m_{\mu\mu} &= m_1 \sin^2\theta + m_2 \cos^2\theta, & (2.13)\\ m_{e\mu} &= (m_2 - m_1)\sin\theta\cos\theta. & (2.14) \end{aligned}$$

In QFT the basic dynamics, i.e. the Lagrangian and the resulting field equations, is given in terms of Heisenberg (or interacting) fields. The physical observables are expressed in terms of asymptotic in- (or out-) fields, also called physical or free fields. In the LSZ formalism of QFT [22, 23], the free fields, say for definitiveness the in-fields, are obtained by the weak limit of the Heisenberg fields for time $t \to -\infty$. The meaning of the weak limit is that the realization of the basic dynamics in terms of the in-fields is not unique so that the limit for $t \to -\infty$ (or $t \to +\infty$ for the out-fields) is representation dependent. The representation dependence of the asymptotic limit arises from the existence in QFT of infinitely many unitarily non-equivalent representations of the canonical (anti-)commutation relations [22, 23]. Of course, since observables are described in terms of asymptotic fields, unitarily inequivalent representations describe different, i.e. physically inequivalent, phases. It is therefore of crucial importance, in order to get physically meaningful results, to investigate with much care the mapping among Heisenberg or interacting fields and free fields. Such a mapping is usually called the Haag expansion or the dynamical map [22, 23]. Only in a very rude and naive approximation we may assume that interacting fields and free fields share the same vacuum state and the same Fock space representation.

We stress that the above remarks apply to QFT, namely to systems with infinite number of degrees of freedom. In quantum mechanics, where finite volume systems are considered, the von Neumann theorem ensures that the representations of the canonical commutation relations are each other unitary equivalent and no problem arises with uniqueness of the asymptotic limit. In QFT, however, the von Neumann theorem does not hold and much more careful attention is required when considering any mapping among interacting and free fields [22, 23].

With this warnings, mixing relations such as the relations Eqs.(2.1) deserve a careful analysis. In fact we will investigate the structure of the Fock spaces $\mathcal{H}_{1,2}$ and $\mathcal{H}_{e,\mu}$ relative to $\nu_1(x)$, $\nu_2(x)$ and $\nu_e(x)$, $\nu_\mu(x)$, respectively. In particular we will study the relation among these spaces in the infinite volume limit. We expect that $\mathcal{H}_{1,2}$ and $\mathcal{H}_{e,\mu}$ become orthogonal in such a limit, since they represent the Hilbert spaces for free and interacting fields, respectively [22, 23]. In the following, we will perform all computations at finite volume V and only at the end we will put $V \to \infty$.

We construct the generator for the mixing transformation Eqs.(2.1) and define [25]:

$$\begin{aligned} \nu_e^\alpha(x) &= G_\theta^{-1}(t)\ \nu_1^\alpha(x)\ G_\theta(t) \\ \nu_\mu^\alpha(x) &= G_\theta^{-1}(t)\ \nu_2^\alpha(x)\ G_\theta(t) \end{aligned} \tag{2.15}$$

where $G_\theta(t)$ is given by

$$G_\theta(t) = exp\left[\theta \int d^3\mathbf{x} \left(\nu_1^\dagger(x)\nu_2(x) - \nu_2^\dagger(x)\nu_1(x)\right)\right] \ , \tag{2.16}$$

and is, at finite volume, an unitary operator: $G_\theta^{-1}(t) = G_{-\theta}(t) = G_\theta^\dagger(t)$, preserving the canonical anticommutation relations Eqs.(2.5). To obtain the Eq.(2.16), we observe that, from Eqs.(2.15),

$$d^2\nu_e^\alpha/d\theta^2 = -\nu_e^\alpha \ , \quad d^2\nu_\mu^\alpha/d\theta^2 = -\nu_\mu^\alpha, \tag{2.17}$$

by using the initial conditions

$$\nu_e^\alpha|_{\theta=0} = \nu_1^\alpha, \quad d\nu_e^\alpha/d\theta|_{\theta=0} = \nu_2^\alpha \quad \text{and} \quad \nu_\mu^\alpha|_{\theta=0} = \nu_2^\alpha, \quad d\nu_\mu^\alpha/d\theta|_{\theta=0} = -\nu_1^\alpha, \tag{2.18}$$

the operator $G_\theta(t)$ generates Eqs.(2.1).

By introducing the operators

$$S_+(t) \equiv \int d^3\mathbf{x}\ \nu_1^\dagger(x)\nu_2(x) \ , \quad S_-(t) \equiv \int d^3\mathbf{x}\ \nu_2^\dagger(x)\nu_1(x) \ = (S_+)^\dagger \ , \tag{2.19}$$

$G_\theta(t)$ can be written as

$$G_\theta(t) = exp[\theta(S_+ - S_-)] \ . \tag{2.20}$$

Introducing S_3 and the Casimir operator S_0 (proportional to the total charge) as follows

$$S_3 \equiv \frac{1}{2}\int d^3\mathbf{x} \left(\nu_1^\dagger(x)\nu_1(x) - \nu_2^\dagger(x)\nu_2(x)\right) \ , \tag{2.21}$$

$$S_0 \equiv \frac{1}{2}\int d^3\mathbf{x} \left(\nu_1^\dagger(x)\nu_1(x) + \nu_2^\dagger(x)\nu_2(x)\right) \ , \tag{2.22}$$

the $su(2)$ algebra is closed:

$$[S_+(t), S_-(t)] = 2S_3 \quad , \quad [S_3, S_\pm(t)] = \pm S_\pm(t) \quad , \quad [S_0, S_3] = [S_0, S_\pm(t)] = 0 \ . \tag{2.23}$$

The momentum expansion of $S_+(t)$, $S_-(t)$, S_3, and S_0, is given by using the Eq.(2.2):

$$\begin{aligned} S_+(t) \ &\equiv \ \sum_{\mathbf{k}} S_+^{\mathbf{k}}(t) = \sum_{\mathbf{k}}\sum_{r,s} \Big(u^{r\dagger}_{\mathbf{k},1}(t) u^s_{\mathbf{k},2}(t) \alpha^{r\dagger}_{\mathbf{k},1}\alpha^s_{\mathbf{k},2} + v^{r\dagger}_{-\mathbf{k},1}(t) u^s_{\mathbf{k},2}(t) \beta^r_{-\mathbf{k},1}\alpha^s_{\mathbf{k},2} + \\ &\quad + u^{r\dagger}_{\mathbf{k},1}(t) v^s_{-\mathbf{k},2}(t) \alpha^{r\dagger}_{\mathbf{k},1}\beta^{s\dagger}_{-\mathbf{k},2} + v^{r\dagger}_{-\mathbf{k},1}(t) v^s_{-\mathbf{k},2}(t) \beta^r_{-\mathbf{k},1}\beta^{s\dagger}_{-\mathbf{k},2} \Big) \ , \end{aligned} \tag{2.24}$$

$$\begin{aligned} S_-(t) \ &\equiv \ \sum_{\mathbf{k}} S_-^{\mathbf{k}}(t) = \sum_{\mathbf{k}}\sum_{r,s} \Big(u^{r\dagger}_{\mathbf{k},2}(t) u^s_{\mathbf{k},1}(t) \alpha^{r\dagger}_{\mathbf{k},2}\alpha^s_{\mathbf{k},1} + v^{r\dagger}_{-\mathbf{k},2}(t) u^s_{\mathbf{k},1}(t) \beta^r_{-\mathbf{k},2}\alpha^s_{\mathbf{k},1} + \\ &\quad + u^{r\dagger}_{\mathbf{k},2}(t) v^s_{-\mathbf{k},1}(t) \alpha^{r\dagger}_{\mathbf{k},2}\beta^{s\dagger}_{-\mathbf{k},1} + v^{r\dagger}_{-\mathbf{k},2}(t) v^s_{-\mathbf{k},1}(t) \beta^r_{-\mathbf{k},2}\beta^{s\dagger}_{-\mathbf{k},1} \Big) \ , \end{aligned} \tag{2.25}$$

$$S_3 \equiv \sum_{\mathbf{k}} S_3^{\mathbf{k}} = \frac{1}{2}\sum_{\mathbf{k},r} \left(\alpha^{r\dagger}_{\mathbf{k},1}\alpha^r_{\mathbf{k},1} - \beta^{r\dagger}_{-\mathbf{k},1}\beta^r_{-\mathbf{k},1} - \alpha^{r\dagger}_{\mathbf{k},2}\alpha^r_{\mathbf{k},2} + \beta^{r\dagger}_{-\mathbf{k},2}\beta^r_{-\mathbf{k},2} \right) , \tag{2.26}$$

$$S_0 \equiv \sum_{\mathbf{k}} S_0^{\mathbf{k}} = \frac{1}{2}\sum_{\mathbf{k},r} \left(\alpha^{r\dagger}_{\mathbf{k},1}\alpha^r_{\mathbf{k},1} - \beta^{r\dagger}_{-\mathbf{k},1}\beta^r_{-\mathbf{k},1} + \alpha^{r\dagger}_{\mathbf{k},2}\alpha^r_{\mathbf{k},2} - \beta^{r\dagger}_{-\mathbf{k},2}\beta^r_{-\mathbf{k},2} \right) . \tag{2.27}$$

We observe that the operatorial structure of Eqs.(2.24) and (2.25) is the one of the rotation generator and of the Bogoliubov generator. Using these expansions it is easy to show that the $su(2)$ algebra does hold for each momentum component:

$$\left[S_+^{\mathbf{k}}(t), S_-^{\mathbf{k}}(t)\right] = 2S_3^{\mathbf{k}}, \qquad \left[S_3^{\mathbf{k}}(t), S_\pm^{\mathbf{k}}(t)\right] = \pm S_\pm^{\mathbf{k}}(t), \qquad \left[S_0^{\mathbf{k}}, S_3^{\mathbf{k}}\right] = \left[S_0^{\mathbf{k}}, S_\pm^{\mathbf{k}}\right] = 0, \tag{2.28}$$

$$\left[S_{\pm}^{\mathbf{k}}(t), S_{\pm}^{\mathbf{p}}(t)\right] = \left[S_{3}^{\mathbf{k}}(t), S_{\pm}^{\mathbf{p}}(t)\right] = \left[S_{3}^{\mathbf{k}}, S_{3}^{\mathbf{p}}\right] = 0, \qquad \mathbf{k} \neq \mathbf{p}. \tag{2.29}$$

This means that the original $su(2)$ algebra given in Eqs.(2.23) splits into $\mathbf{k}$ disjoint $su_{\mathbf{k}}(2)$ algebras, given by Eqs.(2.28), i.e. we have the group structure $\bigotimes_{\mathbf{k}} SU_{\mathbf{k}}(2)$.

To establish the relation between the Hilbert spaces for free fields $\mathcal{H}_{1,2}$ and interacting fields $\mathcal{H}_{e,\mu}$ we consider the generic matrix element ${}_{1,2}\langle a|\nu_1^{\alpha}(x)|b\rangle_{1,2}$ (a similar argument holds for $\nu_2^{\alpha}(x)$), where $|a\rangle_{1,2}$ is the generic element of $\mathcal{H}_{1,2}$. Using the inverse of the first of the Eqs.(2.15), we obtain:

$${}_{1,2}\langle a|G_{\theta}(t)\; \nu_e^{\alpha}(x)\; G_{\theta}^{-1}(t)|b\rangle_{1,2} \;=\; {}_{1,2}\langle a|\nu_1^{\alpha}(x)|b\rangle_{1,2}\ . \tag{2.30}$$

Since the operator field ν_e is defined on the Hilbert space $\mathcal{H}_{e,\mu}$, Eq.(2.30) shows that $G_{\theta}^{-1}(t)|a\rangle_{1,2}$ is a vector of $\mathcal{H}_{e,\mu}$, so $G_{\theta}^{-1}(t)$ maps $\mathcal{H}_{1,2}$ to $\mathcal{H}_{e,\mu}$:

$$G_{\theta}^{-1}(t) : \mathcal{H}_{1,2} \mapsto \mathcal{H}_{e,\mu}. \tag{2.31}$$

In particular for the vacuum $|0\rangle_{1,2}$ we have, at finite volume V:

$$|0(t)\rangle_{e,\mu} = G_{\theta}^{-1}(t)\; |0\rangle_{1,2}\ . \tag{2.32}$$

$|0\rangle_{e,\mu}$ is the vacuum for $\mathcal{H}_{e,\mu}$, which we will refer to as the flavor vacuum. Due to the linearity of $G_{\theta}(t)$, we can define the flavor annihilators, relative to the fields $\nu_e(x)$ and $\nu_{\mu}(x)$ at each time expressed as

$$\begin{aligned}
\alpha_{\mathbf{k},e}^{r}(t)\; |0(t)\rangle_{e,\mu} &= G_{\theta}^{-1}(t)\; \alpha_{\mathbf{k},1}^{r}\; |0\rangle_{1,2} = 0,\\
\alpha_{\mathbf{k},\mu}^{r}(t)\; |0(t)\rangle_{e,\mu} &= G_{\theta}^{-1}(t)\; \alpha_{\mathbf{k},2}^{r}\; |0\rangle_{1,2} = 0,\\
\beta_{\mathbf{k},e}^{r}(t)\; |0(t)\rangle_{e,\mu} &= G_{\theta}^{-1}(t)\; \beta_{\mathbf{k},1}^{r}\; |0\rangle_{1,2} = 0,\\
\beta_{\mathbf{k},\mu}^{r}(t)\; |0(t)\rangle_{e,\mu} &= G_{\theta}^{-1}(t)\; \beta_{\mathbf{k},2}^{r}\; |0\rangle_{1,2} = 0,
\end{aligned} \tag{2.33}$$

in the following way

$$\begin{aligned}
\alpha_{\mathbf{k},e}^{r}(t) &\equiv G_{\theta}^{-1}(t)\; \alpha_{\mathbf{k},1}^{r}\; G_{\theta}(t),\\
\alpha_{\mathbf{k},\mu}^{r}(t) &\equiv G_{\theta}^{-1}(t)\; \alpha_{\mathbf{k},2}^{r}\; G_{\theta}(t),\\
\beta_{\mathbf{k},e}^{r}(t) &\equiv G_{\theta}^{-1}(t)\; \beta_{\mathbf{k},1}^{r}\; G_{\theta}(t),\\
\beta_{\mathbf{k},\mu}^{r}(t) &\equiv G_{\theta}^{-1}(t)\; \beta_{\mathbf{k},2}^{r}\; G_{\theta}(t).
\end{aligned} \tag{2.34}$$

The flavor fields are then rewritten into the form:

$$\nu_e(\mathbf{x},t) \;=\; \frac{1}{\sqrt{V}}\sum_{\mathbf{k},r} e^{i\mathbf{k}.\mathbf{x}} \left[u_{\mathbf{k},1}^{r}(t)\alpha_{\mathbf{k},e}^{r}(t) + v_{-\mathbf{k},1}^{r}(t)\beta_{-\mathbf{k},e}^{r\dagger}(t)\right], \tag{2.35}$$

$$\nu_{\mu}(\mathbf{x},t) \;=\; \frac{1}{\sqrt{V}}\sum_{\mathbf{k},r} e^{i\mathbf{k}.\mathbf{x}} \left[u_{\mathbf{k},2}^{r}(t)\alpha_{\mathbf{k},\mu}^{r}(t) + v_{-\mathbf{k},2}^{r}(t)\beta_{-\mathbf{k},\mu}^{r\dagger}(t)\right], \tag{2.36}$$

i.e. they can be expanded in the same bases as ν_1 and ν_2, respectively.

The flavor annihilation operators can be calculated explicitly, we have

$$\begin{aligned}
\alpha^r_{\mathbf{k},e}(t) &= \cos\theta\;\alpha^r_{\mathbf{k},1} \;+\; \sin\theta\;\sum_s \Big[u^{r\dagger}_{\mathbf{k},1}(t)u^s_{\mathbf{k},2}(t)\;\alpha^s_{\mathbf{k},2} \;+\; u^{r\dagger}_{\mathbf{k},1}(t)v^s_{-\mathbf{k},2}(t)\;\beta^{s\dagger}_{-\mathbf{k},2}\Big] \\
\alpha^r_{\mathbf{k},\mu}(t) &= \cos\theta\;\alpha^r_{\mathbf{k},2} \;-\; \sin\theta\;\sum_s \Big[u^{r\dagger}_{\mathbf{k},2}(t)u^s_{\mathbf{k},1}(t)\;\alpha^s_{\mathbf{k},1} \;+\; u^{r\dagger}_{\mathbf{k},2}(t)v^s_{-\mathbf{k},1}(t)\;\beta^{s\dagger}_{-\mathbf{k},1}\Big] \\
\beta^r_{-\mathbf{k},e}(t) &= \cos\theta\;\beta^r_{-\mathbf{k},1} \;+\; \sin\theta\;\sum_s \Big[v^{s\dagger}_{-\mathbf{k},2}(t)v^r_{-\mathbf{k},1}(t)\;\beta^s_{-\mathbf{k},2} \;+\; u^{s\dagger}_{\mathbf{k},2}(t)v^r_{-\mathbf{k},1}(t)\;\alpha^{s\dagger}_{\mathbf{k},2}\Big] \\
\beta^r_{-\mathbf{k},\mu}(t) &= \cos\theta\;\beta^r_{-\mathbf{k},2} \;-\; \sin\theta\;\sum_s \Big[v^{s\dagger}_{-\mathbf{k},1}(t)v^r_{-\mathbf{k},2}(t)\;\beta^s_{-\mathbf{k},1} \;+\; u^{s\dagger}_{\mathbf{k},1}(t)v^r_{-\mathbf{k},2}(t)\;\alpha^{s\dagger}_{\mathbf{k},1}\Big] .
\end{aligned} \tag{2.37}$$

Without loss of generality, we can choose the reference frame such that $\mathbf{k} = (0,0,|\mathbf{k}|)$. This implies that only the products of wave functions with $r = s$ will survive, i.e. the spins decouple and the Eqs.(2.37) assume the simpler form:

$$\begin{aligned}
\alpha^r_{\mathbf{k},e}(t) &= \cos\theta\;\alpha^r_{\mathbf{k},1} \;+\; \sin\theta\;\Big(U^*_{\mathbf{k}}(t)\;\alpha^r_{\mathbf{k},2} \;+\; \epsilon^r\;V_{\mathbf{k}}(t)\;\beta^{r\dagger}_{-\mathbf{k},2}\Big) \\
\alpha^r_{\mathbf{k},\mu}(t) &= \cos\theta\;\alpha^r_{\mathbf{k},2} \;-\; \sin\theta\;\Big(U_{\mathbf{k}}(t)\;\alpha^r_{\mathbf{k},1} \;-\; \epsilon^r\;V_{\mathbf{k}}(t)\;\beta^{r\dagger}_{-\mathbf{k},1}\Big) \\
\beta^r_{-\mathbf{k},e}(t) &= \cos\theta\;\beta^r_{-\mathbf{k},1} \;+\; \sin\theta\;\Big(U^*_{\mathbf{k}}(t)\;\beta^r_{-\mathbf{k},2} \;-\; \epsilon^r\;V_{\mathbf{k}}(t)\;\alpha^{r\dagger}_{\mathbf{k},2}\Big) \\
\beta^r_{-\mathbf{k},\mu}(t) &= \cos\theta\;\beta^r_{-\mathbf{k},2} \;-\; \sin\theta\;\Big(U_{\mathbf{k}}(t)\;\beta^r_{-\mathbf{k},1} \;+\; \epsilon^r\;V_{\mathbf{k}}(t)\;\alpha^{r\dagger}_{\mathbf{k},1}\Big) ,
\end{aligned} \tag{2.38}$$

with $\epsilon^r = (-1)^r$ and

$$\begin{aligned}
U_{\mathbf{k}}(t) &\equiv u^{r\dagger}_{\mathbf{k},2}(t)u^r_{\mathbf{k},1}(t) = v^{r\dagger}_{-\mathbf{k},1}(t)v^r_{-\mathbf{k},2}(t) \\
V_{\mathbf{k}}(t) &\equiv \epsilon^r\;u^{r\dagger}_{\mathbf{k},1}(t)v^r_{-\mathbf{k},2}(t) = -\epsilon^r\;u^{r\dagger}_{\mathbf{k},2}(t)v^r_{-\mathbf{k},1}(t)
\end{aligned} \tag{2.39}$$

We have:

$$V_{\mathbf{k}}(t) = |V_{\mathbf{k}}|\;e^{i(\omega_{k,2}+\omega_{k,1})t} \quad , \quad U_{\mathbf{k}}(t) = |U_{\mathbf{k}}|\;e^{i(\omega_{k,2}-\omega_{k,1})t}; \tag{2.40}$$

$$\begin{aligned}
|U_{\mathbf{k}}| &= \left(\frac{\omega_{k,1}+m_1}{2\omega_{k,1}}\right)^{\frac{1}{2}}\left(\frac{\omega_{k,2}+m_2}{2\omega_{k,2}}\right)^{\frac{1}{2}}\left(1+\frac{\mathbf{k}^2}{(\omega_{k,1}+m_1)(\omega_{k,2}+m_2)}\right) \\
|V_{\mathbf{k}}| &= \left(\frac{\omega_{k,1}+m_1}{2\omega_{k,1}}\right)^{\frac{1}{2}}\left(\frac{\omega_{k,2}+m_2}{2\omega_{k,2}}\right)^{\frac{1}{2}}\left(\frac{k}{(\omega_{k,2}+m_2)}-\frac{k}{(\omega_{k,1}+m_1)}\right)
\end{aligned} \tag{2.41}$$

$$|U_{\mathbf{k}}|^2 + |V_{\mathbf{k}}|^2 = 1. \tag{2.42}$$

We thus see that, at level of annihilation operators, the structure of the mixing transformation is that of a Bogoliubov transformation nested into a rotation. The two transformations however cannot be disentangled, thus the mixing transformations Eq.(2.38) are essentially different from the usual Bogoliubov transformations.

We observe that $G_\theta^{-1}(t) = exp[\theta(S_- - S_+)]$ is just the generator for generalized coherent states of $SU(2)$: the flavor vacuum state is therefore an $SU(2)$ (time dependent) coherent state. Let us now obtain the explicit expression for $|0\rangle_{e,\mu}$ and investigate the infinite volume limit of Eq.(2.32).

Using the Gaussian decomposition, $G_\theta^{-1}(t)$ can be written as

$$G_\theta^{-1}(t) = exp[\theta(S_- - S_+)] = exp(-tan\theta\ S_+)\ exp(-2ln\ cos\theta\ S_3)\ exp(tan\theta\ S_-) \tag{2.43}$$

where $0 \leq \theta < \frac{\pi}{2}$. Eq.(2.32) then becomes

$$|0\rangle_{e,\mu} = \prod_{\mathbf{k}} |0\rangle^{\mathbf{k}}_{e,\mu} = \prod_{\mathbf{k}} exp(-tan\theta\ S_+^{\mathbf{k}}) exp(-2ln\ cos\theta\ S_3^{\mathbf{k}})\ exp(tan\theta\ S_-^{\mathbf{k}})|0\rangle_{1,2}\ . \tag{2.44}$$

The right hand side of Eq.(2.44) may be computed by using the relations

$$S_3^{\mathbf{k}}|0\rangle_{1,2} = 0\ \ ,\ \ S_\pm^{\mathbf{k}}|0\rangle_{1,2} \neq 0\ \ ,\ \ (S_\pm^{\mathbf{k}})^2|0\rangle_{1,2} \neq 0\ \ ,\ \ (S_\pm^{\mathbf{k}})^3|0\rangle_{1,2} = 0\ , \tag{2.45}$$

and other useful relations which are given in the Appendix B. The expression for $|0\rangle_{e,\mu}$ in terms of $S_\pm^{\mathbf{k}}$ and $S_3^{\mathbf{k}}$ is:

$$\begin{aligned} |0\rangle_{e,\mu} = \prod_{\mathbf{k}} |0\rangle^{\mathbf{k}}_{e,\mu} = \prod_{\mathbf{k}} \Big[& 1 + \sin\theta\cos\theta \left(S_-^{\mathbf{k}} - S_+^{\mathbf{k}}\right) + \frac{1}{2}\sin^2\theta\cos^2\theta \left((S_-^{\mathbf{k}})^2 + (S_+^{\mathbf{k}})^2\right) + \\ & - \sin^2\theta S_+^{\mathbf{k}} S_-^{\mathbf{k}} + \frac{1}{2}\sin^3\theta\cos\theta \left(S_-^{\mathbf{k}}(S_+^{\mathbf{k}})^2 - S_+^{\mathbf{k}}(S_-^{\mathbf{k}})^2\right) + \frac{1}{4}\sin^4\theta (S_+^{\mathbf{k}})^2(S_-^{\mathbf{k}})^2 \Big] |0\rangle_{1,2}\ . \end{aligned} \tag{2.46}$$

The state $|0\rangle_{e,\mu}$ is normalized to 1 (see Eq.(2.32)). Eq.(2.46) and Eqs.(2.24) and (2.25) exhibit the rich coherent state structure of $|0\rangle_{e,\mu}$.

Let us now compute ${}_{1,2}\langle 0|0\rangle_{e,\mu}$. We obtain

$${}_{1,2}\langle 0|0\rangle_{e,\mu} = \prod_{\mathbf{k}} \left(1 - \sin^2\theta\ {}_{1,2}\langle 0|S_+^{\mathbf{k}} S_-^{\mathbf{k}}|0\rangle_{1,2} + \frac{1}{4}\sin^4\theta\ {}_{1,2}\langle 0|(S_+^{\mathbf{k}})^2(S_-^{\mathbf{k}})^2|0\rangle_{1,2}\right) \tag{2.47}$$

where (see Appendix B)

$$\begin{aligned}
& {}_{1,2}\langle 0|S_+^{\mathbf{k}}S_-^{\mathbf{k}}|0\rangle_{1,2} = \\
= \; & {}_{1,2}\langle 0| \left(\sum_{\sigma,\tau}\sum_{r,s} \left[v_{-\mathbf{k},1}^{\sigma\dagger}(t) u_{\mathbf{k},2}^{\tau}(t) \right] \left[u_{\mathbf{k},2}^{s\dagger}(t) v_{-\mathbf{k},1}^{r}(t) \right] \beta_{-\mathbf{k},1}^{\sigma} \alpha_{\mathbf{k},2}^{\tau} \alpha_{\mathbf{k},2}^{s\dagger} \beta_{-\mathbf{k},1}^{r\dagger} \right) |0\rangle_{1,2} = \\
= \; & \sum_{r,s} |\; v_{-\mathbf{k},1}^{r\dagger}(t) u_{\mathbf{k},2}^{s}(t) \;|^2 \equiv 2|V_{\mathbf{k}}|^2 \, . \qquad (2.48)
\end{aligned}$$

In a similar way we find

$$ {}_{1,2}\langle 0|(S_+^{\mathbf{k}})^2(S_-^{\mathbf{k}})^2|0\rangle_{1,2} = 2|V_{\mathbf{k}}|^4 \, . \qquad (2.49)$$

The function $|V_{\mathbf{k}}|^2$ depends on $|\mathbf{k}|$ only through its modulus and it is always in the interval $[0, \frac{1}{2}[$. It has a maximum at $|\mathbf{k}| = \sqrt{m_1 m_2}$ ($\sqrt{m_1 m_2}$ is the scale of the condensation density) and $|V_{\mathbf{k}}|^2 = 0$ when $m_1 = m_2$. Also, $|V_{\mathbf{k}}|^2 \to 0$ when $k \to \infty$.

In conclusion we have

$$\begin{aligned}
{}_{1,2}\langle 0|0\rangle_{e,\mu} &= \prod_{\mathbf{k}} \left(1 - \sin^2\theta \; |V_{\mathbf{k}}|^2\right)^2 \equiv \prod_{k} \Gamma(\mathbf{k}) = \qquad (2.50) \\
&= \prod_{\mathbf{k}} e^{ln\,\Gamma(\mathbf{k})} = e^{\sum_{\mathbf{k}} ln\,\Gamma(\mathbf{k})}.
\end{aligned}$$

From the properties of $|V_{\mathbf{k}}|^2$ we have that $\Gamma(\mathbf{k}) < 1$ for any value of $\mathbf{k}$ and of the parameters m_1 and m_2. By using the continuous limit relation $\sum_{\mathbf{k}} \to \frac{V}{(2\pi)^3}\int d^3\mathbf{k}$, in the infinite volume limit we obtain

$$\lim_{V\to\infty} {}_{1,2}\langle 0|0\rangle_{e,\mu} = \lim_{V\to\infty} \; e^{\frac{V}{(2\pi)^3}\int d^3\mathbf{k}\; ln\,\Gamma(\mathbf{k})} = 0 \qquad (2.51)$$

Notice that Eq.(2.51) shows that the orthogonality between $|0\rangle_{e,\mu}$ and $|0\rangle_{1,2}$ is due to the infrared contributions which are taken in care by the infinite volume limit and therefore high momentum contributions do not influence the result (for this reason here we do not need to consider the regularization problem of the UV divergence of the integral of $ln\,\Gamma(\mathbf{k})$).

Of course, this orthogonality disappears when $\theta = 0$ and/or when $m_1 = m_2$ (because in this case $|V_{\mathbf{k}}|^2 = 0$ and no mixing occurs in Pontecorvo theory).

Eq.(2.51) expresses the unitary inequivalence in the infinite volume limit of the flavor and the mass representations and shows the absolutely non-trivial nature of the mixing transformations Eq.(2.1). In other words, the mixing transformations induce a physically non-trivial structure in the flavor vacuum which indeed turns out to be an $SU(2)$ generalized coherent state. In Section 2.5 we will see how such a vacuum structure may lead to phenomenological consequences in the neutrino oscillations.

From Eq.(2.51) we also see that Eq.(2.32) is a purely formal expression which only holds at finite volume.

We thus realize the limit of validity of the approximation usually adopted when the mass vacuum state (representation for definite mass operators) is identified with the vacuum for the flavor operators. We point out that even at finite volume the vacua identification is actually an approximation since the flavor vacuum is an $SU(2)$ generalized coherent state. In such an approximation, the coherent state structure and many physical features are missed.

It is also interesting to exhibit the explicit expression of $|0\rangle^{\mathbf{k}}_{e,\mu}$, at time $t = 0$, in the reference frame for which $\mathbf{k} = (0, 0, |\mathbf{k}|)$ (see Appendix D):

$$\begin{aligned} |0\rangle^{\mathbf{k}}_{e,\mu} &= \prod_r \Big[(1-\sin^2\theta\,|V_{\mathbf{k}}|^2) - \epsilon^r \sin\theta\,\cos\theta\,|V_{\mathbf{k}}|(\alpha^{r\dagger}_{\mathbf{k},1}\beta^{r\dagger}_{-\mathbf{k},2} + \alpha^{r\dagger}_{\mathbf{k},2}\beta^{r\dagger}_{-\mathbf{k},1}) + \\ &+ \epsilon^r \sin^2\theta\,|V_{\mathbf{k}}||U_{\mathbf{k}}|(\alpha^{r\dagger}_{\mathbf{k},1}\beta^{r\dagger}_{-\mathbf{k},1} - \alpha^{r\dagger}_{\mathbf{k},2}\beta^{r\dagger}_{-\mathbf{k},2}) + \sin^2\theta\,|V_{\mathbf{k}}|^2\alpha^{r\dagger}_{\mathbf{k},1}\beta^{r\dagger}_{-\mathbf{k},2}\alpha^{r\dagger}_{\mathbf{k},2}\beta^{r\dagger}_{-\mathbf{k},1}\Big]|0\rangle_{1,2} \end{aligned} \tag{2.52}$$

We see that the expression of the flavor vacuum $|0\rangle_{e,\mu}$ involves four different particle-antiparticle "couples".

The condensation density is given by

$$_{e,\mu}\langle 0|\alpha^{r\dagger}_{\mathbf{k},i}\alpha^{r}_{\mathbf{k},i}|0\rangle_{e,\mu} \;=\; _{e,\mu}\langle 0|\beta^{r\dagger}_{\mathbf{k},i}\beta^{r}_{\mathbf{k},i}|0\rangle_{e,\mu} \;=\; \sin^2\theta\,|V_{\mathbf{k}}|^2\,, \qquad i = 1, 2\,. \tag{2.53}$$

2.3 Generalization of mixing transformations

In Section 2.2 we have expressed the flavor fields ν_e and ν_μ in the same bases as the (free) fields with definite masses ν_1 and ν_2, respectively. However, it has been noticed [28, 29], that this is actually a special choice, and that a more general possibility exists.

Let us introduce the notation $(\sigma, j) = (e, 1), (\mu, 2)$, the fields ν_e and ν_μ can be rewritten in the following form:

$$\nu_\sigma(x) \;=\; G^{-1}_\theta(t)\,\nu_j(x)\,G_\theta(t) = \frac{1}{\sqrt{V}}\sum_{\mathbf{k},r}\Big[u^r_{\mathbf{k},j}\alpha^r_{\mathbf{k},\sigma}(t) + v^r_{-\mathbf{k},j}\beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\Big]\,e^{i\mathbf{k}\cdot\mathbf{x}}, \tag{2.54}$$

The flavor annihilation operators are rewritten as

$$\left(\begin{array}{c}\alpha^r_{\mathbf{k},\sigma}(t)\\ \beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\end{array}\right) = G^{-1}_\theta(t)\left(\begin{array}{c}\alpha^r_{\mathbf{k},j}(t)\\ \beta^{r\dagger}_{-\mathbf{k},j}(t)\end{array}\right)G_\theta(t) \tag{2.55}$$

and the explicit expression of the flavor annihilation operators is (we choose $\mathbf{k} = (0, 0, |\mathbf{k}|)$):

$$\left(\begin{array}{c} \alpha^r_{\mathbf{k},e}(t) \\ \alpha^r_{\mathbf{k},\mu}(t) \\ \beta^{r\dagger}_{-\mathbf{k},e}(t) \\ \beta^{r\dagger}_{-\mathbf{k},\mu}(t) \end{array}\right) = \left(\begin{array}{cccc} c_\theta & s_\theta\,|U_\mathbf{k}| & 0 & s_\theta\,\epsilon^r\,|V_\mathbf{k}| \\ -s_\theta\,|U_\mathbf{k}| & c_\theta & s_\theta\,\epsilon^r\,|V_\mathbf{k}| & 0 \\ 0 & -s_\theta\,\epsilon^r\,|V_\mathbf{k}| & c_\theta & s_\theta\,|U_\mathbf{k}| \\ -s_\theta\,\epsilon^r\,|V_\mathbf{k}| & 0 & -s_\theta\,|U_\mathbf{k}| & c_\theta \end{array}\right) \left(\begin{array}{c} \alpha^r_{\mathbf{k},1}(t) \\ \alpha^r_{\mathbf{k},2}(t) \\ \beta^{r\dagger}_{-\mathbf{k},1}(t) \\ \beta^{r\dagger}_{-\mathbf{k},2}(t) \end{array}\right) \tag{2.56}$$

where $c_\theta \equiv \cos\theta$, $s_\theta \equiv \sin\theta$, $\epsilon^r \equiv (-1)^r$

In the expansion Eq.(2.54) one could use eigenfunctions with arbitrary masses μ_σ, and therefore not necessarily the same as the masses which appear in the Lagrangian. Indeed, the transformation Eq.(2.55) can be generalized [28, 29] by writing the flavor fields as

$$\nu_\sigma(x) \;=\; \frac{1}{\sqrt{V}}\sum_{\mathbf{k},r}\left[u^r_{\mathbf{k},\sigma}\widetilde{\alpha}^r_{\mathbf{k},\sigma}(t) + v^r_{-\mathbf{k},\sigma}\widetilde{\beta}^{r\dagger}_{-\mathbf{k},\sigma}(t)\right]e^{i\mathbf{k}\cdot\mathbf{x}}, \tag{2.57}$$

where u_σ and v_σ are the helicity eigenfunctions with mass ${\mu_\sigma}$[1]. We denote by a tilde the generalized flavor operators [28, 29] in order to distinguish them from the ones defined in Eq.(2.55). The expansion Eq.(2.57) is more general than the one in Eq.(2.54) since the latter corresponds to the particular choice $\mu_e \equiv m_1$, $\mu_\mu \equiv m_2$.

The relation between the general flavor and the mass operators is now:

$$\left(\begin{array}{c} \widetilde{\alpha}^r_{\mathbf{k},\sigma}(t) \\ \widetilde{\beta}^{r\dagger}_{-\mathbf{k},\sigma}(t) \end{array}\right) = K^{-1}_{\theta,\mu}(t)\left(\begin{array}{c} \alpha^r_{\mathbf{k},j}(t) \\ \beta^{r\dagger}_{-\mathbf{k},j}(t) \end{array}\right) K_{\theta,\mu}(t) \quad , \tag{2.58}$$

with $(\sigma, j) = (e, 1), (\mu, 2)$, where $K_{\theta,\mu}(t)$ is the generator of the transformation Eq.(2.1) and can be expressed as

$$K_{\theta,\mu}(t) \;=\; I_\mu(t)\,G_\theta(t) \tag{2.59}$$

$$I_\mu(t) \;=\; \prod_{\mathbf{k},r}\exp\left\{i\sum_{(\sigma,j)}\xi^{\mathbf{k}}_{\sigma,j}\left[\alpha^{r\dagger}_{\mathbf{k},j}(t)\beta^{r\dagger}_{-\mathbf{k},j}(t) + \beta^r_{-\mathbf{k},j}(t)\alpha^r_{\mathbf{k},j}(t)\right]\right\} \tag{2.60}$$

with

$$\xi^{\mathbf{k}}_{\sigma,j} \equiv (\chi_\sigma - \chi_j)/2, \qquad \cot\chi_\sigma = |\mathbf{k}|/\mu_\sigma, \qquad \cot\chi_j = |\mathbf{k}|/m_j. \tag{2.61}$$

For $\mu_e \equiv m_1$, $\mu_\mu \equiv m_2$, we have $I_\mu(t) = 1$.

---

[1] The use of such a basis simplifies considerably calculations with respect to the original choice of ref.[25].

The explicit matrix form of the flavor operators is [28, 29]:

$$\left(\begin{array}{c} \widetilde{\alpha}^r_{\mathbf{k},e}(t) \\ \widetilde{\alpha}^r_{\mathbf{k},\mu}(t) \\ \widetilde{\beta}^{r\dagger}_{-\mathbf{k},e}(t) \\ \widetilde{\beta}^{r\dagger}_{-\mathbf{k},\mu}(t) \end{array}\right) = \left(\begin{array}{cccc} c_\theta\,\rho^{\mathbf{k}}_{e1} & s_\theta\,\rho^{\mathbf{k}}_{e2} & ic_\theta\,\lambda^{\mathbf{k}}_{e1} & is_\theta\,\lambda^{\mathbf{k}}_{e2} \\ -s_\theta\,\rho^{\mathbf{k}}_{\mu 1} & c_\theta\,\rho^{\mathbf{k}}_{\mu 2} & -is_\theta\,\lambda^{\mathbf{k}}_{\mu 1} & ic_\theta\,\lambda^{\mathbf{k}}_{\mu 2} \\ ic_\theta\,\lambda^{\mathbf{k}}_{e1} & is_\theta\,\lambda^{\mathbf{k}}_{e2} & c_\theta\,\rho^{\mathbf{k}}_{e1} & s_\theta\,\rho^{\mathbf{k}}_{e2} \\ -is_\theta\,\lambda^{\mathbf{k}}_{\mu 1} & ic_\theta\,\lambda^{\mathbf{k}}_{\mu 2} & -s_\theta\,\rho^{\mathbf{k}}_{\mu 1} & c_\theta\,\rho^{\mathbf{k}}_{\mu 2} \end{array}\right) \left(\begin{array}{c} \alpha^r_{\mathbf{k},1}(t) \\ \alpha^r_{\mathbf{k},2}(t) \\ \beta^{r\dagger}_{-\mathbf{k},1}(t) \\ \beta^{r\dagger}_{-\mathbf{k},2}(t) \end{array}\right) \tag{2.62}$$

where $c_\theta \equiv \cos\theta$, $s_\theta \equiv \sin\theta$ and

$$\rho^{\mathbf{k}}_{ab}\delta_{rs} \equiv \cos\left(\frac{\chi_a - \chi_b}{2}\right)\delta_{rs} = u^{r\dagger}_{\mathbf{k},a}u^{s}_{\mathbf{k},b} = v^{r\dagger}_{-\mathbf{k},a}v^{s}_{-\mathbf{k},b} \tag{2.63}$$

$$i\lambda^{\mathbf{k}}_{ab}\delta_{rs} \equiv i\sin\left(\frac{\chi_a - \chi_b}{2}\right)\delta_{rs} = u^{r\dagger}_{\mathbf{k},a}v^{s}_{-\mathbf{k},b} = v^{r\dagger}_{-\mathbf{k},a}u^{s}_{\mathbf{k},b} \tag{2.64}$$

with $a, b = 1, 2, e, \mu$.

Since $\rho^{\mathbf{k}}_{12} = |U_{\mathbf{k}}|$ and $i\lambda^{\mathbf{k}}_{12} = \epsilon^r|V_{\mathbf{k}}|$, etc., the operators Eq.(2.62) reduce to the ones in Eqs.(2.56) when $\mu_e \equiv m_1$ and $\mu_\mu \equiv m_2$[2].

The generalization of the flavor vacuum, which is annihilated by the general flavor operators given by Eq.(2.58), is [28, 29]:

$$|\widetilde{0}(t)\rangle_{e,\mu} \equiv K^{-1}_{\theta,\mu}(t)|0\rangle_{1,2} \quad . \tag{2.65}$$

For $\mu_e \equiv m_1$ and $\mu_\mu \equiv m_2$, this state reduces to the flavor vacuum $|0(t)\rangle_{e,\mu}$ above defined.

The relation between the general flavor operators of Eq.(2.58) and the flavor operators of Eq.(2.55) is [28, 29]:

$$\left(\begin{array}{c} \widetilde{\alpha}^r_{\mathbf{k},\sigma}(t) \\ \widetilde{\beta}^{r\dagger}_{-\mathbf{k},\sigma}(t) \end{array}\right) = J^{-1}_\mu(t)\left(\begin{array}{c} \alpha^r_{\mathbf{k},\sigma}(t) \\ \beta^{r\dagger}_{-\mathbf{k},\sigma}(t) \end{array}\right) J_\mu(t) \quad , \tag{2.66}$$

$$J_\mu(t) = \prod_{\mathbf{k},r} \exp\left\{ i\sum_{(\sigma,j)} \xi^{\mathbf{k}}_{\sigma,j}\left[\alpha^{r\dagger}_{\mathbf{k},\sigma}(t)\beta^{r\dagger}_{-\mathbf{k},\sigma}(t) + \beta^{r}_{-\mathbf{k},\sigma}(t)\alpha^{r}_{\mathbf{k},\sigma}(t)\right]\right\} \tag{2.67}$$

We have shown that the Hilbert space for the flavor fields is not unique: an infinite number of vacua (and consequently infinitely many Hilbert spaces) can be generated by introducing the arbitrary mass parameters μ_e, μ_μ. It is obvious that physical quantities must not depend on these parameters. In Section 2.6 we show indeed that the observable charge operators are invariant under the transformations generated by Eq.(2.67). Thus, physical observables are fully independent of the arbitrary mass parameters μ_σ. See also Appendix I.

---

[2] In performing such an identification, one should take into account that the operators for antiparticles differ for a minus sign, related to the different spinor bases used in the expansions Eqs.(2.54) and (2.57). Such a sign difference is however irrelevant in what follows.

2.4 The current structure for field mixing

We now analize the transformations acting on a doublet of free fields with different masses. The results of this Section clarify the meaning of the $su(2)$ algebraic structure found before and will be useful in the discussion of neutrino oscillations.

Let us consider the Lagrangian describing two free Dirac fields with masses m_1 and m_2:

$$\mathcal{L}(x) \,=\, \bar{\Psi}_m(x) \left(i \not\!\partial - \mathsf{M}_d \right) \Psi_m(x) \,, \tag{2.68}$$

where $\Psi_m^T = (\nu_1, \nu_2)$ and $\mathsf{M}_d = diag(m_1, m_2)$. We introduce a subscript m, in order to distinguish the quantities here introduced, which are in terms of fields with definite masses, from the ones which are in terms of flavor fields.

The Lagrangian $\mathcal{L}(x)$ is invariant under global $U(1)$ phase transformations of the type

$$\Psi_m^{'}(x) = e^{i\alpha} \Psi_m(x), \tag{2.69}$$

then, we have the conservation of the Noether charge

$$Q_{=} \int I^0(x) d^3\mathbf{x} \tag{2.70}$$

(with $I^\mu(x) = \bar{\Psi}_m(x)\gamma^\mu \Psi_m(x)$) which is indeed the total charge of the system, i.e. the total lepton number.

Consider then the global $SU(2)$ transformation [19]:

$$\Psi_m^{'}(x) = e^{i\alpha_j \cdot \tau_j} \Psi_m(x) \qquad\qquad j = 1, 2, 3. \tag{2.71}$$

with α_j real constants, $\tau_j = \sigma_j/2$ and σ_j being the Pauli matrices.

Since the masses m_1 and m_2 are different, the Lagrangian is not invariant under the above transformations. By use of equations of motion, we obtain the variation of the Lagrangian:

$$\delta\mathcal{L} = i\alpha_j \bar{\Psi}_m(x) \left[\tau_j ,\ M_d\right] \Psi_m(x) = -\alpha_j \partial_\mu J^\mu_{m,j}(x), \tag{2.72}$$

where the currents for a complex field $\Psi^\alpha(x)$ are given by:

$$J^\mu(x) = i \left\{ \Psi^{\dagger\alpha}(x) \frac{\partial\mathcal{L}}{\partial\left(\partial_\mu \Psi^{\dagger\alpha}(x)\right)} - \frac{\partial\mathcal{L}}{\partial\left(\partial_\mu \Psi^{\alpha}(x)\right)} \Psi^\alpha(x) \right\}, \tag{2.73}$$

and then, in our case

$$J^\mu_{m,j}(x) = \bar{\Psi}_m(x) \ \gamma^\mu \ \tau_j \ \Psi_m(x), \qquad\qquad j = 1, 2, 3. \tag{2.74}$$

We thus have the following currents:

$$J^{\mu}_{m,1}(x) = \frac{1}{2}\left[\bar{\nu}_1(x)\, \gamma^{\mu}\, \nu_2(x) + \bar{\nu}_2(x)\, \gamma^{\mu}\, \nu_1(x)\right], \tag{2.75}$$

$$J^{\mu}_{m,2}(x) = \frac{i}{2}\left[\bar{\nu}_1(x)\, \gamma^{\mu}\, \nu_2(x) - \bar{\nu}_2(x)\, \gamma^{\mu}\, \nu_1(x)\right], \tag{2.76}$$

$$J^{\mu}_{m,3}(x) = \frac{1}{2}\left[\bar{\nu}_1(x)\, \gamma^{\mu}\, \nu_1(x) - \bar{\nu}_2(x)\, \gamma^{\mu}\, \nu_2(x)\right]. \tag{2.77}$$

The related charges, defined as

$$Q_{m,j}(t) = \int J^{0}_{m,j}(x) d^3\mathbf{x}, \qquad\qquad j = 1, 2, 3, \tag{2.78}$$

satisfy the $su(2)$ algebra:

$$[Q_{m,i}(t), Q_{m,j}(t)] = i\varepsilon_{ijk} Q_{m,k}(t). \tag{2.79}$$

We note that the Casimir operator is proportional to the total (conserved) charge:

$$Q_{m,0} = \frac{1}{2} Q. \tag{2.80}$$

Also $Q_{m,3}$ is conserved, due to the fact that the mass matrix M_d is diagonal. This implies the conservation of charge separately for ν_1 and ν_2, which is what we expect for a system of two non-interacting fields. We can thus define the combinations:

$$Q_1 = \frac{1}{2} Q + Q_{m,3}, \tag{2.81}$$

$$Q_2 = \frac{1}{2} Q - Q_{m,3}, \tag{2.82}$$

$$Q_i = \sum_{\mathbf{k},r} \left(\alpha^{r\dagger}_{\mathbf{k},i}(t) \alpha^{r}_{\mathbf{k},i}(t) - \beta^{r\dagger}_{-\mathbf{k},i}(t) \beta^{r}_{-\mathbf{k},i}(t) \right), \qquad i = 1, 2. \tag{2.83}$$

These are the Noether charges associated with the non interacting fields ν_1 and ν_2. Explicitly, the transformations induced by the three above generators τ_1, τ_2, τ_3 are

$$\Psi'_m = \begin{pmatrix} \cos\theta_1 & i\sin\theta_1 \\ i\sin\theta_1 & \cos\theta_1 \end{pmatrix} \Psi_m, \tag{2.84}$$

$$\Psi_m^{'} = \begin{pmatrix} \cos\theta_2 & \sin\theta_2 \\ -\sin\theta_2 & \cos\theta_2 \end{pmatrix} \Psi_m, \tag{2.85}$$

$$\Psi_m^{'} = \begin{pmatrix} e^{i\theta_3} & 0 \\ 0 & e^{-i\theta_3} \end{pmatrix} \Psi_m. \tag{2.86}$$

We observe that the transformation induced by $Q_{m,2}(t)$,

$$\Psi_f(x) = e^{-2i\theta Q_{m,2}(t)}\ \Psi_m(x)\ e^{2i\theta Q_{m,2}(t)} \tag{2.87}$$

is just the mixing transformation Eq.(2.1).

Let us consider the Lagrangian written in the flavor basis (the subscript f denotes here flavor)

$$\mathcal{L}(x) \ = \ \bar{\Psi}_f(x)\,(i\ \partial\!\!\!/ - \mathsf{M})\,\Psi_f(x)\,, \tag{2.88}$$

where $\Psi_f^T = (\nu_e,\, \nu_\mu)$, and $\mathsf{M} = \begin{pmatrix} m_e & m_{e\mu} \\ m_{e\mu} & m_\mu \end{pmatrix}$.

In analogy with was done above, consider now the variation of the Lagrangian Eq.(2.88) under the $SU(2)$ transformation:

$$\Psi_f^{'}(x) = e^{i\alpha_j\cdot\tau_j}\Psi_f(x) \qquad\qquad j = 1, 2, 3. \tag{2.89}$$

We have

$$\delta\mathcal{L}(x) = i\alpha_j\bar{\Psi}_f(x)\,[\tau_j, \mathsf{M}]\,\Psi_f(x) = -\alpha_j\partial_\mu J^\mu_{f,j}(x), \tag{2.90}$$

$$J^\mu_{f,j}(x) = \bar{\Psi}_f(x)\ \gamma^\mu\ \tau_j\ \Psi_f(x) \tag{2.91}$$

and obtain the currents:

$$J^\mu_{f,1}(x) = \frac{1}{2}\left[\bar{\nu}_e(x)\ \gamma^\mu\ \nu_\mu(x) + \bar{\nu}_\mu(x)\ \gamma^\mu\ \nu_e(x)\right], \tag{2.92}$$

$$J^\mu_{f,2}(x) = \frac{i}{2}\left[\bar{\nu}_e(x)\ \gamma^\mu\ \nu_\mu(x) - \bar{\nu}_\mu(x)\ \gamma^\mu\ \nu_e(x)\right], \tag{2.93}$$

$$J^\mu_{f,3}(x) = \frac{1}{2}\left[\bar{\nu}_e(x)\ \gamma^\mu\ \nu_e(x)\ - \bar{\nu}_\mu(x)\gamma^\mu\ \nu_\mu(x)\right], \tag{2.94}$$

and

$$J^\mu_{f,0}(x) = \frac{1}{2}\left[\bar{\nu}_e(x)\ \gamma^\mu\ \nu_e(x) + \bar{\nu}_\mu(x)\ \gamma^\mu\ \nu_\mu(x)\right]. \tag{2.95}$$

Again, the charges

$$Q_{f,j}(t) = \int J^0_{f,j}(x) d^3\mathbf{x} \qquad j = 1, 2, 3, \tag{2.96}$$

satisfy the $su(2)$ algebra: $[Q_{f,i}(t), Q_{f,j}(t)] = i\varepsilon_{i,jk} Q_{f,k}(t)$.

The Casimir $Q_{f,0}$ is proportional to the total charge $Q_{f,0} = Q_0 = \frac{1}{2}Q$.

However, now, because of the off-diagonal (mixing) terms in the mass matrix M, $Q_{f,3}$ is not conserved anymore. This implies an exchange of charge between ν_e and ν_μ, resulting in the phenomenon of neutrino oscillations.

Let us indeed define the *flavor charges* as

$$Q_e(t) \equiv \frac{1}{2}Q + Q_{f,3}(t), \tag{2.97}$$

$$Q_\mu(t) \equiv \frac{1}{2}Q - Q_{f,3}(t), \tag{2.98}$$

where

$$Q_e(t) + Q_\mu(t) = Q, \tag{2.99}$$

the oscillation formulas are obtained by taking expectation values of the above charges on the neutrino state.

2.5 The exact formula for neutrino mixing

In terms of the flavor operators, the flavor charge operators are

$$Q_\sigma(t) = \sum_{\mathbf{k},r} \left(\alpha^{r\dagger}_{\mathbf{k},\sigma}(t) \alpha^r_{\mathbf{k},\sigma}(t) - \beta^{r\dagger}_{-\mathbf{k},\sigma}(t) \beta^r_{-\mathbf{k},\sigma}(t) \right), \qquad \sigma = e, \mu. \tag{2.100}$$

At time $t = 0$, the vacuum state is $|0\rangle_{e,\mu}$ and the one electron neutrino state is (for $\mathbf{k} = (0, 0, |\mathbf{k}|)$):

$$|\nu_e\rangle \equiv \alpha^{r\dagger}_{\mathbf{k},e}|0\rangle_{e,\mu} = \left[\cos\theta\, \alpha^{r\dagger}_{\mathbf{k},1} + |U_\mathbf{k}|\, \sin\theta\, \alpha^{r\dagger}_{\mathbf{k},2} + \epsilon^r\, |V_\mathbf{k}|\, \sin\theta\, \alpha^{r\dagger}_{\mathbf{k},2} \alpha^{r\dagger}_{\mathbf{k},1} \beta^{r\dagger}_{-\mathbf{k},1} \right] |0\rangle_{1,2}\,. \tag{2.101}$$

In this state a multiparticle component is present, disappearing in the relativistic limit $|\mathbf{k}| \gg \sqrt{m_1 m_2}$: in this limit the (quantum-mechanical) Pontecorvo state is recovered.

If we now assume that the neutrino state at time t is given by $|\nu_e(t)\rangle = e^{-iHt}|\nu_e\rangle$, we see that it is not possible to compare directly this state with the one at time $t = 0$ given in Eq.(2.101).

In fact, $|\nu_e(t)\rangle$ is given by:

$$\begin{aligned}|\nu_e(t)\rangle &= e^{-iHt}\,\alpha^{r\dagger}_{\mathbf{k},e}\,|0\rangle_{e,\mu} = e^{-i\omega_1 t}\Big[\cos\theta\,\alpha^{r\dagger}_{\mathbf{k},1} + e^{-i(\omega_2-\omega_1)t}\,|U_{\mathbf{k}}|\,\sin\theta\,\alpha^{r\dagger}_{\mathbf{k},2} + \\ &+ \epsilon^r\,e^{-i(\omega_2+\omega_1)t}\,|V_{\mathbf{k}}|\,\sin\theta\,\alpha^{r\dagger}_{\mathbf{k},2}\alpha^{r\dagger}_{\mathbf{k},1}\beta^{r\dagger}_{\mathbf{k},1}\Big]|0\rangle_{1,2} \qquad (2.102)\end{aligned}$$

and then (see the Appendix E)

$$\lim_{V\to\infty}\langle\nu_e(t)|\nu_e(0)\rangle = 0. \qquad (2.103)$$

The reason is that the flavor vacuum $|0\rangle_{e,\mu}$ is not eigenstate of the free Hamiltonian H and it "rotates" under the action of the time evolution generator: one indeed finds $\lim_{V\to\infty}\ {}_{e,\mu}\langle 0\mid 0(t)\rangle_{e,\mu} = 0$ (Appendix E). Thus at different times we have unitarily inequivalent flavor vacua (in the limit $V\to\infty$): this expresses the different particle content of these (coherent) states and it is direct consequence of the fact that flavor states are not mass eigenstates.

As already observed, this implies that we cannot directly compare flavor states at different times. However we can consider the flavor charge operators, defined as in Eq.(2.100). We then have (in the Heisenberg representation)

$${}_{e,\mu}\langle 0|Q_e(t)|0\rangle_{e,\mu} =_{e,\mu} \langle 0|Q_\mu(t)|0\rangle_{e,\mu} = 0, \qquad (2.104)$$

$$\mathcal{Q}^e_{\mathbf{k},e}(t) = \langle\nu_e|Q_e(t)|\nu_e\rangle = \left|\left\{\alpha^r_{\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2, \qquad (2.105)$$

$$\mathcal{Q}^e_{\mathbf{k},\mu}(t) = \langle\nu_e|Q_\mu(t)|\nu_e\rangle = \left|\left\{\alpha^r_{\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2. \qquad (2.106)$$

Charge conservation is obviously ensured at any time:

$$\mathcal{Q}^e_{\mathbf{k},e}(t) + \mathcal{Q}^e_{\mathbf{k},\mu}(t) = 1. \qquad (2.107)$$

The oscillation formula for the flavor charges are then [20]:

$$\begin{aligned}\mathcal{Q}^e_{\mathbf{k},e}(t) &= \left|\left\{\alpha^r_{\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 = \\ &= 1 - \sin^2(2\theta)\left[|U_{\mathbf{k}}|^2\sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) + |V_{\mathbf{k}}|^2\sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right],\end{aligned} \qquad (2.108)$$

$$\begin{aligned}\mathcal{Q}^e_{\mathbf{k},\mu}(t) &= \left|\left\{\alpha^r_{\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 = \\ &= \sin^2(2\theta)\left[|U_{\mathbf{k}}|^2\sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) + |V_{\mathbf{k}}|^2\sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right].\end{aligned} \qquad (2.109)$$

This result is exact. There are two differences with respect to the usual formula for neutrino oscillations: the amplitudes are energy dependent, and there is an additional oscillating term.

For $|\mathbf{k}| \gg \sqrt{m_1 m_2}$ we have $|U_{\mathbf{k}}|^2 \longrightarrow 1$ and $|V_{\mathbf{k}}|^2 \longrightarrow 0$ and the traditional formula is recovered.

2.6 Discussion

We conclude the chapter with a number of considerations about the oscillation formulas Eqs.(2.108), (2.109).

The Eqs.(2.108), (2.109) have a sense as statistical averages, i.e. as mean values. This is because the structure of the theory for mixed field is that of a many-body theory, where does not make sense to talk about single particle states.

This situation contrast with the quantum mechanical picture, which however is recovered in the ultra-relativistic limit. There, the approximate Pontecorvo result is recovered.

We now show [21] that the exact oscillation formulae are independent of the arbitrary mass parameters, indeed we have

$$\begin{aligned} &\left|\left\{\widetilde{\alpha}^r_{\mathbf{k},e}(t), \widetilde{\alpha}^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\widetilde{\beta}^{r\dagger}_{\mathbf{k},e}(t), \widetilde{\alpha}^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 = \\ &= \left|\left\{\alpha^r_{\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2, \end{aligned} \tag{2.110}$$

$$\begin{aligned} &\left|\left\{\widetilde{\alpha}^r_{\mathbf{k},\mu}(t), \widetilde{\alpha}^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\widetilde{\beta}^{r\dagger}_{-\mathbf{k},\mu}(t), \widetilde{\alpha}^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 = \\ &= \left|\left\{\alpha^r_{\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2, \end{aligned} \tag{2.111}$$

which ensure the cancellation of the arbitrary mass parameters.

However, the important point for the full understanding of the result Eqs.(2.110), (2.111) is that the charge operators Q_σ are *invariant* under the action of the Bogoliubov generator Eq.(2.67), i.e. $\widetilde{Q}_\sigma = Q_\sigma$, where $\widetilde{Q}_\sigma \equiv \widetilde{\alpha}^\dagger_\sigma \widetilde{\alpha}_\sigma - \widetilde{\beta}^\dagger_\sigma \widetilde{\beta}_\sigma$. Besides the direct computations leading to Eqs.(2.110), (2.111), such an invariance provides a strong and immediate proof of the independence of the oscillation formula from the μ_σ parameters. Thus, the expectation values of the flavor charge operators are the only physical relevant quantities in the context of the above theory, all other operators having expectation values depending on the arbitrary parameters above introduced.

Chapter 3

Quantum Field Theory of three flavor neutrino mixing and oscillations with CP violation

We analyze the Quantum Field Theory of mixing among three generations of Dirac fermions (neutrinos). We construct the Hilbert space for the flavor fields and determine the generators of the mixing transformations. By use of these generators, we recover all the known parameterizations of the three-flavor mixing matrix and we find a number of new ones. The algebra of the currents associated with the mixing transformations is shown to be a deformed $su(3)$ algebra, when CP violating phases are present. We then derive the exact oscillation formulas, exhibiting new features with respect to the usual ones. CP and T violation are also discussed.

3.1 Introduction

In this chapter we study in detail the case of three flavor fermion (neutrino) mixing. This is not a simple extension of the previous results [20, 21, 25] since the existence of a CP violating phase in the parameterization of the three-flavor mixing matrix introduces novel features which are absent in the two-flavor case. We determine the generators of the mixing transformations and by use of them, we recover the known parameterizations of the three-flavor mixing matrix and find a number of new ones. We construct the flavor Hilbert space, for which the ground state (flavor vacuum) turns out to be a generalized coherent state. We also study the algebraic structure of currents and charges associated with the mixing transformations and we find (as will show in the next chapter) that, in presence of CP violation, it is that of a deformed $su(3)$. The construction of the flavor Hilbert space is an essential step in the derivation of exact oscillation formulas, which account for CP violation and reduce to the corresponding quantum–mechanical ones in the relativistic limit.

The chapter is organized as follows. In Section 3.2 we construct the Hilbert space for three-flavor mixed fermions. In Section 3.3 we study the various parameterization of the unitary 3 × 3 mixing matrix obtained by use of the algebraic generators. In Section 3.4, we study the currents and charges for three-flavor mixing, which are then used in Section 3.5 to derive the exact neutrino oscillation formulas. Finally in Section 3.6, CP an T violation in QFT neutrino oscillations are discussed. The Section 3.7 is devoted to conclusions. In the Appendices F, G, H, I we put some useful formulas and a discussion of the arbitrary mass parameterization in the expansion of flavor fields as recently reported in [29, 30, 31].

3.2 Three flavor fermion mixing

We start by considering the following Lagrangian density describing three Dirac fields with a mixed mass term:

$$\mathcal{L}(x) \,=\, \bar{\Psi}_f(x)\,(i\,\not\partial - \mathsf{M})\,\Psi_f(x)\,, \tag{3.1}$$

where $\Psi_f^T = (\nu_e, \nu_\mu, \nu_\tau)$ and $\mathsf{M} = \mathsf{M}^\dagger$ is the mixed mass matrix.

Among the various possible parameterizations of the mixing matrix for three fields, we choose to work with the following one since it is the familiar parameterization of the CKM matrix [4]:

$$\Psi_f(x) \,= \mathcal{U}\,\Psi_m(x) = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \Psi_m(x), \tag{3.2}$$

with $c_{ij} = \cos\theta_{ij}$ and $s_{ij} = \sin\theta_{ij}$, being θ_{ij} the mixing angle between ν_i, ν_j and $\Psi_m^T = (\nu_1, \nu_2, \nu_3)$.

Using Eq.(3.2), we diagonalize the quadratic form of Eq.(3.1), which then reduces to the Lagrangian for three Dirac fields, with masses m_1, m_2 and m_3:

$$\mathcal{L}(x) \,=\, \bar{\Psi}_m(x)\,(i\,\not\partial - \mathsf{M}_d)\,\Psi_m(x)\,, \tag{3.3}$$

where $\mathsf{M}_d = diag(m_1, m_2, m_3)$.

As in the case of two flavor fermion mixing [25], we construct the generator for the mixing transformation (3.2) and define[1]

$$\nu_\sigma^\alpha(x) \equiv G_\theta^{-1}(t)\,\nu_i^\alpha(x)\,G_\theta(t), \tag{3.4}$$

---

[1] Let us consider for example the generation of the first row of the mixing matrix $\mathcal{U}$. We have

where $(\sigma, i) = (e,1), (\mu,2), (\tau,3)$, and

$$G_\theta(t) = G_{23}(t)\; G_{13}(t)\; G_{12}(t)\,, \tag{3.5}$$

where

$$G_{12}(t) \equiv \exp\Big[\theta_{12}L_{12}(t)\Big]; \qquad L_{12}(t) = \int d^3\mathbf{x}\left[\nu_1^\dagger(x)\nu_2(x) - \nu_2^\dagger(x)\nu_1(x)\right], \tag{3.6}$$

$$G_{23}(t) \equiv \exp\Big[\theta_{23}L_{23}(t)\Big]; \qquad L_{23}(t) = \int d^3\mathbf{x}\left[\nu_2^\dagger(x)\nu_3(x) - \nu_3^\dagger(x)\nu_2(x)\right], \tag{3.7}$$

$$G_{13}(t) \equiv \exp\Big[\theta_{13}L_{13}(\delta,t)\Big]; \quad L_{13}(\delta,t) = \int d^3\mathbf{x}\left[\nu_1^\dagger(x)\nu_3(x)e^{-i\delta} - \nu_3^\dagger(x)\nu_1(x)e^{i\delta}\right]. \tag{3.8}$$

It is evident from the above form of the generators, that the phase δ is unavoidable for three field mixing, while it can be incorporated in the definition of the fields in the two flavor case.

The free fields ν_i (i=1,2,3) can be quantized in the usual way [22] (we use $t \equiv x_0$):

$$\nu_i(x) = \sum_r \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\left[u^r_{\mathbf{k},i}(t)\alpha^r_{\mathbf{k},i} + v^r_{-\mathbf{k},i}(t)\beta^{r\dagger}_{-\mathbf{k},i}\right]e^{i\mathbf{k}\cdot\mathbf{x}}, \qquad i = 1,2,3\,, \tag{3.9}$$

with $u^r_{\mathbf{k},i}(t) = e^{-i\omega_{k,i}t}u^r_{\mathbf{k},i}$, $v^r_{\mathbf{k},i}(t) = e^{i\omega_{k,i}t}v^r_{\mathbf{k},i}$ and $\omega_{k,i} = \sqrt{\mathbf{k}^2 + m_i^2}$. The vacuum for the mass eigenstates is denoted by $|0\rangle_m$: $\alpha^r_{\mathbf{k},i}|0\rangle_m = \beta^r_{\mathbf{k},i}|0\rangle_m = 0$. The anticommutation relations are the usual ones; the wave function orthonormality and completeness relations are given in Eqs.(2.7).

An important result of previous chapter is the unitary inequivalence [25] (in the infinite volume limit) of the vacua for the flavor fields and for the fields with definite masses. There such an inequivalence was proved for the case of two generations; subsequently, in Ref.[32], a rigorous general proof of such inequivalence for any number

---

(see also Appendix H) $\partial\nu_e/\partial\theta_{23} = 0$; and

$$\partial\nu_e/\partial\theta_{13} = G_{12}^{-1}G_{13}^{-1}[\nu_1, L_{13}]G_{13}G_{12} = G_{12}^{-1}G_{13}^{-1}e^{-i\delta}\nu_3G_{13}G_{12},$$

thus:

$$\partial^2\nu_e/\partial\theta_{13}^2 = -\nu_e \quad\Rightarrow\quad \nu_e = f(\theta_{12})\cos\theta_{13} + g(\theta_{12})\sin\theta_{13};$$

with the initial conditions (from Eq.(3.4)): $f(\theta_{12}) = \nu_e|_{\theta_{13}=0}$ and $g(\theta_{12}) = \partial\nu_e/\partial\theta_{13}|_{\theta_{13}=0} = e^{-i\delta}\nu_3$. We also have

$$\partial^2 f(\theta_{12})/\partial\theta_{13}^2 = -f(\theta_{12}) \quad\Rightarrow\quad f(\theta_{12}) = A\cos\theta_{12} + B\sin\theta_{12}$$

with the initial conditions $A = \nu_e|_{\theta=0} = \nu_1$ and $B = \partial f(\theta_{12})/\partial\theta_{12}|_{\theta=0} = \nu_2$, and $\theta = (\theta_{12}, \theta_{13}, \theta_{23})$.

of generations has been given (see also Ref.[33]). Thus we define the *flavor vacuum* as:

$$|0(t)\rangle_f \;\equiv\; G_\theta^{-1}(t)\;|0\rangle_m\;. \tag{3.10}$$

The form of this state is considerably more complicated of the one for two generations. When $\delta = 0$, the generator G_θ is an element of the $SU(3)$ group (see Chapter 4) and the flavor vacuum is classified as an $SU(3)$ generalized coherent state à la Perelomov [44]. A nonzero CP violating phase introduces an interesting modification of the algebra associated with the mixing transformations Eq.(3.2): we discuss this in Chapter 4.

By use of $G_\theta(t)$, the flavor fields can be expanded as:

$$\nu_\sigma(x) = \sum_r \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\left[u^r_{\mathbf{k},i}(t)\alpha^r_{\mathbf{k},\sigma}(t) + v^r_{-\mathbf{k},i}(t)\beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\right]e^{i\mathbf{k}\cdot\mathbf{x}}, \tag{3.11}$$

with $(\sigma, i) = (e,1),(\mu,2),(\tau,3)$.

The flavor annihilation operators are defined as

$$\begin{aligned}\alpha^r_{\mathbf{k},\sigma}(t) &\equiv G_\theta^{-1}(t)\;\alpha^r_{\mathbf{k},i}\;G_\theta(t) && (3.12)\\ \beta^{r\dagger}_{-\mathbf{k},\sigma}(t) &\equiv G_\theta^{-1}(t)\;\beta^{r\dagger}_{-\mathbf{k},i}\;G_\theta(t). && (3.13)\end{aligned}$$

They clearly act as annihilators for the flavor vacuum Eq.(3.10). For further reference, it is useful to list explicitly the flavor annihilation/creation operators [30]. In the reference frame $\mathbf{k} = (0,0,|\mathbf{k}|)$ the spins decouple and their form is particularly simple:

$$\begin{aligned}\alpha^r_{\mathbf{k},e}(t) &= c_{12}c_{13}\;\alpha^r_{\mathbf{k},1} + s_{12}c_{13}\left(U^{\mathbf{k}*}_{12}(t)\;\alpha^r_{\mathbf{k},2} + \epsilon^r V^{\mathbf{k}}_{12}(t)\;\beta^{r\dagger}_{-\mathbf{k},2}\right) + \\ &\quad + e^{-i\delta}\;s_{13}\left(U^{\mathbf{k}*}_{13}(t)\;\alpha^r_{\mathbf{k},3} + \epsilon^r V^{\mathbf{k}}_{13}(t)\;\beta^{r\dagger}_{-\mathbf{k},3}\right)\;, && (3.14)\end{aligned}$$

$$\begin{aligned}\alpha^r_{\mathbf{k},\mu}(t) &= \left(c_{12}c_{23} - e^{i\delta}\;s_{12}s_{23}s_{13}\right)\;\alpha^r_{\mathbf{k},2} - \left(s_{12}c_{23} + e^{i\delta}\;c_{12}s_{23}s_{13}\right)\times \\ &\quad \times\left(U^{\mathbf{k}}_{12}(t)\;\alpha^r_{\mathbf{k},1} - \epsilon^r V^{\mathbf{k}}_{12}(t)\;\beta^{r\dagger}_{-\mathbf{k},1}\right) + \;s_{23}c_{13}\left(U^{\mathbf{k}*}_{23}(t)\;\alpha^r_{\mathbf{k},3} + \epsilon^r V^{\mathbf{k}}_{23}(t)\;\beta^{r\dagger}_{-\mathbf{k},3}\right)\;, && (3.15)\end{aligned}$$

$$\begin{aligned}\alpha^r_{\mathbf{k},\tau}(t) &= c_{23}c_{13}\;\alpha^r_{\mathbf{k},3} - \left(c_{12}s_{23} + e^{i\delta}\;s_{12}c_{23}s_{13}\right)\left(U^{\mathbf{k}}_{23}(t)\;\alpha^r_{\mathbf{k},2} - \epsilon^r V^{\mathbf{k}}_{23}(t)\;\beta^{r\dagger}_{-\mathbf{k},2}\right) + \\ &\quad + \;\left(s_{12}s_{23} - e^{i\delta}\;c_{12}c_{23}s_{13}\right)\left(U^{\mathbf{k}}_{13}(t)\;\alpha^r_{\mathbf{k},1} - \epsilon^r V^{\mathbf{k}}_{13}(t)\;\beta^{r\dagger}_{-\mathbf{k},1}\right)\;, && (3.16)\end{aligned}$$

$$\begin{aligned}\beta^r_{-\mathbf{k},e}(t) &= c_{12}c_{13}\ \beta^r_{-\mathbf{k},1} + s_{12}c_{13}\left(U^{\mathbf{k}*}_{12}(t)\ \beta^r_{-\mathbf{k},2} - \epsilon^r V^{\mathbf{k}}_{12}(t)\ \alpha^{r\dagger}_{\mathbf{k},2}\right) + \\ &\quad + e^{i\delta}\ s_{13}\left(U^{\mathbf{k}*}_{13}(t)\ \beta^r_{-\mathbf{k},3} - \epsilon^r V^{\mathbf{k}}_{13}(t)\ \alpha^{r\dagger}_{\mathbf{k},3}\right)\ ,\end{aligned} \tag{3.17}$$

$$\begin{aligned}\beta^r_{-\mathbf{k},\mu}(t) &= \left(c_{12}c_{23} - e^{-i\delta}\ s_{12}s_{23}s_{13}\right)\ \beta^r_{-\mathbf{k},2} - \left(s_{12}c_{23} + e^{-i\delta}\ c_{12}s_{23}s_{13}\right) \times \\ &\quad \times \left(U^{\mathbf{k}}_{12}(t)\ \beta^r_{-\mathbf{k},1} + \epsilon^r\ V^{\mathbf{k}}_{12}(t)\ \alpha^{r\dagger}_{\mathbf{k},1}\right) + \ s_{23}c_{13}\left(U^{\mathbf{k}*}_{23}(t)\ \beta^r_{-\mathbf{k},3} - \epsilon^r V^{\mathbf{k}}_{23}(t)\ \alpha^{r\dagger}_{\mathbf{k},3}\right)\ ,\end{aligned} \tag{3.18}$$

$$\begin{aligned}\beta^r_{-\mathbf{k},\tau}(t) &= c_{23}c_{13}\ \beta^r_{-\mathbf{k},3} - \left(c_{12}s_{23} + e^{-i\delta}\ s_{12}c_{23}s_{13}\right)\left(U^{\mathbf{k}}_{23}(t)\ \beta^r_{-\mathbf{k},2} + \epsilon^r V^{\mathbf{k}}_{23}(t)\ \alpha^{r\dagger}_{\mathbf{k},2}\right) \\ &\quad + \ \left(s_{12}s_{23} - e^{-i\delta}\ c_{12}c_{23}s_{13}\right)\left(U^{\mathbf{k}}_{13}(t)\ \beta^r_{-\mathbf{k},1} + \epsilon^r V^{\mathbf{k}}_{13}(t)\ \alpha^{r\dagger}_{\mathbf{k},1}\right)\ .\end{aligned} \tag{3.19}$$

These operators satisfy canonical (anti)commutation relations at equal times. The main difference with respect to their "naive" quantum-mechanical counterparts is in the anomalous terms proportional to the V_{ij} factors. In fact, $U^{\mathbf{k}}_{ij}$ and $V^{\mathbf{k}}_{ij}$ are Bogoliubov coefficients defined as:

$$V^{\mathbf{k}}_{ij}(t) = |V^{\mathbf{k}}_{ij}|\ e^{i(\omega_{k,j}+\omega_{k,i})t} \quad , \quad U^{\mathbf{k}}_{ij}(t) = |U^{\mathbf{k}}_{ij}|\ e^{i(\omega_{k,j}-\omega_{k,i})t} \tag{3.20}$$

$$|U^{\mathbf{k}}_{ij}| = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}}\left(\frac{\omega_{k,j}+m_j}{2\omega_{k,j}}\right)^{\frac{1}{2}}\left(1 + \frac{|\mathbf{k}|^2}{(\omega_{k,i}+m_i)(\omega_{k,j}+m_j)}\right) = \cos(\xi^{\mathbf{k}}_{ij}) \tag{3.21}$$

$$|V^{\mathbf{k}}_{ij}| = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}}\left(\frac{\omega_{k,j}+m_j}{2\omega_{k,j}}\right)^{\frac{1}{2}}\left(\frac{|\mathbf{k}|}{(\omega_{k,j}+m_j)} - \frac{|\mathbf{k}|}{(\omega_{k,i}+m_i)}\right) = \sin(\xi^{\mathbf{k}}_{ij}) \tag{3.22}$$

$$|U^{\mathbf{k}}_{ij}|^2 + |V^{\mathbf{k}}_{ij}|^2 = 1 \tag{3.23}$$

where $i, j = 1, 2, 3$ and $j > i$. The following identities hold:

$$V^{\mathbf{k}}_{23}(t)V^{\mathbf{k}*}_{13}(t) + U^{\mathbf{k}*}_{23}(t)U^{\mathbf{k}}_{13}(t) = U^{\mathbf{k}}_{12}(t), \tag{3.24}$$

$$V^{\mathbf{k}}_{23}(t)U^{\mathbf{k}*}_{13}(t) - U^{\mathbf{k}*}_{23}(t)V^{\mathbf{k}}_{13}(t) = -V^{\mathbf{k}}_{12}(t), \tag{3.25}$$

$$U^{\mathbf{k}}_{12}(t)U^{\mathbf{k}}_{23}(t) - V^{\mathbf{k}*}_{12}(t)V^{\mathbf{k}}_{23}(t) = U^{\mathbf{k}}_{13}(t), \tag{3.26}$$

$$U^{\mathbf{k}}_{23}(t)V^{\mathbf{k}}_{12}(t) + U^{\mathbf{k}*}_{12}(t)V^{\mathbf{k}}_{23}(t) = V^{\mathbf{k}}_{13}(t), \tag{3.27}$$

$$V^{\mathbf{k}*}_{12}(t)V^{\mathbf{k}}_{13}(t) + U^{\mathbf{k}*}_{12}(t)U^{\mathbf{k}}_{13}(t) = U^{\mathbf{k}}_{23}(t), \tag{3.28}$$

$$V^{\mathbf{k}}_{12}(t)U^{\mathbf{k}}_{13}(t) - U^{\mathbf{k}}_{12}(t)V^{\mathbf{k}}_{13}(t) = -V^{\mathbf{k}}_{23}(t)\ , \tag{3.29}$$

$$\xi^{\mathbf{k}}_{13} = \xi^{\mathbf{k}}_{12} + \xi^{\mathbf{k}}_{23} \qquad , \qquad \xi^{\mathbf{k}}_{ij} = \arctan\left(|V^{\mathbf{k}}_{ij}|\ /\ |U^{\mathbf{k}}_{ij}|\right)\ . \tag{3.30}$$

We observe that, in contrast with the case of two flavor mixing, the condensation densities are now different for particles of different masses [30]:

$$\mathcal{N}_1^{\mathbf{k}} = {}_f\langle 0(t)|N_{\alpha_1}^{\mathbf{k},r}|0(t)\rangle_f = {}_f\langle 0(t)|N_{\beta_1}^{\mathbf{k},r}|0(t)\rangle_f = s_{12}^2 c_{13}^2\,|V_{12}^{\mathbf{k}}|^2 + s_{13}^2\,|V_{13}^{\mathbf{k}}|^2\,, \tag{3.31}$$

$$\begin{aligned}\mathcal{N}_2^{\mathbf{k}} &= {}_f\langle 0(t)|N_{\alpha_2}^{\mathbf{k},r}|0(t)\rangle_f = {}_f\langle 0(t)|N_{\beta_2}^{\mathbf{k},r}|0(t)\rangle_f = \\ &= \left|-s_{12}c_{23} + e^{i\delta}\, c_{12}s_{23}s_{13}\right|^2 |V_{12}^{\mathbf{k}}|^2 + s_{23}^2 c_{13}^2\,|V_{23}^{\mathbf{k}}|^2\,,\end{aligned} \tag{3.32}$$

$$\begin{aligned}\mathcal{N}_3^{\mathbf{k}} &= {}_f\langle 0(t)|N_{\alpha_3}^{\mathbf{k},r}|0(t)\rangle_f = {}_f\langle 0(t)|N_{\beta_3}^{\mathbf{k},r}|0(t)\rangle_f = \\ &= \left|-c_{12}s_{23} + e^{i\delta}\, s_{12}c_{23}s_{13}\right|^2 |V_{23}^{\mathbf{k}}|^2 + \left|s_{12}s_{23} + e^{i\delta}\, c_{12}c_{23}s_{13}\right|^2 |V_{13}^{\mathbf{k}}|^2\,.\end{aligned} \tag{3.33}$$

We plot in Figure 3.1 the condensation densities for sample values of parameters as given in the table 3.1[2]:

m_1	m_2	m_3	θ_{12}	θ_{13}	θ_{23}	δ
1	200	3000	$\pi/4$	$\pi/4$	$\pi/4$	$\pi/4$

Table 3.1: The values of masses and mixing angles used for plots

---

[2] Here and in the following plots, we use the same (energy) units for the values of masses and momentum.



Figure 3.1: Plot of the condensation densities $\mathcal{N}_i^{\mathbf{k}}$ in function of $|\mathbf{k}|$ for the values of parameters as in Tab.(1).

3.3 The parameterizations of the three flavor mixing matrix

In Section 3.2 we have studied the generator of the mixing matrix $\mathcal{U}$ of Eq.(3.2). However, this matrix is only one of the various forms in which a 3×3 unitary matrix can be parameterized. Indeed, the generator Eq.(3.5) can be used for generating such alternative parameterizations. To see this, let us first define in a more general way the generators G_{ij} including phases for all of them:

$$G_{12}(t) \equiv \exp\Big[\theta_{12}L_{12}(\delta_{12},t)\Big]\ ;\quad L_{12}(\delta_{12},t) = \int d^3\mathbf{x}\left[\nu_1^\dagger(x)\nu_2(x)e^{-i\delta_{12}} - \nu_2^\dagger(x)\nu_1(x)e^{i\delta_{12}}\right], \tag{3.34}$$

$$G_{23}(t) \equiv \exp\Big[\theta_{23}L_{23}(\delta_{23},t)\Big]\ ;\quad L_{23}(\delta_{23},t) = \int d^3\mathbf{x}\left[\nu_2^\dagger(x)\nu_3(x)e^{-i\delta_{23}} - \nu_3^\dagger(x)\nu_2(x)e^{i\delta_{23}}\right], \tag{3.35}$$

$$G_{13}(t) \equiv \exp\Big[\theta_{13}L_{13}(\delta_{13},t)\Big]\ ;\quad L_{13}(\delta_{13},t) = \int d^3\mathbf{x}\left[\nu_1^\dagger(x)\nu_3(x)e^{-i\delta_{13}} - \nu_3^\dagger(x)\nu_1(x)e^{i\delta_{13}}\right]. \tag{3.36}$$

Six different matrices can be obtained by permuting the order of the G_{ij} (useful relations are listed in Appendices F and H) in Eq.(3.5). We obtain:

$$G_1 \equiv G_{23}G_{13}G_{12} \tag{3.37}$$

$$\mathcal{U}_1 = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13}e^{-i\delta_{12}} & s_{13}e^{-i\delta_{13}} \\ -s_{12}c_{23}e^{i\delta_{12}} - s_{23}s_{13}c_{12}e^{i(\delta_{13}-\delta_{23})} & c_{12}c_{23} - s_{23}s_{13}s_{12}e^{-i(\delta_{23}-\delta_{13}+\delta_{12})} & s_{23}c_{13}e^{-i\delta_{23}} \\ -s_{13}c_{12}c_{23}e^{i\delta_{13}} + s_{12}s_{23}e^{i(\delta_{12}+\delta_{23})} & -c_{23}s_{13}s_{12}e^{i(\delta_{13}-\delta_{12})} - s_{23}c_{12}e^{i\delta_{23}} & c_{23}c_{13} \end{pmatrix}$$

$$G_2 \equiv G_{23}G_{12}G_{13} \tag{3.38}$$

$$\mathcal{U}_2 = \begin{pmatrix} c_{12}c_{13} & s_{12}e^{-i\delta_{12}} & s_{13}c_{12}e^{-i\delta_{13}} \\ -s_{12}c_{13}c_{23}e^{i\delta_{12}} - s_{23}s_{13}e^{i(\delta_{13}-\delta_{23})} & c_{12}c_{23} & -s_{13}c_{23}s_{12}e^{i(\delta_{12}-\delta_{13})} + s_{23}c_{13}e^{-i\delta_{23}} \\ -s_{13}c_{23}e^{i\delta_{13}} + s_{12}s_{23}c_{13}e^{i(\delta_{12}+\delta_{23})} & -c_{12}s_{23}e^{i\delta_{23}} & c_{23}c_{13} + s_{12}s_{13}s_{23}e^{i(\delta_{12}+\delta_{23}-\delta_{13})} \end{pmatrix}$$

$$G_3 \equiv G_{13}G_{23}G_{12} \tag{3.39}$$

$$\mathcal{U}_3 = \begin{pmatrix} c_{12}c_{13} + s_{13}s_{23}s_{12}e^{i(\delta_{12}-\delta_{13}+\delta_{23})} & s_{12}c_{13}e^{-i\delta_{12}} - s_{13}s_{23}c_{12}e^{i(\delta_{23}-\delta_{13})} & s_{13}c_{23}e^{-i\delta_{13}} \\ -s_{12}c_{23}e^{i\delta_{12}} & c_{12}c_{23} & s_{23}e^{-i\delta_{23}} \\ c_{13}s_{23}s_{12}e^{i(\delta_{23}+\delta_{12})} - s_{13}c_{12}e^{i\delta_{13}} & -c_{13}s_{23}c_{12}e^{i\delta_{23}} - s_{12}s_{13}e^{i(\delta_{13}-\delta_{12})} & c_{23}c_{13} \end{pmatrix}$$

$$G_4 \equiv G_{13}G_{12}G_{23} \tag{3.40}$$

$$\mathcal{U}_4 = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13}c_{23}e^{-i\delta_{12}} - s_{13}s_{23}e^{i(\delta_{23}-\delta_{13})} & s_{12}s_{23}c_{13}e^{-i(\delta_{12}+\delta_{23})} + s_{13}c_{23}e^{-i\delta_{13}} \\ -s_{12}e^{i\delta_{12}} & c_{12}c_{23} & s_{23}c_{12}e^{-i\delta_{23}} \\ -c_{12}s_{13}e^{i\delta_{13}} & -c_{13}s_{23}e^{i\delta_{23}} - s_{12}c_{23}s_{13}e^{i(\delta_{13}-\delta_{12})} & c_{23}c_{13} - s_{12}s_{23}s_{13}e^{-i(\delta_{12}+\delta_{23}-\delta_{13})} \end{pmatrix}$$

$$G_5 \equiv G_{12}G_{13}G_{23} \tag{3.41}$$

$$
\mathcal{U}_5 \;=\; \begin{pmatrix} c_{12}c_{13} & s_{12}c_{23}e^{-i\delta_{12}} - s_{13}c_{12}s_{23}e^{-i(\delta_{13}-\delta_{23})} & s_{13}c_{12}c_{23}e^{-i\delta_{13}} + s_{12}s_{23}e^{-i(\delta_{12}+\delta_{23})} \\ -s_{12}c_{13}e^{i\delta_{12}} & c_{12}c_{23} + s_{12}s_{23}s_{13}e^{i(\delta_{12}-\delta_{13}+\delta_{23})} & s_{23}c_{12}e^{-i\delta_{23}} - s_{12}c_{23}s_{13}e^{i(\delta_{12}-\delta_{13})} \\ -s_{13}e^{i\delta_{13}} & -c_{13}s_{23}e^{i\delta_{23}} & c_{23}c_{13} \end{pmatrix}
$$

$$
G_6 \;\equiv\; G_{12}G_{23}G_{13} \tag{3.42}
$$

$$
\mathcal{U}_6 \;=\; \begin{pmatrix} c_{12}c_{13} - s_{12}s_{23}s_{13}e^{-i(\delta_{12}+\delta_{23}-\delta_{13})} & s_{12}c_{23}e^{-i\delta_{12}} & c_{12}s_{13}e^{-i\delta_{13}} + s_{12}s_{23}c_{13}e^{-i(\delta_{12}+\delta_{23})} \\ -c_{12}s_{23}s_{13}e^{i(\delta_{13}-\delta_{23})} - s_{12}c_{13}e^{i\delta_{12}} & c_{12}c_{23} & c_{12}s_{23}c_{13}e^{-i\delta_{23}} - s_{12}s_{13}e^{i(\delta_{12}-\delta_{13})} \\ -c_{23}s_{13}e^{i\delta_{13}} & -s_{23}e^{i\delta_{23}} & c_{23}c_{13} \end{pmatrix}
$$

The above matrices are generated for a particular set of initial conditions, namely for those of Eq.(3.4). The freedom in the choice of the initial conditions reflects into the possibility of obtaining other unitary matrices from the above ones by permuting rows and columns and by multiplying row or columns for a phase factor.

We thus can easily recover all the existing parameterizations of the CKM matrix [4, 17, 36, 37, 38, 39, 40]:

- the Maiani parameterization [36, 38] is obtained from $\mathcal{U}_1$ by setting $\theta_{12} \to \theta$, $\theta_{13} \to \beta$, $\theta_{23} \to \gamma$, $\delta_{12} \to 0$, $\delta_{13} \to 0$, $\delta_{23} \to -\delta$;

- the Chau–Keung parameterization [36, 39] is recovered from $\mathcal{U}_1$ by setting $\delta_{12} \to 0$ and $\delta_{23} \to 0$;

- the Kobayshi–Maskawa [4, 36] is recovered from $\mathcal{U}_5$ by setting $\theta_{12} \to \theta_2$, $\theta_{13} \to \theta_1$, $\theta_{23} \to \theta_3$, $\delta_{12} \to -\delta$, $\delta_{13} \to 0$ and $\delta_{23} \to 0$, $\theta_i \to \frac{3}{2}\pi - \theta_i$, with $i = 1, 2, 3$, and multiply the last column for (-1);

- the Anselm parameterization [36, 40] is obtained from $\mathcal{U}_1$ by setting $\theta_{12} \leftrightarrow \theta_{13}$, then $\delta_{12} \to 0$, $\delta_{13} \to 0$, $\theta_{12} \to \pi + \theta_{12}$, $\theta_{13} \to \pi - \theta_{13}$, $\theta_{23} \to \frac{3}{2}\pi + \theta_{23}$, exchanging second and third column and multiplying the last row for (-1).

From the above analysis it is clear that a number of new parameterizations of the mixing matrix can be generated and that a clear physical meaning can be attached to each of them, by considering the order in which the generators G_{ij} act and the initial conditions used for getting that particular matrix.

3.4 Currents and charges for three flavor fermion mixing

In this Section we study the currents associated to the Lagrangians Eqs.(3.1) and (3.3). To this end, let us consider the transformations acting on the triplet of free fields with different masses Ψ_m [19, 30].

$\mathcal{L}$ is invariant under global $U(1)$ phase transformations of the type $\Psi'_m = e^{i\alpha}\,\Psi_m$: as a result, we have the conservation of the Noether charge $Q = \int d^3\mathbf{x}\, I^0(x)$ (with $I^\mu(x) = \bar{\Psi}_m(x)\,\gamma^\mu\,\Psi_m(x)$) which is indeed the total charge of the system (i.e. the total lepton number).

Consider then the $SU(3)$ global transformations acting on Ψ_m:

$$\Psi'_m(x) \,=\, e^{i\alpha_j F_j}\,\Psi_m(x) \qquad , \qquad j=1,2,...,8. \tag{3.43}$$

with α_j real constants, $F_j = \frac{1}{2}\lambda_j$ being the generators of $SU(3)$ and λ_j the Gell-Mann matrices [4].

The Lagrangian is not generally invariant under (3.43) and we obtain, by use of the equations of motion,

$$\begin{aligned} \delta\mathcal{L}(x) \,&=\, i\alpha_j\,\bar{\Psi}_m(x)\,[F_j, \mathsf{M}_d]\,\Psi_m(x) \,=\, -\alpha_j\,\partial_\mu J^\mu_{m,j}(x) \\ J^\mu_{m,j}(x) \,&=\, \bar{\Psi}_m(x)\,\gamma^\mu\,F_j\,\Psi_m(x) \qquad , \qquad j=1,2,...,8. \end{aligned} \tag{3.44}$$

It is useful to list explicitly the eight currents:

$$J^\mu_{m,1}(x) \,=\, \frac{1}{2}\,[\bar{\nu}_1(x)\,\gamma^\mu\,\nu_2(x) \,+\, \bar{\nu}_2(x)\,\gamma^\mu\,\nu_1(x)] \tag{3.45}$$

$$J^\mu_{m,2}(x) \,=\, -\frac{i}{2}\,[\bar{\nu}_1(x)\,\gamma^\mu\,\nu_2(x) \,-\, \bar{\nu}_2(x)\,\gamma^\mu\,\nu_1(x)] \tag{3.46}$$

$$J^\mu_{m,3}(x) \,=\, \frac{1}{2}\,[\bar{\nu}_1(x)\,\gamma^\mu\,\nu_1(x) \,-\, \bar{\nu}_2(x)\,\gamma^\mu\,\nu_2(x)] \tag{3.47}$$

$$J^\mu_{m,4}(x) \,=\, \frac{1}{2}\,[\bar{\nu}_1(x)\,\gamma^\mu\,\nu_3(x) \,+\, \bar{\nu}_3(x)\,\gamma^\mu\,\nu_1(x)] \tag{3.48}$$

$$J^\mu_{m,5}(x) \,=\, -\frac{i}{2}\,[\bar{\nu}_1(x)\,\gamma^\mu\,\nu_3(x) \,-\, \bar{\nu}_3(x)\,\gamma^\mu\,\nu_1(x)] \tag{3.49}$$

$$J^\mu_{m,6}(x) \,=\, \frac{1}{2}\,[\bar{\nu}_2(x)\,\gamma^\mu\,\nu_3(x) \,+\, \bar{\nu}_3(x)\,\gamma^\mu\,\nu_2(x)\,] \tag{3.50}$$

$$J^\mu_{m,7}(x) \,=\, -\frac{i}{2}\,[\bar{\nu}_2(x)\,\gamma^\mu\,\nu_3(x) \,-\, \bar{\nu}_3(x)\,\gamma^\mu\,\nu_2(x)] \tag{3.51}$$

$$J^\mu_{m,8}(x) \,=\, \frac{1}{2\sqrt{3}}\,[\bar{\nu}_1(x)\,\gamma^\mu\,\nu_1(x) \,+\, \bar{\nu}_2(x)\,\gamma^\mu\,\nu_2(x) \,-2\bar{\nu}_3(x)\,\gamma^\mu\,\nu_3(x)]\,. \tag{3.52}$$

The related charges $Q_{m,j}(t) \equiv \int d^3\mathbf{x}\, J^0_{m,j}(x)$, satisfy the $su(3)$ algebra

$$[Q_{m,j}(t), Q_{m,k}(t)] \,=\, i\,f_{jkl}\,Q_{m,l}(t). \tag{3.53}$$

Note that only two of the above charges are time-independent, namely $Q_{m,3}$ and

$Q_{m,8}$. We can thus define the combinations:

$$\begin{aligned}
Q_1 &\equiv \frac{1}{3}Q + Q_{m,3} + \frac{1}{\sqrt{3}}Q_{m,8}, \\
Q_2 &\equiv \frac{1}{3}Q - Q_{m,3} + \frac{1}{\sqrt{3}}Q_{m,8}, \\
Q_3 &\equiv \frac{1}{3}Q - \frac{2}{\sqrt{3}}Q_{m,8},
\end{aligned} \tag{3.54}$$

$$Q_i = \sum_r \int d^3\mathbf{k} \left(\alpha^{r\dagger}_{\mathbf{k},i}\alpha^r_{\mathbf{k},i} - \beta^{r\dagger}_{-\mathbf{k},i}\beta^r_{-\mathbf{k},i}\right), \qquad i = 1,2,3. \tag{3.55}$$

These are nothing but the Noether charges associated with the non-interacting fields ν_1, ν_2 and ν_3: in the absence of mixing, they are the flavor charges, separately conserved for each generation.

As already observed in Section 3.2, in the case when CP is conserved ($\delta = 0$), the mixing generator Eq.(3.5) is an element of the $SU(3)$ group and can be expressed in terms of the above charges as:

$$G_\theta(t)|_{\delta=0} = e^{i2\theta_{23}\,Q_{m,7}(t)}\, e^{i2\theta_{13}\,Q_{m,5}(t)}\, e^{i2\theta_{12}\,Q_{m,2}(t)} \tag{3.56}$$

We can now perform the $SU(3)$ transformations on the flavor triplet Ψ_f and obtain another set of currents for the flavor fields:

$$\Psi'_f(x) = e^{i\alpha_j F_j}\,\Psi_f(x) \qquad , \qquad j = 1,2,...,8, \tag{3.57}$$

which leads to

$$\begin{aligned}
\delta\mathcal{L}(x) &= i\alpha_j\,\bar{\Psi}_f(x)\,[F_j, \mathsf{M}]\,\Psi_f(x) = -\alpha_j\,\partial_\mu J^\mu_{f,j}(x)\,, \\
J^\mu_{f,j}(x) &= \bar{\Psi}_f(x)\,\gamma^\mu\,F_j\,\Psi_f(x) \qquad , \qquad j = 1,2,...,8.
\end{aligned} \tag{3.58}$$

Alternatively, the same currents can be obtained by applying on the $J^\mu_{m,j}(x)$ the mixing generator Eq.(3.5):

$$J^\mu_{f,j}(x) = G^{-1}_\theta(t)\,J^\mu_{m,j}(x)\,G_\theta(t) \qquad , \qquad j = 1,2,...,8. \tag{3.59}$$

The related charges $Q_{f,j}(t) \equiv \int d^3\mathbf{x}\, J^0_{f,j}(x)$ still close the $su(3)$ algebra. Due to the off–diagonal (mixing) terms in the mass matrix M, $Q_{f,3}(t)$ and $Q_{f,8}(t)$ are time–dependent. This implies an exchange of charge between ν_e, ν_μ and ν_τ, resulting in the flavor oscillations.

In accordance with Eqs.(3.54) we define the *flavor charges* for mixed fields as

$$\begin{aligned} Q_e(t) &\equiv \frac{1}{3}Q + Q_{f,3}(t) + \frac{1}{\sqrt{3}}Q_{f,8}(t), \\ Q_\mu(t) &\equiv \frac{1}{3}Q - Q_{f,3}(t) + \frac{1}{\sqrt{3}}Q_{f,8}(t), \\ Q_\tau(t) &\equiv \frac{1}{3}Q - \frac{2}{\sqrt{3}}Q_{f,8}(t). \end{aligned} \tag{3.60}$$

with

$$Q_e(t) + Q_\mu(t) + Q_\tau(t) = Q. \tag{3.61}$$

These charges have a simple expression in terms of the flavor ladder operators:

$$Q_\sigma(t) = \sum_r \int d^3\mathbf{k} \left(\alpha^{r\dagger}_{\mathbf{k},\sigma}(t)\alpha^r_{\mathbf{k},\sigma}(t) - \beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\beta^r_{-\mathbf{k},\sigma}(t) \right) \tag{3.62}$$

with $\sigma = e, \mu, \tau$, because of the connection with the Noether charges of Eq.(3.55) via the mixing generator:

$$Q_\sigma(t) = G^{-1}_\theta(t)\; Q_i \; G_\theta(t). \tag{3.63}$$

Notice also that the operator $\Delta Q_\sigma(t) \equiv Q_\sigma(t) - Q_i$ with $(\sigma, i) = (e,1), (\mu,2), (\tau,3)$, describes how much the mixing violates the (lepton) charge conservation for a given generation.

3.5 Neutrino Oscillations

The oscillation formulas are obtained by taking expectation values of the above charges on the (flavor) neutrino state. Consider for example an initial electron neutrino state defined as $|\nu_e\rangle \equiv \alpha^{r\dagger}_{\mathbf{k},e}(0)|0\rangle_f$ [20, 21, 26]. Working in the Heisenberg picture, we obtain

$$\begin{aligned} \mathcal{Q}^\rho_{\mathbf{k},\sigma}(t) &\equiv \langle \nu_\rho | Q_\sigma(t) | \nu_\rho \rangle - {}_f\langle 0 | Q_\sigma(t) | 0 \rangle_f = \\ &= \left| \left\{ \alpha^r_{\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},\rho}(0) \right\} \right|^2 + \left| \left\{ \beta^{r\dagger}_{-\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},\rho}(0) \right\} \right|^2, \end{aligned} \tag{3.64}$$

$$\begin{aligned} \mathcal{Q}^{\bar\rho}_{\mathbf{k},\sigma}(t) &\equiv \langle \bar\nu_\rho | Q_\sigma(t) | \bar\nu_\rho \rangle - {}_f\langle 0 | Q_\sigma(t) | 0 \rangle_f = \\ &= -\left| \left\{ \beta^r_{\mathbf{k},\sigma}(t), \beta^{r\dagger}_{\mathbf{k},\rho}(0) \right\} \right|^2 - \left| \left\{ \alpha^{r\dagger}_{-\mathbf{k},\sigma}(t), \beta^{r\dagger}_{\mathbf{k},\rho}(0) \right\} \right|^2, \end{aligned} \tag{3.65}$$

where $|0\rangle_f \equiv |0(0)\rangle_f$. Overall charge conservation is obviously ensured at any time:

$$\mathcal{Q}^{\rho}_{\mathbf{k},e}(t) + \mathcal{Q}^{\rho}_{\mathbf{k},\mu}(t) + \mathcal{Q}^{\rho}_{\mathbf{k},\tau}(t) \; = \; 1, \qquad\qquad \rho = e, \mu, \tau. \tag{3.66}$$

We remark that the expectation value of Q_σ cannot be taken on vectors of the Fock space built on $|0\rangle_m$, [20, 21, 26]. Also we observe that ${}_f\langle 0|Q_\sigma(t)|0\rangle_f \neq 0$, in contrast with the two flavor case [21, 30, 31]. We introduce the following notation:

$$\Delta^{\mathbf{k}}_{ij} \equiv \frac{\omega_{k,j} - \omega_{k,i}}{2} \quad , \quad \Omega^{\mathbf{k}}_{ij} \equiv \frac{\omega_{k,i} + \omega_{k,j}}{2}$$

Then the oscillation (in time) formulae for the flavor charges, on an initial electron neutrino state, follow as [30]:

$$\begin{aligned}
\mathcal{Q}^{e}_{\mathbf{k},e}(t) \; &= \; 1 \, - \, \sin^2(2\theta_{12}) \cos^4\theta_{13} \Big[|U^{\mathbf{k}}_{12}|^2 \, \sin^2\left(\Delta^{\mathbf{k}}_{12} t\right) \, + \, |V^{\mathbf{k}}_{12}|^2 \, \sin^2\left(\Omega^{\mathbf{k}}_{12} t\right) \Big] \\
&\quad - \, \sin^2(2\theta_{13}) \cos^2\theta_{12} \Big[|U^{\mathbf{k}}_{13}|^2 \, \sin^2\left(\Delta^{\mathbf{k}}_{13} t\right) \, + \, |V^{\mathbf{k}}_{13}|^2 \, \sin^2\left(\Omega^{\mathbf{k}}_{13} t\right) \Big] \\
&\quad - \, \sin^2(2\theta_{13}) \sin^2\theta_{12} \Big[|U^{\mathbf{k}}_{23}|^2 \, \sin^2\left(\Delta^{\mathbf{k}}_{23} t\right) \, + \, |V^{\mathbf{k}}_{23}|^2 \, \sin^2\left(\Omega^{\mathbf{k}}_{23} t\right) \Big] \, ,
\end{aligned} \tag{3.67}$$

$$\begin{aligned}
\mathcal{Q}^{e}_{\mathbf{k},\mu}(t) \; &= \; 2J_{CP} \Big[|U^{\mathbf{k}}_{12}|^2 \, \sin(2\Delta^{\mathbf{k}}_{12} t) - |V^{\mathbf{k}}_{12}|^2 \, \sin(2\Omega^{\mathbf{k}}_{12} t) + (|U^{\mathbf{k}}_{12}|^2 - |V^{\mathbf{k}}_{13}|^2) \sin(2\Delta^{\mathbf{k}}_{23} t) \\
&+ \; (|V^{\mathbf{k}}_{12}|^2 - |V^{\mathbf{k}}_{13}|^2) \sin(2\Omega^{\mathbf{k}}_{23} t) - |U^{\mathbf{k}}_{13}|^2 \, \sin(2\Delta^{\mathbf{k}}_{13} t) + |V^{\mathbf{k}}_{13}|^2 \, \sin(2\Omega^{\mathbf{k}}_{13} t) \Big] \\
&+ \; \cos^2\theta_{13} \sin\theta_{13} \Big[\cos\delta \sin(2\theta_{12}) \sin(2\theta_{23}) + 4\cos^2\theta_{12} \sin\theta_{13} \sin^2\theta_{23} \Big] \times \\
&\times \; \Big[|U^{\mathbf{k}}_{13}|^2 \sin^2\left(\Delta^{\mathbf{k}}_{13} t\right) + |V^{\mathbf{k}}_{13}|^2 \; \sin^2\left(\Omega^{\mathbf{k}}_{13} t\right) \Big] \\
&- \; \cos^2\theta_{13} \sin\theta_{13} \Big[\cos\delta \sin(2\theta_{12}) \sin(2\theta_{23}) - 4\sin^2\theta_{12} \sin\theta_{13} \sin^2\theta_{23} \Big] \times \\
&\times \; \Big[|U^{\mathbf{k}}_{23}|^2 \; \sin^2\left(\Delta^{\mathbf{k}}_{23} t\right) + |V^{\mathbf{k}}_{23}|^2 \; \sin^2\left(\Omega^{\mathbf{k}}_{23} t\right) \Big] \\
&+ \; \cos^2\theta_{13} \sin(2\theta_{12}) \Big[(\cos^2\theta_{23} - \sin^2\theta_{23} \sin^2\theta_{13}) \sin(2\theta_{12}) \\
&+ \; \cos\delta \cos(2\theta_{12}) \sin\theta_{13} \sin(2\theta_{23}) \Big] \Big[|U^{\mathbf{k}}_{12}|^2 \; \sin^2\left(\Delta^{\mathbf{k}}_{12} t\right) + |V^{\mathbf{k}}_{12}|^2 \; \sin^2\left(\Omega^{\mathbf{k}}_{12} t\right) \Big] \, ,
\end{aligned} \tag{3.68}$$

$$
\begin{aligned}
\mathcal{Q}^{e}_{\mathbf{k},\tau}(t) &= -2J_{_{CP}}\Big[|U^{\mathbf{k}}_{12}|^2\ \sin(2\Delta^{\mathbf{k}}_{12}t) - |V^{\mathbf{k}}_{12}|^2\ \sin(2\Omega^{\mathbf{k}}_{12}t) + (|U^{\mathbf{k}}_{12}|^2\ - |V^{\mathbf{k}}_{13}|^2)\sin(2\Delta^{\mathbf{k}}_{23}t) \\
&+ \ (|V^{\mathbf{k}}_{12}|^2\ - |V^{\mathbf{k}}_{13}|^2)\sin(2\Omega^{\mathbf{k}}_{23}t)\ - |U^{\mathbf{k}}_{13}|^2\ \sin(2\Delta^{\mathbf{k}}_{13}t) + |V^{\mathbf{k}}_{13}|^2\ \sin(2\Omega^{\mathbf{k}}_{13}t)\Big] \\
&- \ \cos^2\theta_{13}\sin\theta_{13}\Big[\cos\delta\sin(2\theta_{12})\sin(2\theta_{23}) - 4\cos^2\theta_{12}\sin\theta_{13}\cos^2\theta_{23}\Big]\times \\
&\times \ \Big[|U^{\mathbf{k}}_{13}|^2\ \sin^2\left(\Delta^{\mathbf{k}}_{13}t\right) + |V^{\mathbf{k}}_{13}|^2\ \sin^2\left(\Omega^{\mathbf{k}}_{13}t\right)\Big] \\
&+ \ \cos^2\theta_{13}\sin\theta_{13}\Big[\cos\delta\sin(2\theta_{12})\sin(2\theta_{23}) + 4\sin^2\theta_{12}\sin\theta_{13}\cos^2\theta_{23}\Big]\times \\
&\times \ \Big[|U^{\mathbf{k}}_{23}|^2\ \sin^2\left(\Delta^{\mathbf{k}}_{23}t\right) + |V^{\mathbf{k}}_{23}|^2\ \sin^2\left(\Omega^{\mathbf{k}}_{23}t\right)\Big] \\
&+ \ \cos^2\theta_{13}\sin(2\theta_{12})\Big[(\sin^2\theta_{23} - \sin^2\theta_{13}\cos^2\theta_{23})\sin(2\theta_{12}) \\
&- \ \cos\delta\cos(2\theta_{12})\sin\theta_{13}\sin(2\theta_{23})\Big]\Big[|U^{\mathbf{k}}_{12}|^2\ \sin^2\left(\Delta^{\mathbf{k}}_{12}t\right) + |V^{\mathbf{k}}_{12}|^2\ \sin^2\left(\Omega^{\mathbf{k}}_{12}t\right)\Big]\,,
\end{aligned}
\tag{3.69}
$$

where we used the relations Eqs.(3.24)-(3.30). We also introduced the Jarlskog factor $J_{_{CP}}$ defined as [45]

$$
J_{_{CP}} \equiv Im(u_{i\alpha}u_{j\beta}u^*_{i\beta}u^*_{j\alpha}), \tag{3.70}
$$

where the u_{ij} are the elements of mixing matrix $\mathcal{U}$ and $i \neq j$, $\alpha \neq \beta$. In the parameterization Eq.(3.2), $J_{_{CP}}$ is given by

$$
J_{_{CP}} = \frac{1}{8}\sin\delta\ \sin(2\theta_{12})\sin(2\theta_{13})\cos\theta_{13}\sin(2\theta_{23}). \tag{3.71}
$$

Evidently, $J_{_{CP}}$ vanishes if $\theta_{ij} = 0, \pi/2$ and/or $\delta = 0, \pi$: all CP–violating effects are proportional to it.

The above oscillation formulas are exact. The differences with respect to the usual formulas for neutrino oscillations are in the energy dependence of the amplitudes and in the additional oscillating terms. For $|\mathbf{k}| \gg \sqrt{m_1 m_2}$, we have $|U^{\mathbf{k}}_{ij}|^2 \to 1$ and $|V^{\mathbf{k}}_{ij}|^2 \to 0$ and the traditional (Pontecorvo) oscillation formulas are approximately recovered. Indeed, for sufficiently small time arguments, a correction to the Pontecorvo formula is present even in the relativistic limit.

In Appendix G the oscillation formulas for the flavor charges on an initial electron anti-neutrino state are given.

We plot in Figures 3.2 and 3.4 the QFT oscillation formulas $\mathcal{Q}^{e}_{\mathbf{k},e}(t)$ and $\mathcal{Q}^{e}_{\mathbf{k},\mu}(t)$ as a function of time, and in Figures 3.3 and 3.5 the corresponding Pontecorvo oscillation formulas $P^{\mathbf{k}}_{e\to e}(t)$ and $P^{\mathbf{k}}_{e\to\mu}(t)$. The time scale is in T_{12} units, where $T_{12} = \pi/\Delta^{\mathbf{k}}_{12}$ is, for the values of parameters of Tab.(1), the largest oscillation period.



Figure 3.2: Plot of QFT oscillation formula: $\mathcal{Q}^e_{\mathbf{k},e}(t)$ in function of time for $k = 55$ and parameters as in Tab.(1).



Figure 3.3: Plot of QM oscillation formula: $P^{\mathbf{k}}_{e\to e}(t)$ in function of time for $k = 55$ and parameters as in Tab.(1).



Figure 3.4: Plot of QFT oscillation formula: $\mathcal{Q}^e_{\mathbf{k},\mu}(t)$ in function of time for $k = 55$ and parameters as in Tab(1).



Figure 3.5: Plot of QM oscillation formula: $P^{\mathbf{k}}_{e\rightarrow\mu}(t)$ in function of time for $k = 55$ and parameters as in Tab.(1).

3.6 CP and T violations in neutrino oscillations

In this Section we consider the oscillation induced CP and T violation in the context of the present QFT framework. Let us first briefly recall the situation in QM[3]: there, the CP asymmetry between the probabilities of two conjugate neutrino transitions, due to CPT invariance and unitarity of the mixing matrix, is given as [37]

$$\hat{\Delta}^{\rho\sigma}_{_{CP}}(t) \equiv P_{\nu_\sigma\to\nu_\rho}(t) - P_{\overline{\nu}_\sigma\to\overline{\nu}_\rho}(t), \tag{3.72}$$

where $\sigma, \rho = e, \mu, \tau$. The T violating asymmetry can be obtained in similar way as [37]

$$\hat{\Delta}^{\rho\sigma}_{_{T}}(t) \;\equiv\; P_{\nu_\sigma\to\nu_\rho}(t) - P_{\nu_\rho\to\nu_\sigma}(t) \;=\; P_{\nu_\sigma\to\nu_\rho}(t) - P_{\nu_\sigma\to\nu_\rho}(-t)\,. \tag{3.73}$$

The relationship $\hat{\Delta}^{\rho\sigma}_{_{CP}}(t) = \hat{\Delta}^{\rho\sigma}_{_{T}}(t)$ is a consequence of CPT invariance.

The corresponding quantities in QFT have to be defined in the framework of the previous Section, i.e. as expectation values of the flavor charges on states belonging to the flavor Hilbert space. We thus have for the CP violation:

$$\begin{aligned} \Delta^{\rho\sigma}_{_{CP}}(t) &\equiv \mathcal{Q}^{\rho}_{\mathbf{k},\sigma}(t) \,+\, \mathcal{Q}^{\bar{\rho}}_{\mathbf{k},\sigma}(t) \\ &= \left|\left\{\alpha^{r}_{\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},\rho}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},\rho}(0)\right\}\right|^2 - \\ &- \left|\left\{\alpha^{r\dagger}_{-\mathbf{k},\sigma}(t), \beta^{r\dagger}_{\mathbf{k},\rho}(0)\right\}\right|^2 - \left|\left\{\beta^{r}_{\mathbf{k},\sigma}(t), \beta^{r\dagger}_{\mathbf{k},\rho}(0)\right\}\right|^2 . \end{aligned} \tag{3.74}$$

We have

$$\sum_{\sigma} \Delta^{\rho\sigma}_{_{CP}} \;=\; 0 \qquad , \quad \rho, \sigma = e, \mu, \tau, \tag{3.75}$$

which follows from the fact that

$$\sum_{\sigma} Q_\sigma(t) = Q, \qquad \langle \nu_\rho | Q | \nu_\rho \rangle \;=\; 1 \quad \text{ and } \quad \langle \bar{\nu}_\rho | Q | \bar{\nu}_\rho \rangle \;=\; -1. \tag{3.76}$$

We can calculate the CP asymmetry Eq.(3.74) for a specific case, namely for the transition $\nu_e \longrightarrow \nu_\mu$. We obtain

$$\begin{aligned} \Delta^{e\mu}_{_{CP}}(t) \;&=\; 4J_{_{CP}}\Big[|U^{\mathbf{k}}_{12}|^2 \sin(2\Delta^{\mathbf{k}}_{12}t) - |V^{\mathbf{k}}_{12}|^2 \sin(2\Omega^{\mathbf{k}}_{12}t) + (|U^{\mathbf{k}}_{12}|^2 - |V^{\mathbf{k}}_{13}|^2)\sin(2\Delta^{\mathbf{k}}_{23}t) \\ &+(|V^{\mathbf{k}}_{12}|^2 - |V^{\mathbf{k}}_{13}|^2)\sin(2\Omega^{\mathbf{k}}_{23}t) - |U^{\mathbf{k}}_{13}|^2 \sin(2\Delta^{\mathbf{k}}_{13}t) + |V^{\mathbf{k}}_{13}|^2 \sin(2\Omega^{\mathbf{k}}_{13}t)\Big]\,, \end{aligned} \tag{3.77}$$

---

[3] We use here the ”hat” for QM quantities. For notational simplicity, we also suppress momentum indices where unnecessary.



Figure 3.6: Plot of the QFT CP asymmetry $\Delta^{e\mu}_{CP}(t)$, in function of time for $k = 55$ and parameters as in Tab.(1).

---

and $\Delta^{e\tau}_{CP}(t) = -\Delta^{e\mu}_{CP}(t)$. As already observed for oscillation formulas, high-frequency oscillating terms and Bogoliubov coefficients in the oscillation amplitudes appear in Eq.(3.77) as a QFT correction to the QM formula.

The definition of the QFT analogue of the T-violating quantity Eq.(3.73) is more delicate. Indeed, defining Δ_T as $\Delta^{e\mu}_T \equiv \mathcal{Q}^e_\mu(t) - \mathcal{Q}^\mu_e(t)$ does not seem to work, since we obtain $\Delta^{e\mu}_T - \Delta^{e\mu}_{CP} \neq 0$ in contrast with CPT conservation.

A more consistent definition of the time-reversal violation in QFT is then:

$$\Delta^{\rho\sigma}_T(t) \equiv \mathcal{Q}^\rho_{\mathbf{k},\sigma}(t) - \mathcal{Q}^\rho_{\mathbf{k},\sigma}(-t) \qquad , \quad \rho,\sigma = e,\mu,\tau\,. \tag{3.78}$$

With such definition, the equality $\Delta^{\rho\sigma}_T(t) = \Delta^{\rho\sigma}_{CP}(t)$ follows from $\mathcal{Q}^\rho_{\mathbf{k},\sigma}(-t) = -\mathcal{Q}^{\bar\rho}_{\mathbf{k},\sigma}(t)$.

We plot in Fig. 3.6 the CP asymmetry Eq.(3.77) for sample values of the parameters as in Tab.(1). In Fig. 3.7 the corresponding standard QM quantity is plotted for the same values of parameters.



Figure 3.7: Plot of the QM CP asymmetry $\hat{\Delta}^{e\mu}_{CP}(t)$, in function of time for $k = 55$ and parameters as in Tab.(1).

3.7 Conclusions

In this chapter, we have discussed the mixing of (Dirac) fermionic fields in Quantum Field Theory for the case of three flavors with CP violation. We constructed the flavor Hilbert space and studied the currents and charges for mixed fields (neutrinos). We will show in the next chapter that the algebraic structure associated with the mixing for the case of three generation turned out to be that of a deformed $su(3)$ algebra, when a CP violating phase is present.

We have derived all the known parameterization of the three-flavor mixing matrix and a number of new ones. We have shown that these parameterizations actually reflect the group theoretical structure of the generator of the mixing transformations.

By use of the flavor Hilbert space, we have calculated the exact QFT oscillation formulas, a generalization of the usual QM Pontecorvo formulas. The comparison between the exact oscillation formulas and the usual ones has been explicitly exhibited for sample values of the neutrino masses and mixings. CP and T violation induced by neutrino oscillations have also been discussed.

As already remarked in the Introduction, the corrections introduced by the present formalism to the usual Pontecorvo formulas are in principle experimentally testable. The fact that these corrections may be quantitatively below the experimental accuracy reachable at the present state of the art in the detection of the neutrino oscillations, does not justify neglecting them in the analysis of the particle mixing and oscillation mechanism. The exact oscillation formulas here derived are the result of a mathematically consistent analysis which cannot be ignored in a correct treatment of the field mixing phenomenon. As we have seen above, the QFT formalism accounts for all the known parameterizations of the mixing matrix and explains their origin and their reciprocal relations, thus unifying the phenomenological proposals scattered in the literature where such parameterizations have been presented. Moreover, the QFT formalism clearly points to the truly non-perturbative character of the particle mixing phenomenon. A lot of Physics must be there waiting to be discovered.

Chapter 4

Group theoretical aspects of neutrino mixing in Quantum Field Theory

We analyze some aspects of three flavor neutrino mixing. Particular emphasis is given to the related algebraic structures and their deformation in the presence of CP violation. A novel geometric phase related to CP violation is introduced.

4.1 Introduction

In this chapter we discuss the group structure involved in the mixing and the related representations, both for two- and three-flavor mixing. The deformation of the associated algebra as well as the geometric phase due to CP violation are also discussed [30, 35]. In Section 4.2 we study the deformation of SU(3) algebra induced by CP violation; in Section 4.3 we analyze the group representations, considering the case of two and three generations. The Section 4.4 is devoted to conclusions.

4.2 SU(3) deformed algebra

We investigate the algebraic structures associated with the mixing generator of three Dirac neutrinos fields Eq.(3.5). To this end, we consider the Lagrangian density describing three Dirac neutrinos fields:

$$\mathcal{L}(x) \;=\; \bar{\Psi}_m(x)\,(i\,\not\partial - \mathsf{M}_d)\,\Psi_m(x) \;=\; \bar{\Psi}_f(x)\,(i\,\not\partial - \mathsf{M})\,\Psi_f(x)\,, \tag{4.1}$$

where $\mathsf{M}_d = diag(m_1, m_2, m_3)$ and the matrix M is non-diagonal.

The above Lagrangian is invariant under global $U(1)$ phase transformations, leading to a conserved (total) charge $Q = \int d^3\mathbf{x}\,\Psi_m^\dagger(x)\,\Psi_m(x) \;=\; \int d^3\mathbf{x}\,\Psi_f^\dagger(x)\,\Psi_f(x)$.

We then study the invariance of $\mathcal{L}$ under global phase transformations of the kind:

$$\Psi'_m(x) \;=\; e^{i\alpha_j \tilde{F}_j}\, \Psi_m(x) \qquad , \qquad j = 1, 2, ..., 8. \tag{4.2}$$

where

$$\tilde{F}_j = \frac{1}{2}\tilde{\lambda}_j \quad , \quad j = 1, .., 8 \tag{4.3}$$

and the $\tilde{\lambda}_j$ are a generalization of the usual Gell-Mann matrices λ_j:

$$\tilde{\lambda}_1 = \begin{pmatrix} 0 & e^{i\delta_2} & 0 \\ e^{-i\delta_2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad , \qquad \tilde{\lambda}_2 = \begin{pmatrix} 0 & -ie^{i\delta_2} & 0 \\ ie^{-i\delta_2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\tilde{\lambda}_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad , \qquad \tilde{\lambda}_4 = \begin{pmatrix} 0 & 0 & e^{-i\delta_5} \\ 0 & 0 & 0 \\ e^{i\delta_5} & 0 & 0 \end{pmatrix}$$

$$\tilde{\lambda}_5 = \begin{pmatrix} 0 & 0 & -ie^{-i\delta_5} \\ 0 & 0 & 0 \\ ie^{i\delta_5} & 0 & 0 \end{pmatrix} \quad , \qquad \tilde{\lambda}_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & e^{i\delta_7} \\ 0 & e^{-i\delta_7} & 0 \end{pmatrix}$$

$$\tilde{\lambda}_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -ie^{i\delta_7} \\ 0 & ie^{-i\delta_7} & 0 \end{pmatrix} \quad , \qquad \tilde{\lambda}_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} . \tag{4.4}$$

These are normalized as the Gell-Mann matrices: $tr(\lambda_j \lambda_k) = 2\delta_{jk}$. One then obtains the following set of charges [35]:

$$\tilde{Q}_{m,j}(t) \;=\; \int d^3\mathbf{x}\, \Psi_m^\dagger(x)\, \tilde{F}_j\, \Psi_m(x) \;, \qquad j = 1, 2, ..., 8. \tag{4.5}$$

Thus the CKM matrix Eq.(3.2) is generated by $\tilde{Q}_{m,2}(t)$, $\tilde{Q}_{m,5}(t)$ and $\tilde{Q}_{m,7}(t)$, with $\{\delta_2, \delta_5, \delta_7\} \to \{0, \delta, 0\}$. An interesting point is that the algebra generated by the matrices Eqs.(4.4) *is not* $su(3)$ unless the condition $\Delta \equiv \delta_2 + \delta_5 + \delta_7 = 0$ is imposed: such a condition is however incompatible with the presence of a CP violating phase. In the parameterizations of the mixing matrices of Section 3.2 and Section 3.3 (cf. e.g. the discussion after Eq.(3.42)); we have the correspondence $\{\delta_2, \delta_5, \delta_7\} \leftrightarrow \{\delta_{12}, \delta_{13}, \delta_{23}\}$.

The $\tilde{F}_j$ satisfy a deformed $su(3)$ algebra with deformed commutation relations given by [30, 35]:

$$\begin{aligned}
[\tilde{F}_2,\tilde{F}_5] &= \frac{i}{2}\,\tilde{F}_7\,e^{-i\,\Delta\,(\tilde{F}_3-\sqrt{3}\tilde{F}_8)} \quad , & [\tilde{F}_1,\tilde{F}_4] &= \frac{i}{2}\,\tilde{F}_7\,e^{-i\,\Delta\,(\tilde{F}_3-\sqrt{3}\tilde{F}_8)} \quad , \\
[\tilde{F}_1,\tilde{F}_5] &= -\frac{i}{2}\,\tilde{F}_6\,e^{-i\,\Delta\,(\tilde{F}_3-\sqrt{3}\tilde{F}_8)} \quad , & [\tilde{F}_2,\tilde{F}_4] &= \frac{i}{2}\,\tilde{F}_6\,e^{-i\,\Delta\,(\tilde{F}_3-\sqrt{3}\tilde{F}_8)} \quad , \\
[\tilde{F}_2,\tilde{F}_7] &= -\frac{i}{2}\,\tilde{F}_5\,e^{-i\,\Delta\,(\tilde{F}_3+\sqrt{3}\tilde{F}_8)} \quad , & [\tilde{F}_1,\tilde{F}_7] &= -\,\frac{i}{2}\,\tilde{F}_4\,e^{-i\,\Delta\,(\tilde{F}_3+\sqrt{3}\tilde{F}_8)} \quad , \\
[\tilde{F}_1,\tilde{F}_6] &= \frac{i}{2}\,\tilde{F}_5\,e^{-i\,\Delta\,(\tilde{F}_3+\sqrt{3}\tilde{F}_8)} \quad , & [\tilde{F}_2,\tilde{F}_6] &= -\frac{i}{2}\,\tilde{F}_4\,e^{-i\,\Delta\,(\tilde{F}_3+\sqrt{3}\tilde{F}_8)} \quad , \\
[\tilde{F}_5,\tilde{F}_7] &= \frac{i}{2}\,\tilde{F}_2\,e^{2i\,\Delta\,\tilde{F}_3} \quad , & [\tilde{F}_4,\tilde{F}_7] &= \frac{i}{2}\,\tilde{F}_1\,e^{2i\,\Delta\,\tilde{F}_3} \quad , \\
[\tilde{F}_5,\tilde{F}_6] &= -\frac{i}{2}\,\tilde{F}_1\,e^{2i\,\Delta\,\tilde{F}_3} \quad , & [\tilde{F}_4,\tilde{F}_6] &= \frac{i}{2}\,\tilde{F}_2\,e^{2i\,\Delta\,\tilde{F}_3}, \qquad (4.6)
\end{aligned}$$

where $\Delta \equiv \delta_2+\delta_5+\delta_7$. The other commutators are the usual $su(3)$ ones. For $\Delta = 0$, the $su(3)$ algebra is recovered.

When CP violation is allowed, then $\Delta \neq 0$ and the $su(3)$ algebra is deformed. Let us introduce the raising and lowering operators, defined as [4]:

$$\tilde{T}_\pm \equiv \tilde{F}_1 \pm i\tilde{F}_2 \quad , \quad \tilde{U}_\pm \equiv \tilde{F}_6 \pm i\tilde{F}_7 \quad , \quad \tilde{V}_\pm \equiv \tilde{F}_4 \pm i\tilde{F}_5 \qquad (4.7)$$

We also define:

$$\tilde{Y} = \frac{2}{\sqrt{3}}\tilde{F}_8 \quad , \quad \tilde{T}_3 \equiv \tilde{F}_3 \quad , \quad \tilde{U}_3 \equiv \frac{1}{2}\left(\sqrt{3}\tilde{F}_8 - \tilde{F}_3\right) \quad , \quad \tilde{V}_3 \equiv \frac{1}{2}\left(\sqrt{3}\tilde{F}_8 + \tilde{F}_3\right) \qquad (4.8)$$

The commutation relations are

$$[\tilde{T}_3,\tilde{T}_\pm] = \pm\tilde{T}_\pm \quad , \quad [\tilde{T}_3,\tilde{U}_\pm] = \mp\frac{1}{2}\tilde{U}_\pm \quad , \quad [\tilde{T}_3,\tilde{V}_\pm] = \pm\frac{1}{2}\tilde{V}_\pm \quad , \quad [\tilde{T}_3,\tilde{Y}] = 0, \qquad (4.9)$$

$$[\tilde{Y},\tilde{T}_\pm] = 0 \quad , \quad [\tilde{Y},\tilde{U}_\pm] = \pm\tilde{U}_\pm \quad , \quad [\tilde{Y},\tilde{V}_\pm] = \pm\tilde{V}_\pm \quad , \qquad (4.10)$$

$$[\tilde{T}_+,\tilde{T}_-] = 2\tilde{T}_3 \quad , \quad [\tilde{U}_+,\tilde{U}_-] = 2\tilde{U}_3 \quad , \quad [\tilde{V}_+,\tilde{V}_-] = 2\tilde{V}_3 \quad , \qquad (4.11)$$

$$[\tilde{T}_+,\tilde{V}_+] = [\tilde{T}_+,\tilde{U}_-] = [\tilde{U}_+,\tilde{V}_+] = 0, \qquad (4.12)$$

that are similar to the standard SU(3) commutation relations. However, the following commutators are deformed:

$$[\tilde{T}_+,\tilde{V}_-] = -\tilde{U}_-\,e^{2i\Delta\tilde{U}_3} \quad , \quad [\tilde{T}_+,\tilde{U}_+] = \tilde{V}_+\,e^{-2i\Delta\tilde{V}_3} \quad , \quad [\tilde{U}_+,\tilde{V}_-] = \tilde{T}_-\,e^{2i\Delta\tilde{T}_3}. \qquad (4.13)$$

In a similar way with the above derivation, we can study the invariance properties of the Lagrangian Eq.(4.1) under the transformations:

$$\Psi'_f(x) = e^{i\alpha_j\tilde{F}_j}\,\Psi_f(x)\,, \qquad j = 1,2,...,8. \qquad (4.14)$$

Then the following charges are obtained

$$\tilde{Q}_{f,j}(t) \;=\; \int d^3\mathbf{x}\, \Psi_f^\dagger(x)\, \tilde{F}_j\, \Psi_f(x)\;, \qquad j = 1, 2, ..., 8. \tag{4.15}$$

4.3 Group representations and the oscillation formula

We now study the group representations. Let us first consider the simple case of two generations and then discuss the three flavor case.

4.3.1 Two flavors

In this case, the group is $SU(2)$ and the charges in the mass basis read:

$$Q_{m,j}(t) \;=\; \frac{1}{2}\int d^3\mathbf{x}\, \Psi_m^\dagger(x)\, \tau_j\, \Psi_m(x), \qquad j \;=\; 1, 2, 3, \tag{4.16}$$

where $\Psi_m^T = (\nu_1, \nu_2)$ and $\tau_j = \sigma_j/2$ with σ_j being the Pauli matrices.

The states with definite masses can then be defined as eigenstates of $Q_{m,3}$:

$$Q_{m,3}|\nu_1\rangle = \frac{1}{2}|\nu_1\rangle \qquad ; \qquad Q_{m,3}|\nu_2\rangle = -\frac{1}{2}|\nu_2\rangle \tag{4.17}$$

and similar ones for antiparticles, where

$$Q_{m,3} = \frac{1}{2}\int d^3\mathbf{x} \left[\nu_1^\dagger(x)\nu_1(x) - \nu_2^\dagger(x)\nu_2(x)\right]. \tag{4.18}$$



Figure 4.1: Plot of the mass doublet $SU(2)$.

We have $|\nu_i\rangle = \alpha_{\mathbf{k},i}^{r\dagger}|0\rangle_m$, $i = 1, 2$.

The Eq.(4.17) expresses the obvious fact that the mass eigenstates, treated as free particle states, are eigenstates of the conserved $U(1)$ charges associated to ν_1 and ν_2:

$$Q_1 \;\equiv\; \frac{1}{2}Q \,+\, Q_{m,3} \qquad ; \qquad Q_2 \;\equiv\; \frac{1}{2}Q \,-\, Q_{m,3}. \tag{4.19}$$

The next step is to define flavor states using a similar procedure. We need to be careful here since the diagonal $SU(2)$ generator $Q_{f,3}$ is time-dependent in the flavor basis. Thus we define states (Hilbert space) at a reference time $t = 0$ from:

$$Q_{f,3}(0)|\nu_e\rangle = \frac{1}{2}|\nu_e\rangle \quad ; \quad Q_{f,3}(0)|\nu_\mu\rangle = -\frac{1}{2}|\nu_\mu\rangle. \tag{4.20}$$

with $|\nu_\sigma\rangle = \alpha^{r\dagger}_{\mathbf{k},\sigma}(0)|0(0)\rangle_f$, $\sigma = e, \mu$ and similar ones for antiparticles.



Figure 4.2: Plot of the flavor doublet $SU(2)$.

The flavor states so defined are eigenstates of the flavor charges at time $t = 0$:

$$Q_e(t) = \frac{1}{2}Q + Q_{f,3}(t) \qquad ; \qquad Q_\mu(t) = \frac{1}{2}Q - Q_{f,3}(t), \tag{4.21}$$

$$Q_e(0)|\nu_e\rangle = |\nu_e\rangle \qquad ; \qquad Q_\mu(0)|\nu_\mu\rangle = |\nu_\mu\rangle. \tag{4.22}$$

and $Q_e(0)|\nu_\mu\rangle = Q_\mu(0)|\nu_e\rangle = 0$, with $Q_e(t)$ given by

$$\begin{aligned} Q_e(t) &= \int d^3\mathbf{x} \left[\nu_e^\dagger(x)\nu_e(x)\right] = \int d^3\mathbf{x}\Big[\nu_1^\dagger(x)\nu_1(x)\cos^2\theta + \\ &+ \nu_2^\dagger(x)\nu_2(x)\sin^2\theta + \Big(\nu_1^\dagger(x)\nu_2(x) + \nu_2^\dagger(x)\nu_1(x)\Big)\sin\theta\cos\theta\Big] \end{aligned} \tag{4.23}$$

and similar equation for $Q_\mu(t)$.

This result is far from being trivial since the usual Pontecorvo states [1]:

$$|\nu_e\rangle_P = \cos\theta\,|\nu_1\rangle + \sin\theta\,|\nu_2\rangle \tag{4.24}$$

$$|\nu_\mu\rangle_P = -\sin\theta\,|\nu_1\rangle + \cos\theta\,|\nu_2\rangle\,, \tag{4.25}$$

are *not* eigenstates of the flavor charges, indeed, for example, we have

$$\begin{aligned} &Q_e(0)|\nu_e\rangle_P = \cos^3\theta|\nu_1\rangle + \sin^3\theta|\nu_2\rangle + \sin\theta\cos\theta\,[U^*_{\mathbf{k}}|\nu_1\rangle\sin\theta + U_{\mathbf{k}}|\nu_2\rangle\cos\theta- \\ &-\sum_r\int d^3\mathbf{k}\left(\cos\theta u^{r\dagger}_{\mathbf{k},1}v^r_{-\mathbf{q},2}\beta^{r\dagger}_{-\mathbf{k},2}\alpha^{r\dagger}_{\mathbf{k},1}\alpha^{s\dagger}_{\mathbf{q},1}|0\rangle_{1,2} + \sin\theta u^{r\dagger}_{\mathbf{k},1}v^r_{-\mathbf{q},2}\beta^{r\dagger}_{-\mathbf{k},2}\alpha^{r\dagger}_{\mathbf{k},1}\alpha^{s\dagger}_{\mathbf{q},2}|0\rangle_{1,2}\right) - \\ &-\sum_r\int d^3\mathbf{k}\left(\cos\theta u^{r\dagger}_{\mathbf{k},2}v^s_{-\mathbf{q},1}\beta^{r\dagger}_{-\mathbf{k},1}\alpha^{r\dagger}_{\mathbf{k},2}\alpha^{s\dagger}_{\mathbf{q},1}|0\rangle_{1,2} + \sin\theta u^{r\dagger}_{\mathbf{k},2}v^s_{-\mathbf{q},1}\beta^{s\dagger}_{-\mathbf{k},1}\alpha^{r\dagger}_{\mathbf{k},2}\alpha^{s\dagger}_{\mathbf{q},2}|0\rangle_{1,2}\right). \end{aligned} \tag{4.26}$$

At a time $t \neq 0$, oscillation formulae can be derived for the flavor charges from the following relation

$$\begin{aligned}\langle \nu_e | Q_{f,3}(t) | \nu_e \rangle &= \frac{1}{2} - |U^{\mathbf{k}}_{12}|^2 \sin^2(2\theta) \sin^2\left(\frac{\omega_{k,2} - \omega_{k,1}}{2} t\right) - \\ &- |V^{\mathbf{k}}_{12}|^2 \sin^2(2\theta) \sin^2\left(\frac{\omega_{k,2} + \omega_{k,1}}{2} t\right) \qquad (4.27)\end{aligned}$$

where the non-standard oscillation term do appear and $Q_{f,3}(t)$ is

$$Q_{f,3}(t) = \frac{1}{2}[Q_e(t) - Q_\mu(t)]. \qquad (4.28)$$

4.3.2 Three flavors

Having discussed above the procedure for the definition of flavor states in the case of two flavors, we can directly write the *flavor charges* for three flavor mixed fields as

$$\begin{aligned} Q_e(t) &\equiv \frac{1}{3} Q + Q_{f,3}(t) + \frac{1}{\sqrt{3}} Q_{f,8}(t), \\ Q_\mu(t) &\equiv \frac{1}{3} Q - Q_{f,3}(t) + \frac{1}{\sqrt{3}} Q_{f,8}(t), \qquad (4.29) \\ Q_\tau(t) &\equiv \frac{1}{3} Q - \frac{2}{\sqrt{3}} Q_{f,8}(t). \end{aligned}$$

$$Q_\sigma(0)|\nu_\sigma\rangle = |\nu_\sigma\rangle \quad , \qquad Q_\sigma(0)|\bar{\nu}_\sigma\rangle = -|\bar{\nu}_\sigma\rangle \quad , \qquad \sigma = e, \mu, \tau, \qquad (4.30)$$

leading to

$$|\nu_\sigma\rangle \equiv \alpha^{r\dagger}_{\mathbf{k},\sigma}(0)|0(0)\rangle_f \quad , \qquad |\bar{\nu}_\sigma\rangle \equiv \beta^{r\dagger}_{\mathbf{k},\sigma}(0)|0(0)\rangle_f \quad , \qquad \sigma = e, \mu, \tau. \qquad (4.31)$$

These neutrino and antineutrino states can be related to the fundamental representation **3** and $\mathbf{3}^*$ of the (deformed) $SU(3)$ mixing group above introduced, as shown in Fig.4.3 for neutrinos.



Figure 4.3: Plot of the flavor triplet $SU(3)$.

To have the Fig.4.3, we have compared the generalized Gell-Mann–Nishijima relation that defines the charge operator as a function of $SU(3)$ generators:

$$q = \frac{1}{3}Q + (T_3 + \frac{Y}{2}), \tag{4.32}$$

to the Eqs.(4.29).

Note that the position of the points in the $\tilde{Y} - \tilde{T}_3$ is the same as for the ordinary $SU(3)$, since the diagonal matrices $\tilde{\lambda}_3$, $\tilde{\lambda}_8$ do not contain phases. However, a closed loop around the triangle gives a non-zero phase which is of geometrical origin [46] and only depends on the CP phase. A similar situation is valid for antineutrinos.

To see this more in detail, let us consider the octet representation as in Fig.4.4 and define the normalized state $|A\rangle$: $\langle A|A\rangle = 1$. Then all the other states are also normalized, except for $|G\rangle$: $|G\rangle = \frac{1}{\sqrt{2}}\tilde{T}_-|A\rangle$.



Figure 4.4: Plot of the $SU(3)$ octet.

We obtain the following paths

$$
\begin{aligned}
(AGBA): &\quad \tilde{V}_+\tilde{U}_-\tilde{T}_-|A\rangle = \tilde{V}_+([\tilde{U}_-,\tilde{T}_-]+\tilde{T}_-\tilde{U}_-)|A\rangle = \tilde{V}_+\tilde{V}_-e^{2i\Delta\tilde{V}_3}|A\rangle = e^{i\Delta}|A\rangle \\
(ABGA): &\quad \tilde{T}_+\tilde{U}_+\tilde{V}_-|A\rangle = e^{-i\Delta}|A\rangle \\
(AFGA): &\quad \tilde{T}_+\tilde{V}_-\tilde{U}_+|A\rangle = -e^{-i\Delta}|A\rangle \\
(AGFA): &\quad \tilde{U}_-\tilde{V}_+\tilde{T}_-|A\rangle = -e^{i\Delta}|A\rangle \\
(AFGBA): &\quad \tilde{V}_+\tilde{U}_-\tilde{V}_-\tilde{U}_+|A\rangle = |A\rangle \\
(AFEDCBA): &\quad \tilde{V}_+\tilde{T}_+\tilde{U}_-\tilde{V}_-\tilde{T}_-\tilde{U}_+|A\rangle = |A\rangle
\end{aligned}
\tag{4.33}
$$

where we have used

$$
\tilde{U}_-|A\rangle = \tilde{T}_+|A\rangle = \tilde{V}_+|A\rangle = 0, \quad \tilde{T}_3|A\rangle = |A\rangle\ , \quad \tilde{V}_3|A\rangle = \frac{1}{2}|A\rangle\ , \quad \tilde{U}_3|A\rangle = -\frac{1}{2}|A\rangle \tag{4.34}
$$

and the commutation relations.

We thus see that the phase sign change if we change the versus of the path on the triangles; the paths on two opposite triangles and around the hexagon bring no phase.

4.4 Conclusions

In this chapter, we have discussed some aspects of the quantization of mixed fermions (neutrinos) in the context of Quantum Field Theory.

In particular, we have analyzed the algebraic structures arising in connection with field mixing and their deformation due to the presence of CP violating phase, in the case of neutrino mixing among three generations.

We have defined flavor states in terms of the representations of the group associated with field mixing. A new geometric phase arising from CP violation was discovered.

Chapter 5

Mixing and oscillations of Majorana fermions

We study the mixing of Majorana fermions in Quantum Field Theory: Majorana field are treated in the case of mixing among two generations. We show how to consistently calculate oscillation formulas, which agree with previous results for Dirac fields.

5.1 Introduction

In the derivation of the oscillation formulas by use of the flavor Hilbert space a central role is played by the flavor charges [19] and indeed it was found that these operators satisfy very specifical physical requirements [21, 28]. However, these charges vanish identically in the case of neutral fields. In this chapter we will provide a consistent treatment of Majorana fermions. In order to keep the discussion transparent, we limit ourselves to the case of two generations.

Apart from the explicit quantization of the neutral mixed fields, the main point of this chapter is the study of the momentum operator (and more in general of the energy-momentum tensor) for those fields [41]. We show how to define it in a consistent way and by its use we then derive the oscillation formulas, which match the ones already obtained for Dirac fields. We also comment on its relevance for the study of Dirac mixed fields, where, when CP violation is present, the charge interpretation requires a further effort [29, 30, 31].

In Section 5.2 we treat Majorana fields. In Section 5.3 we discuss some general consequences of the results presented in this chapter and draw conclusions.

5.2 Majorana fermions

We consider the case of mixing of two Majorana fermion fields. The charge-conjugation operator C is defined as satisfying the relations

$$C^{-1}\gamma_\mu\, C = -\gamma_\mu^T \quad , \quad C^\dagger = C^{-1} \quad , \quad C^T = -C\ . \tag{5.1}$$

from which we define the charge conjugate ψ^c of ψ as

$$\psi^c(x) \equiv \gamma_0\, C\, \psi^*(x)\ . \tag{5.2}$$

Now we define a Majorana fermion as the field that satisfies the Dirac equation

$$(i\,\partial\!\!\!/ - m)\psi = 0 \tag{5.3}$$

and the self-conjugation relation

$$\psi = \psi^c\ . \tag{5.4}$$

Thus the two equations (5.3) and (5.4) ensure that the Majorana field is a neutral fermion field.

We now proceed by introducing the following Lagrangian:

$$\mathcal{L}(x) \;=\; \bar\psi_f(x)(i\,\partial\!\!\!/ - M)\psi_f(x) \;=\; \bar\psi_m(x)(i\,\partial\!\!\!/ - M_d)\psi_m(x)\,, \tag{5.5}$$

with $\psi_f^T = (\nu_e, \nu_\mu)$ being the flavor fields and $M = \begin{pmatrix} m_e & m_{e\mu} \\ m_{e\mu} & m_\mu \end{pmatrix}$. The flavor fields are connected to the free fields $\psi_m^T = (\nu_1, \nu_2)$ with $M_d = diag(m_1, m_2)$ by the rotation:

$$\nu_e(x) = \nu_1(x)\ \cos\theta + \nu_2(x)\ \sin\theta\ , \tag{5.6}$$

$$\nu_\mu(x) = -\nu_1(x)\ \sin\theta + \nu_2(x)\ \cos\theta\ . \tag{5.7}$$

The quantization of the free fields is given by [11]

$$\nu_i(x) = \sum_{r=1,2}\int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\, e^{i\mathbf{k}\cdot\mathbf{x}} \left[u^r_{\mathbf{k},i}(t)\alpha^r_{\mathbf{k},i} + v^r_{-\mathbf{k},i}(t)\alpha^{r\dagger}_{-\mathbf{k},i}\right], \quad i = 1, 2. \tag{5.8}$$

where $u^r_{\mathbf{k},i}(t) = e^{-i\omega_{k,i}t}u^r_{\mathbf{k},i}$, $v^r_{\mathbf{k},i}(t) = e^{i\omega_{k,i}t}v^r_{\mathbf{k},i}$, with $\omega_{k,i} = \sqrt{\mathbf{k}^2 + m_i^2}$. In order for the Majorana condition (5.4) to be satisfied, the four spinors must also satisfy the following condition:

$$v^s_{\mathbf{k},i} = \gamma_0\, C(u^s_{\mathbf{k},i})^* \quad ; \quad u^s_{\mathbf{k},i} = \gamma_0\, C(v^s_{\mathbf{k},i})^*\,. \tag{5.9}$$

The equal time anticommutation relations are:

$$\{\nu_i^\alpha(x), \nu_j^{\beta\dagger}(y)\}_{t=t'} = \delta^3(\mathbf{x}-\mathbf{y})\delta_{\alpha\beta}\delta_{ij} \quad , \quad \{\nu_i^\alpha(x), \nu_j^{\beta}(y)\}_{t=t'} = \delta^3(\mathbf{x}-\mathbf{y})(\gamma_0 C)^{\alpha\beta}\delta_{ij}, \tag{5.10}$$

with $\alpha, \beta = 1, .., 4$ and

$$\{\alpha_{\mathbf{k},i}^r, \alpha_{\mathbf{q},j}^{s\dagger}\} = \delta^3(\mathbf{k}-\mathbf{q})\delta_{rs}\delta_{ij} \, , \qquad i,j = 1,2 \, . \tag{5.11}$$

All other anticommutators are zero. The orthonormality and completeness relations are:

$$u_{\mathbf{k},i}^{r\dagger} u_{\mathbf{k},i}^s = v_{\mathbf{k},i}^{r\dagger} v_{\mathbf{k},i}^s = \delta_{rs} \, , \quad u_{\mathbf{k},i}^{r\dagger} v_{-\mathbf{k},i}^s = v_{-\mathbf{k},i}^{r\dagger} u_{\mathbf{k},i}^s = 0 \, , \quad \sum_{r=1,2} (u_{\mathbf{k},i}^r u_{\mathbf{k},i}^{r\dagger} + v_{-\mathbf{k},i}^r v_{-\mathbf{k},i}^{r\dagger}) = I \, . \tag{5.12}$$

We can recast Eqs.(5.6),(5.7) into the form:

$$\nu_e^\alpha(x) = G_\theta^{-1}(t) \; \nu_1^\alpha(x) \; G_\theta(t) \, , \tag{5.13}$$

$$\nu_\mu^\alpha(x) = G_\theta^{-1}(t) \; \nu_2^\alpha(x) \; G_\theta(t) \, , \tag{5.14}$$

where $G_\theta(t)$ is given by

$$G_\theta(t) = \exp\left[\frac{\theta}{2}\int d^3\mathbf{x} \left(\nu_1^\dagger(x)\nu_2(x) - \nu_2^\dagger(x)\nu_1(x)\right)\right] \, . \tag{5.15}$$

We have $G_\theta(t) = \prod_{\mathbf{k}} G_\theta^{\mathbf{k}}(t)$. Moreover for a given $\mathbf{k}$, in the reference frame where $\mathbf{k} = (0,0,|\mathbf{k}|)$, the spins decouple [25] and one has $G_\theta^{\mathbf{k}}(t) = \prod_r G_\theta^{\mathbf{k},r}(t)$ with

$$\begin{aligned} G_\theta^{\mathbf{k},r}(t) = \exp\Big\{\theta\Big[& U_{\mathbf{k}}^*(t) \; \alpha_{\mathbf{k},1}^{r\dagger}\alpha_{\mathbf{k},2}^r - U_{\mathbf{k}}(t) \; \alpha_{-\mathbf{k},2}^{r\dagger}\alpha_{-\mathbf{k},1}^r - \\ & -\epsilon^r V_{\mathbf{k}}^*(t)\alpha_{-\mathbf{k},1}^r\alpha_{\mathbf{k},2}^r + \epsilon^r V_{\mathbf{k}}(t)\alpha_{\mathbf{k},1}^{r\dagger}\alpha_{-\mathbf{k},2}^{r\dagger}\Big]\Big\} \, , \end{aligned} \tag{5.16}$$

where $U_{\mathbf{k}}(t)$ and $V_{\mathbf{k}}(t)$ are Bogoliubov coefficients given by

$$U_{\mathbf{k}}(t) \equiv |U_{\mathbf{k}}| \; e^{i(\omega_{k,2}-\omega_{k,1})t} \quad , \quad V_{\mathbf{k}}(t) \equiv |V_{\mathbf{k}}| \; e^{i(\omega_{k,2}+\omega_{k,1})t} \, , \tag{5.17}$$

$$|U_{\mathbf{k}}| \equiv \left(\frac{\omega_{k,1}+m_1}{2\omega_{k,1}}\right)^{\frac{1}{2}} \left(\frac{\omega_{k,2}+m_2}{2\omega_{k,2}}\right)^{\frac{1}{2}} \left(1 + \frac{|\mathbf{k}|^2}{(\omega_{k,1}+m_1)(\omega_{k,2}+m_2)}\right) , \tag{5.18}$$

$$|V_{\mathbf{k}}| \equiv \left(\frac{\omega_{k,1}+m_1}{2\omega_{k,1}}\right)^{\frac{1}{2}} \left(\frac{\omega_{k,2}+m_2}{2\omega_{k,2}}\right)^{\frac{1}{2}} \left(\frac{|\mathbf{k}|}{(\omega_{k,2}+m_2)} - \frac{|\mathbf{k}|}{(\omega_{k,1}+m_1)}\right) , \tag{5.19}$$

$$|U_{\mathbf{k}}|^2 + |V_{\mathbf{k}}|^2 = 1 \, . \tag{5.20}$$

The flavor fields can be thus expanded as:

$$\nu_\sigma(x) = \sum_{r=1,2} \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} e^{i\mathbf{k}\cdot\mathbf{x}} \left[u^r_{\mathbf{k},j}(t)\alpha^r_{\mathbf{k},\sigma}(t) + v^r_{-\mathbf{k},j}(t)\alpha^{r\dagger}_{-\mathbf{k},\sigma}(t) \right] , \tag{5.21}$$

with $\sigma, j = (e,1), (\mu,2)$ and the flavor annihilation operators given by (for $\mathbf{k} = (0,0,|\mathbf{k}|)$):

$$\begin{aligned} \alpha^r_{\mathbf{k},e}(t) &\equiv G^{-1}_\theta(t)\ \alpha^r_{\mathbf{k},1}\ G_\theta(t) = \cos\theta\,\alpha^r_{\mathbf{k},1} + \sin\theta\, \left(U^*_{\mathbf{k}}(t)\,\alpha^r_{\mathbf{k},2} + \epsilon^r V_{\mathbf{k}}(t)\,\alpha^{r\dagger}_{-\mathbf{k},2} \right) \ , \\ \alpha^r_{\mathbf{k},\mu}(t) &\equiv G^{-1}_\theta(t)\ \alpha^r_{\mathbf{k},2}\ G_\theta(t) = \cos\theta\,\alpha^r_{\mathbf{k},2} - \sin\theta\, \left(U_{\mathbf{k}}(t)\,\alpha^r_{\mathbf{k},1} - \epsilon^r V_{\mathbf{k}}(t)\,\alpha^{r\dagger}_{-\mathbf{k},1} \right) \ . \end{aligned} \tag{5.22}$$

We now consider the action of the generator of the mixing transformations on the vacuum $|0\rangle_{1,2}$. The flavor vacuum is defined as:

$$|0(\theta,t)\rangle_{e,\mu} \equiv G^{-1}_\theta(t)|0\rangle_{1,2} \ . \tag{5.23}$$

We define the state for a mixed particle with definite flavor, spin and momentum as:

$$|\alpha^r_{\mathbf{k},e}(t)\rangle \equiv \alpha^{r\dagger}_{\mathbf{k},e}(t)|0(t)\rangle_{e,\mu} = G^{-1}_\theta(t)\alpha^{r\dagger}_{\mathbf{k},1}|0\rangle_{1,2} \ . \tag{5.24}$$

The anticommutators of the flavor ladder operators at different times are:

$$\left\{ \alpha^r_{\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} = \cos^2\theta + \sin^2\theta \left(|U_{\mathbf{k}}|^2 e^{-i(\omega_2-\omega_1)(t-t')} + |V_{\mathbf{k}}|^2 e^{i(\omega_2+\omega_1)(t-t')} \right) , \tag{5.25}$$

$$\left\{ \alpha^{r\dagger}_{-\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} = \epsilon^r\ \sin^2\theta\,|U_{\mathbf{k}}||V_{\mathbf{k}}| \left(e^{i\omega_2(t-t')} - e^{-i\omega_2(t-t')} \right) e^{-i\omega_1(t+t')} , \tag{5.26}$$

$$\left\{ \alpha^r_{\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} = \cos\theta\sin\theta\,|U_{\mathbf{k}}| \left(e^{i(\omega_2-\omega_1)t'} - e^{i(\omega_2-\omega_1)t} \right) , \tag{5.27}$$

$$\left\{ \alpha^{r\dagger}_{-\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} = \epsilon^r\ \cos\theta\sin\theta\,|V_{\mathbf{k}}| \left(e^{-i(\omega_2+\omega_1)t'} - e^{-i(\omega_2+\omega_1)t} \right) . \tag{5.28}$$

The following quantity is constant in time:

$$\begin{aligned} &\left| \left\{ \alpha^r_{\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} \right|^2 + \left| \left\{ \alpha^{r\dagger}_{-\mathbf{k},e}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} \right|^2 + \\ &+ \left| \left\{ \alpha^r_{\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} \right|^2 + \left| \left\{ \alpha^{r\dagger}_{-\mathbf{k},\mu}(t), \alpha^{r\dagger}_{\mathbf{k},e}(t') \right\} \right|^2 = 1 \ . \end{aligned} \tag{5.29}$$

The corresponding of Eq.(5.29) for Dirac fields, was consistently interpreted as expressing the conservation of total charge. In the present case we are dealing with a neutral field and thus the charge operator vanishes identically. Nevertheless the quantities in Eq.(5.29) are well defined and are the Majorana field counterpart of the corresponding ones for the case of Dirac fields. Thus we look for a physical interpretation of such oscillating quantities.

Let us consider the momentum operator defined as

$$P^j \equiv \int d^3\mathbf{x}\, T^{0j}(x), \tag{5.30}$$

where the energy-momentum tensor for the fermion field, $T^{\mu\nu}$, is defined by

$$T^{\mu\nu} \equiv i\bar{\psi}\gamma^\nu\partial_\mu\psi. \tag{5.31}$$

For the free fields ψ_i we have:

$$\mathbf{P}_i = \int d^3\mathbf{x}\, \psi_i^\dagger(x)(-i\nabla)\psi_i(x) = \int d^3\mathbf{k} \sum_{r=1,2} \mathbf{k} \left(\alpha^{r\dagger}_{\mathbf{k},i}\alpha^r_{\mathbf{k},i} - \alpha^{r\dagger}_{-\mathbf{k},i}\alpha^r_{-\mathbf{k},i}\right), \quad i = 1, 2\ . \tag{5.32}$$

We then define the momentum operator for mixed fields:

$$\mathbf{P}_\sigma(t) = \int d^3\mathbf{x}\, \psi_\sigma^\dagger(x)(-i\nabla)\psi_\sigma(x) = \int d^3\mathbf{k} \sum_{r=1,2} \mathbf{k} \left(\alpha^{r\dagger}_{\mathbf{k},\sigma}(t)\alpha^r_{\mathbf{k},\sigma}(t) - \alpha^{r\dagger}_{-\mathbf{k},\sigma}(t)\alpha^r_{-\mathbf{k},\sigma}(t)\right), \tag{5.33}$$

with $\sigma = e, \mu$. We have

$$\mathbf{P}_\sigma(t) = G_\theta^{-1}(t)\mathbf{P}_i G_\theta(t) \tag{5.34}$$

and the conservation of total momentum as a consequence of

$$\mathbf{P}_e(t) + \mathbf{P}_\mu(t) = \mathbf{P}_1 + \mathbf{P}_2 \equiv \mathbf{P} \quad , \quad [\mathbf{P}, G_\theta(t)] = 0 \quad , \quad [\mathbf{P}, H] = 0\ . \tag{5.35}$$

We now consider the expectation values on the flavor state $|\alpha^r_{\mathbf{k},e}\rangle \equiv |\alpha^r_{\mathbf{k},e}(0)\rangle$. At time $t = 0$, this state is an eigenstate of the momentum operator $\mathbf{P}_e(0)$:

$$\mathbf{P}_e(0)\, |\alpha^r_{\mathbf{k},e}\rangle \;=\; \mathbf{k}\, |\alpha^r_{\mathbf{k},e}\rangle\ . \tag{5.36}$$

At $t \neq 0$ the expectation value for the momentum (normalized to initial value) gives:

$$\mathcal{P}^e_{\mathbf{k},\sigma}(t) \equiv \frac{\langle\alpha^r_{\mathbf{k},e}|\mathbf{P}_\sigma(t)|\alpha^r_{\mathbf{k},e}\rangle}{\langle\alpha^r_{\mathbf{k},e}|\mathbf{P}_\sigma(0)|\alpha^r_{\mathbf{k},e}\rangle} = \left|\left\{\alpha^r_{\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2 + \left|\left\{\alpha^{r\dagger}_{-\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},e}(0)\right\}\right|^2, \tag{5.37}$$

with $\sigma = e, \mu$, which is the same form of the expression one obtains for the expectation values of the flavor charges in the case of Dirac fields [20]. The flavor vacuum expectation value of the momentum operator $\mathbf{P}_\sigma(t)$ vanishes at all times:

$$ {}_{e,\mu}\langle 0|\mathbf{P}_\sigma(t)|0\rangle_{e,\mu} = 0 \quad , \quad \sigma = e, \mu \; . \tag{5.38}$$

The explicit calculation of the oscillating quantities $\mathcal{P}^e_{\mathbf{k},\sigma}(t)$ gives:

$$\mathcal{P}^e_{\mathbf{k},e}(t) \;=\; 1 - \sin^2 2\theta \left[|U_{\mathbf{k}}|^2 \sin^2\left(\frac{\omega_{k,2} - \omega_{k,1}}{2} t\right) + |V_{\mathbf{k}}|^2 \sin^2\left(\frac{\omega_{k,2} + \omega_{k,1}}{2} t\right)\right] , \tag{5.39}$$

$$\mathcal{P}^e_{\mathbf{k},\mu}(t) \;=\; \sin^2 2\theta \left[|U_{\mathbf{k}}|^2 \sin^2\left(\frac{\omega_{k,2} - \omega_{k,1}}{2} t\right) + |V_{\mathbf{k}}|^2 \sin^2\left(\frac{\omega_{k,2} + \omega_{k,1}}{2} t\right)\right] , \tag{5.40}$$

in complete agreement with the Dirac field case [20].

5.3 Discussion and Conclusions

In this chapter, we have studied the mixing among two generations in the case of Majorana in the context of Quantum Field Theory.

The main point is the calculation of oscillation formulas, which we obtained by use of the momentum operator, that is well defined for (mixed) neutral fields, whereas the charge operator vanishes identically. The results confirm the oscillation formulas already obtained in the case of Dirac fields by use of the flavor charges, and it also reveals to be useful in the case of three flavor mixing, where the presence of the CP violating phase introduces ambiguities in the treatment based on flavor charges [30].

It is indeed interesting to comment on this point: for Dirac fields, the momentum operator is given as

$$\begin{aligned}\mathbf{P}_\sigma(t) = \int d^3\mathbf{k} \sum_r \frac{\mathbf{k}}{2}\Big(&\alpha^{r\dagger}_{\mathbf{k},\sigma}(t)\alpha^r_{\mathbf{k},\sigma}(t) - \alpha^{r\dagger}_{-\mathbf{k},\sigma}(t)\alpha^r_{-\mathbf{k},\sigma}(t) + \\ &+\beta^{r\dagger}_{\mathbf{k},\sigma}(t)\beta^r_{\mathbf{k},\sigma}(t) - \beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\beta^r_{-\mathbf{k},\sigma}(t)\Big) \quad , \quad \sigma = e, \mu, \tau \; .\end{aligned} \tag{5.41}$$

This operator can be used for the calculation of the oscillation formulas for Dirac neutrinos in analogy with what done above in the Majorana case. Although in this case the charge operator is available and it has been used successfully for deriving the oscillation formula [20] in the two-flavor case, it has emerged that for three-flavor mixing, the CP violating phase introduces complications in the identification of the observables and indeed the matter is still object of discussion [29, 30, 31]. The main problem there is that the flavor charges at time t do not annihilate the flavor vacuum:

${}_f\langle 0|Q_\sigma(t)|0\rangle_f \neq 0$ and this expectation value needs to be subtracted by hand in order to get the correct oscillation formulas [30].

However, we see easily how the use of the momentum operator confirms the results of Ref.[30], without presenting any ambiguity. We have indeed:

$$
\begin{aligned}
{}_f\langle 0|\mathbf{P}_\sigma(t)|0\rangle_f \;&=\; 0 && (5.42)\\
\frac{\langle\nu_\rho|\mathbf{P}_\sigma(t)|\nu_\rho\rangle}{\langle\nu_\rho|\mathbf{P}_\sigma(0)|\nu_\rho\rangle} \;&=\; \left|\left\{\alpha^r_{\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},\rho}(0)\right\}\right|^2 + \left|\left\{\beta^{r\dagger}_{-\mathbf{k},\sigma}(t), \alpha^{r\dagger}_{\mathbf{k},\rho}(0)\right\}\right|^2 , && (5.43)
\end{aligned}
$$

with $\sigma, \rho = e, \mu, \tau$ and $|\nu_\rho\rangle \equiv \alpha^{r\dagger}_{\mathbf{k},\rho}(0)|0\rangle_f$. This follows from the following relations:

$$
\begin{aligned}
{}_f\langle 0|\alpha^{r\dagger}_{\mathbf{k},\sigma}(t)\alpha^r_{\mathbf{k},\sigma}(t)|0\rangle_f \;&=\; {}_f\langle 0|\alpha^{r\dagger}_{-\mathbf{k},\sigma}(t)\alpha^r_{-\mathbf{k},\sigma}(t)|0\rangle_f && (5.44)\\
{}_f\langle 0|\beta^{r\dagger}_{\mathbf{k},\sigma}(t)\beta^r_{\mathbf{k},\sigma}(t)|0\rangle_f \;&=\; {}_f\langle 0|\beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\beta^r_{-\mathbf{k},\sigma}(t)|0\rangle_f && (5.45)
\end{aligned}
$$

which are valid even in presence of CP violation, when ${}_f\langle 0|\alpha^{r\dagger}_{\mathbf{k},\sigma}(t)\alpha^r_{\mathbf{k},\sigma}(t)|0\rangle_f \neq {}_f\langle 0|\beta^{r\dagger}_{-\mathbf{k},\sigma}(t)\beta^r_{-\mathbf{k},\sigma}(t)|0\rangle_f$.

These results seem to suggest that perhaps a redefinition of the flavor charge operators is necessary in presence of CP violation and further study in this direction is in progress.

Chapter 6

Neutrino mixing and cosmological constant

We show that the non–perturbative vacuum structure associated with neutrino mixing leads to a non–zero contribution to the value of the cosmological constant. Such a contribution comes from the specific nature of the mixing phenomenon. Its origin is completely different from the one of the ordinary contribution of a massive spinor field. We estimate this neutrino mixing contribution by using the natural cut–off appearing in the quantum field theory formalism for neutrino mixing and oscillation.

6.1 Introduction

By resorting to the recent discovery of the unitary inequivalence between the mass and the flavor vacua for neutrino fields in quantum field theory (QFT) [20, 21, 25, 26, 27, 28, 29, 31, 33, 42, 43], we show that the non–perturbative vacuum structure associated with neutrino mixing may lead to a non–zero contribution to the value of the cosmological constant [34].

The contribution we find comes from the specific nature of the field mixing and is therefore of different origin with respect to the ordinary well known perturbative vacuum energy contribution of a massive spinor field.

The nature of the cosmological constant, say Λ, is one of the most intriguing issues in modern theoretical physics and cosmology. Data coming from observations indicate that not only Λ is different from zero, but it also dominates the universe dynamics driving an accelerated expansion (see for example [47]).

In the classical framework, Λ can be considered as an intrinsic (i.e. not induced by matter) curvature of space-time or a sort of shift in the matter Lagrangian. In the latter case Λ can be considered as an unclustered, non interacting component of the cosmic fluid with a constant energy density ρ and the equation of state $p = -\rho$. These properties and the fact that the presence of a cosmological constant fluid has to be compatible with the structure formation, allow to set the upper bound $\Lambda < 10^{-56} cm^{-2}$ [48].

In the quantum framework, the standard approach is to consider the cosmological constant as a gravitational effect of vacuum energy [49, 50, 51]. A common problem of all these approaches is that they do not provide a value of Λ in the bound given above. This is known as the *cosmological constant problem* [51] and it has been tried to address it in different ways (see for example [52]).

In this chapter, we show that the vacuum energy induced by the neutrino mixing may contribute to the value of cosmological constant in a fundamentally different way from the usual zero-point energy contribution, as already stated above.

Indeed, it has been realized [20, 21, 25] that the mixing of massive neutrino fields is a highly non-trivial transformation in QFT. The vacuum for neutrinos with definite mass is not invariant under the mixing transformation and in the infinite volume limit it is unitarily inequivalent to the vacuum for the neutrino fields with definite flavor number. This affects the oscillation formula which turns out to be different from the usual Pontecorvo formula [1] and a number of consequences have been already discussed [21].

The existence of the two inequivalent vacua for the flavor and the mass eigenstate neutrino fields, respectively, is crucial in order to obtain a non–zero contribution to the cosmological constant as we show below.

In Section 6.2, we show that the neutrino contribution to the value of the cosmological constant is non–zero and then we estimate its value by using the natural scale of neutrino mixing as cut–off. The result turns out to be compatible with the above mentioned upper bound on Λ. Section 6.3 is devoted to the conclusions. The Appendix J is devoted to the tetradic formalism in the Friedmann Robertson Walker (FRW) space–time.

6.2 Neutrino mixing contribution to the cosmological constant

The connection between the vacuum energy density $\langle\rho_{vac}\rangle$ and the cosmological constant Λ is provided by the well known relation

$$\langle\rho_{vac}\rangle = \frac{\Lambda}{4\pi G}, \tag{6.1}$$

where G is the gravitational constant.

In order to calculate $\langle\rho_{vac}\rangle$ we have to consider tetrads and spinorial connection. The symmetries of the cosmological metric make this task easier.

Tetrads are defined as a local inertial coordinate system at every space-time point defined by

$$g_{\mu\nu} = e^a_\mu \, e^b_\nu \, \eta_{ab}, \tag{6.2}$$

where $g_{\mu\nu}$ it the curved space-time metric and η_{ab} is the Minkowski metric. In the tetrads framework the parallel transport in a torsion free space-time is defined by the affine spin connection one form

$$\omega^a_b = \omega^a_{b\mu} dx^\mu, \tag{6.3}$$

which satisfies the Cartan Structure equations

$$de^a + \omega^a_b \wedge e^b = 0, \tag{6.4}$$

where e^a is the tetrad one-form defined by

$$e^a = e^a_\nu dx^\nu. \tag{6.5}$$

The energy–momentum tensor density $\mathcal{T}_{\mu\nu}$ is obtained by varying the action with respect to the metric $g_{\mu\nu}$:

$$\mathcal{T}_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta S}{\delta g^{\mu\nu}(x)}, \tag{6.6}$$

where the action is

$$S = \int \sqrt{-g} \mathcal{L}(x) d^4x. \tag{6.7}$$

In the present case, the energy momentum tensor density is given by

$$\mathcal{T}_{\mu\nu}(x) = \frac{i}{2} \left(\bar{\Psi}_m(x) \gamma_\mu \overleftrightarrow{D}_\nu \Psi_m(x) \right) \; , \tag{6.8}$$

where $\overleftrightarrow{D}_\nu$ is the covariant derivative:

$$D_\nu = \partial_\nu + \Gamma_\nu, \qquad \Gamma_\nu = \frac{1}{8} \omega^{ab}_\nu [\gamma_a, \gamma_b], \qquad \gamma_\mu(x) = \gamma^c e_{c\mu}(x), \tag{6.9}$$

being γ^c the standard Dirac matrices, and $\bar{\Psi} \overleftrightarrow{D}_\nu \Psi = \bar{\Psi} D \Psi - (D\bar{\Psi})\Psi$. Let us consider the FRW metric of the form

$$ds^2 = dt^2 - a^2(t) \left(\frac{dr^2}{\rho} + r^2 d\theta^2 + r^2 \sin^2(\theta) d\phi^2 \right), \tag{6.10}$$

with $\rho = 1 - \kappa r^2$, and $\kappa = 0, 1, -1$.

In the FRW metric Eq.(6.10), the most natural choice for tetrads is

$$\begin{aligned} e^0 &= e^0_0\, dx^0 = dt, & e^1 = e^1_1\, dx^1 = \frac{a(t)}{\rho}\, dr, \\ e^2 &= e^2_2\, dx^2 = a(t)\; r\; d\theta, & e^3 = e^3_3\, dx^3 = a(t)\; r\; \sin(\theta) dr. \end{aligned} \tag{6.11}$$

Using Eq.(6.4) and the definitions in Eq.(6.9) we have (see the Appendix J)

$$D_0 = \partial_0 + \frac{1}{4}[\gamma_i, \gamma_0]\, \mathcal{H}\, e^i_0 + \frac{1}{4}[\gamma_1, \gamma_j]\, \frac{\rho}{ar}\, e^j_0 + \frac{1}{4}[\gamma_2, \gamma_3]\, \frac{\tan(\theta)}{ar}\, e^3_0, \tag{6.12}$$

with $i = 1, 2, 3$, $j = 2, 3$.

Since we choose diagonal tetrads every term e^k_0 with $k = 1, 2, 3$ is null. This implies that all the terms but the first are null whatever the value of the commutators of the Dirac matrices. It is worth to stress that this result is independent of the choice of tetrads because the (0,0) component of the energy momentum tensor of a fluid is equivalent to the energy density only if the tetrads (or in general the chosen coordinates) are time-orthogonal as in Eq.(6.11). If we choose non time-orthogonal tetrads, i.e. e^0_0 not constant, $\mathcal{T}_{00}(x)$ does not represent the energy density because it acquires "pressure components" due to the different orientation of the tetrad. In our calculation these terms are the second, third and forth term of the Eq.(6.12).

Thus the temporal component of the spinorial derivative in the FRW metric is just the standard time derivative [34]:

$$D_0 = \partial_0. \tag{6.13}$$

This is not surprising if we consider the symmetries of the metric element Eq.(6.10). Thus, the (0,0) component of the stress energy tensor density is

$$\mathcal{T}_{00} = \mathcal{T}^{Flat}_{00}. \tag{6.14}$$

This allows us to use $\mathcal{T}^{Flat}_{00}$ to compute the cosmological constant. From Eq.(6.8) we thus obtain

$$\mathcal{T}_{00}(x) = \frac{i}{2} : \Big(\bar{\Psi}_m(x)\gamma_0 \overleftrightarrow{\partial}_0 \Psi_m(x)\Big) : \tag{6.15}$$

where : ... : denotes the customary normal ordering with respect to the mass vacuum in the flat space-time.

In terms of the annihilation and creation operators of fields ν_1 and ν_2, the energy-momentum tensor

$$T_{00} = \int d^3x \mathcal{T}_{00}(x) \tag{6.16}$$

is given by

$$T_{00}^{(i)} = \sum_r \int d^3\mathbf{k}\, \omega_{k,i} \left(\alpha_{\mathbf{k},i}^{r\dagger} \alpha_{\mathbf{k},i}^{r} + \beta_{-\mathbf{k},i}^{r\dagger} \beta_{-\mathbf{k},i}^{r} \right), \qquad i = 1, 2, \tag{6.17}$$

and we note that $T_{00}^{(i)}$ is time independent.

Next, our task is to compute the expectation value of $T_{00}^{(i)}$ in the flavor vacuum $|0\rangle_f$, which, as already recalled, is the one relevant to mixing and oscillations. Thus, the contribution $\langle \rho_{vac}^{mix} \rangle$ of the neutrino mixing to the vacuum energy density is:

$${}_f\langle 0| \sum_i T_{00}^{(i)}(0) |0\rangle_f = \langle \rho_{vac}^{mix} \rangle \eta_{00} \ . \tag{6.18}$$

We observe that within the above QFT formalism for neutrino mixing we have

$${}_f\langle 0| T_{00}^{(i)} |0\rangle_f = {}_f\langle 0(t)| T_{00}^{(i)} |0(t)\rangle_f \tag{6.19}$$

for any t. We then obtain

$${}_f\langle 0| \sum_i T_{00}^{(i)}(0) |0\rangle_f = \sum_{i,r} \int d^3\mathbf{k}\, \omega_{k,i} \left({}_f\langle 0| \alpha_{\mathbf{k},i}^{r\dagger} \alpha_{\mathbf{k},i}^{r} |0\rangle_f + {}_f\langle 0| \beta_{\mathbf{k},i}^{r\dagger} \beta_{\mathbf{k},i}^{r} |0\rangle_f \right). \tag{6.20}$$

For simplicity we restrict ourselves to the two flavor mixing and we use Dirac neutrino fields. Since [25]

$${}_f\langle 0| \alpha_{\mathbf{k},i}^{r\dagger} \alpha_{\mathbf{k},i}^{r} |0\rangle_f = {}_f\langle 0| \beta_{\mathbf{k},i}^{r\dagger} \beta_{\mathbf{k},i}^{r} |0\rangle_f \ = \sin^2\theta |V_{\mathbf{k}}|^2, \tag{6.21}$$

we get

$${}_f\langle 0| \sum_i T_{00}^{(i)}(0) |0\rangle_f = \ 8 \sin^2\theta \int d^3\mathbf{k}\, (\omega_{k,1} + \omega_{k,2})\, |V_{\mathbf{k}}|^2 = \langle \rho_{vac}^{mix} \rangle \eta_{00}, \tag{6.22}$$

i.e.

$$\langle \rho_{vac}^{mix} \rangle = 32 \pi^2 \sin^2\theta \int_0^K dk\, k^2 (\omega_{k,1} + \omega_{k,2}) |V_{\mathbf{k}}|^2, \tag{6.23}$$

where the cut-off K has been introduced. Eq.(6.23) is our result: it shows that the cosmological constant gets a non-zero contribution induced from the neutrino mixing [34]. Notice that such a contribution is indeed zero in the no-mixing limit when the mixing angle $\theta = 0$ and/or $m_1 = m_2$. It is to be remarked that the contribution is zero also in the limit of $V_{\mathbf{k}} \to 0$, namely in the limit of the traditional phenomenological mixing treatment.

It is interesting to note that, for high momenta, the function $|V_{\mathbf{k}}|^2$ produces a drastic decrease in the degree of divergency of the above integral, in comparison with the case of a free field. Thus, if for example we chose $K \gg \sqrt{m_1 m_2}$, we obtain:

$$\langle \rho_{vac}^{mix} \rangle \propto \sin^2\theta \, (m_2 - m_1)^2 \, K^2, \tag{6.24}$$

whereas the usual zero-point energy contribution would be going like K^4.

Of course, we are not in a position to make our result independent on the cut-off choice. However, this was not our goal. What we have shown is that a non - zero contribution to the value of the cosmological constant may come from the mixing of the neutrinos. We have not solved the cosmological constant problem. Although it might be unsatisfactory from a general theoretical point of view, we may try to estimate the neutrino mixing contribution by making our choice for the cut-off. Since we are dealing with neutrino mixing, at a first trial it might be reasonable to chose the cut-off proportional to the natural scale we have in the mixing phenomenon, namely $\mathbf{k}_0^2 \simeq m_1 m_2$ [25].

With such a choice, using $K \sim k_0$, $m_1 = 7 \times 10^{-3} eV$, $m_2 = 5 \times 10^{-2} eV$, $k_0 = 10^{-3} eV$ and $\sin^2\theta \simeq 1$ in Eq.(6.23), we obtain

$$\langle \rho_{vac}^{mix} \rangle = 1.3 \times 10^{-47} GeV^4 \tag{6.25}$$

Using Eq.(6.1), we have agreement with the upper bound given in Section 6.1:

$$\Lambda \sim 10^{-56} cm^{-2}, \tag{6.26}$$

Another possible choice is to use the electro-weak scale cut-off: $K \approx 100 GeV$. We then have

$$\langle \rho_{vac}^{mix} \rangle = 1.5 \times 10^{-15} GeV^4 \tag{6.27}$$

and

$$\Lambda \sim 10^{-24} cm^{-2}, \tag{6.28}$$

which is, however, beyond the accepted upper bound.

6.3 Conclusions

In this chapter we have shown that the neutrino mixing can give rise to a non zero contribution to the cosmological constant [34]. We have shown that this contribution is of a different nature with respect to that given by the zero-point energy of free fields and we estimated it by using the cut–off given by the natural scale of the neutrino mixing phenomenon. The different origin of the mixing contribution also manifests in the different ultraviolet divergency order (quadratic rather than quartic, see Eq.(6.24)). The obtained value is consistent with the accepted upper bound for the value of Λ. On the contrary, by using as a cut off the one related with the electroweak scale, the Λ value is greater than such upper bound.

It is worth to stress once more that the origin of the present contribution is completely different from that of the ordinary contribution to the vacuum zero energy of a massive spinor field. As we have shown, the effect we find is not originated from a radiative correction at some perturbative order [53]. Our effect is exact at any order. It comes from the property of QFT of being endowed with infinitely many representations of the canonical (anti-)commutation relations in the infinite volume limit. Therefore, the new result we find is that it is the *mixing* phenomenon which provides such a vacuum energy contribution, and this is so since the field mixing involves unitary inequivalent representations. Indeed, as Eqs.(6.21)-(6.23) show the contribution vanishes as $V_{\mathbf{k}} \to 0$, namely in the quantum mechanical limit where the representations of the (anti-)commutation relations are all each other unitarily equivalent. Our result thus discloses a new possible mechanism contributing to the cosmological constant value.

As a final consideration, we observe that this effect could also be exploited in the issue of dark energy without introducing exotic fields like quintessence. In fact, neutrinos constitute a cosmic background of unclustered components whose mixing and oscillations could drive the observed accelerated expansion.

PART 2. BOSON MIXING

Chapter 7

Boson mixing in Quantum Mechanics

In this chapter we present the theoretical model describing the meson mixing and oscillations in Quantum Mechanics. In particular, the boson oscillations will be treated by using the Gell-Mann and Pais formalism.

7.1 Usual representation of boson oscillations: the Gell-Mann Pais model

Quark mixing and meson mixing are widely accepted and verified [4]. However, many features of the physics of mixing are still obscure, for example the issue related to its origin in the context of Standard Model and the related problem of the generation of masses.

The problem of the boson mixing is known since 1955 when Gell-Mann and Pais predicted the existence of two neutral kaons [54] : K^0 of strangeness $S = 1$ and $\bar{K}^0$ of strangeness $S = -1$. These are particle and antiparticle, and are connected by the process of charge conjugation, which involves a reversal of values of I_3 and a change of strangeness $\Delta S = 2$. Strong interactions conserve I_3 and S, so that as far as production is concerned, the separate neutral-kaon eigenstates are $K^0 - \bar{K}^0$.

Now suppose K^0 and $\bar{K}^0$ particles propagate through empty space. Since both are neutral, both can decay to pions by weak interaction, with $\Delta S = 1$. Thus, mixing can occur via (virtual) intermediate pion states:

$$\begin{aligned} &K^0 \leftrightarrows 2\pi \leftrightarrows \bar{K}^0 \\ &K^0 \leftrightarrows 3\pi \leftrightarrows \bar{K}^0 \end{aligned} \tag{7.1}$$

These transitions are $\Delta S = 2$ and thus second order weak interactions. Although extremely weak, this implies that if one has a pure K^0-state at $t = 0$, at any later time $t > 0$ one will have a superposition of both K^0 and $\bar{K}^0$, so that the state can be written

$$|K(t)\rangle = \alpha(t)|K^0\rangle + \beta(t)|\bar{K}^0\rangle. \tag{7.2}$$

The phenomena has been explained by realizing that what we observe is the mixture of two mass and mean life eigenstates K_S and K_L expressed by

$$\begin{aligned} |K_S\rangle &= \frac{1}{\sqrt{2(1+|\varepsilon|^2)}}[(1+\varepsilon)|K^0\rangle + (1-\varepsilon)|\bar{K}^0\rangle] = \frac{1}{\sqrt{(1+|\varepsilon|^2)}}(|K_1\rangle + \varepsilon|K_2\rangle), \\ |K_L\rangle &= \frac{1}{\sqrt{2(1+|\varepsilon|^2)}}[(1+\varepsilon)|K^0\rangle - (1-\varepsilon)|\bar{K}^0\rangle] = \frac{1}{\sqrt{(1+|\varepsilon|^2)}}(|K_2\rangle + \varepsilon|K_1\rangle), \end{aligned} \tag{7.3}$$

where ε is a small, complex parameter responsible for CP symmetry breaking and K_1, K_2 are CP eigenstates:

$$\begin{aligned} |K_1\rangle &= \frac{1}{\sqrt{2}}(|K^0\rangle + |\bar{K}^0\rangle) \qquad && CP|K_1\rangle = |K_1\rangle, \\ |K_2\rangle &= \frac{1}{\sqrt{2}}(|K^0\rangle - |\bar{K}^0\rangle) \qquad && CP|K_2\rangle = -|K_2\rangle \end{aligned} \tag{7.4}$$

with the convention

$$\begin{aligned} CP|K^0\rangle &= |\bar{K}^0\rangle \\ CP|\bar{K}^0\rangle &= |K^0\rangle. \end{aligned} \tag{7.5}$$

Unlike K^0 and $\bar{K}^0$, distinguished by their mode of production (the K^0 can be produced by nonstrange particles in association with a hyperon and $\bar{K}^0$ can be produced only in association with a kaon or antihyperon, of strangeness $S = 1$), K_S and K_L are distinguished by their mode of decay. Consider 2π and 3π decay modes.

Since pions have no spin, angular momentum conservation requires that the two pions resulting from $K^0 \longrightarrow 2\pi$ decay carry relative angular momentum equal to the spin of the kaon. A neutral 2π state with specified angular momentum l is an eigenstate of C with eigenvalue $C = (-1)^l$, since the action of C is just to exchange the two pions.

Being K^0 a spinless particle, Gell-Mann e Pais concluded that only the component K_1, CP eigenstate with eigenvalue 1, would be capable of 2π decay; while K_2 would only decay in 3π state with $CP = -1$. But, in 1964 Christenson, Cronin, Fitch and Turlay [2] demonstrate that the K_2 state could also decay to $\pi^+\pi^-$ with a branching ratio of order 10^{-3}. Then there is a CP violation in the K^0 decay and the physical component of K^0 are K_S (short lived component) and K_L (long lived component), where K_S consists principally of a $CP = +1$ amplitude, but with a little $CP = -1$, and K_L vice versa. Experimentally the mean lives of K_S and K_L are $\tau_S = (0.8934 \mp 0.0008) \times 10^{-10}\,\text{sec}$, $\tau_L = (5.17 \mp 0.04) \times 10^{-8}\,\text{sec}$ respectively.

An important phenomenon is the K^0 regeneration [55]. Suppose we produce a pure K^0 beam and let it travel in vacuo for the order of $100K_S$ mean lives, so that all the K_S component has decayed and we are left with K_L only. Now let the K_L beam traverse a slab of material and interact. Immediately, the strong interactions will pick out the strangeness $S = +1$ and $S = -1$ components of the beam.

Thus, of the original K^0 beam intensity, about 50% has disappeared by K_S decay. The remainder, K_L upon traversing a slab where its nuclear interaction can be observed, should consist of 50% K^0 and 50% $\bar{K}^0$.

The K^0 and $\bar{K}^0$ must be absorbed differently; K^0 particles can only undergo elastic and charge exchange scattering, while $\bar{K}^0$ particles can also undergo strangeness exchange giving hyperons:

$$K^0 + p \to K^+ + n \tag{7.6}$$
$$K^0 + n \to K^0 + n \tag{7.7}$$

and

$$\bar{K}^0 + p \to \begin{cases} \Lambda^0 + \pi^+ \\ \Sigma^+ + \pi^0 \\ p + K^+ + K^- \end{cases} \tag{7.8}$$

$$\bar{K}^0 + n \to \Lambda^0 + \pi^0. \tag{7.9}$$

With more strong channels open, the $\bar{K}^0$ is therefore absorbed more strongly than K^0. After emerging from the slab, we shall therefore have a K^0 amplitude $f|K^0\rangle$ and a $\bar{K}^0$ amplitude $\bar{f}|\bar{K}^0\rangle$, where $\bar{f} < f < 1$. If we neglect the CP symmetry breaking, the emergent beam will be

$$\begin{aligned} \frac{1}{2}(f|K^0\rangle - \bar{f}|\bar{K}^0\rangle) &= \frac{f+\bar{f}}{2\sqrt{2}}(|K^0\rangle - |\bar{K}^0\rangle) + \frac{f-\bar{f}}{2\sqrt{2}}(|K^0\rangle + |\bar{K}^0\rangle) \\ &= \frac{1}{2}(f+\bar{f})|K_L\rangle + \frac{1}{2}(f-\bar{f})|K_S\rangle. \end{aligned} \tag{7.10}$$

Since $f \neq \bar{f}$, it follows that some of the K_S state has been regenerated.

The main prediction of the particle mixture hypothesis is the possibility of observing the $K_S - K_L$ interference. If we make the assumption of exponential decay, each component will have a time dependence of the form $e^{-(\Gamma_i/2\hbar + iE_i/\hbar)t}$ where E_i is the total energy of the particle $i = L, S$ and $\Gamma_i = \hbar/\tau_i$ is the width of the state, τ_i being the mean life in the frame in which the energy E_i is defined.

Set $\hbar = c = 1$, and measure all times in the rest frame, so that τ_i is the proper lifetime and $E_i = m_i$, the particle rest mass, then the time dependence became $e^{-(\Gamma_i/2+im_i)t}$.

Suppose that we produce K^0 at $t = 0$, from Eqs.(7.3), the K^0 state at $t = 0$ is a coherent superposition of K_S and K_L

$$|K^0\rangle = \sqrt{\frac{1+|\varepsilon|^2}{2(1+\varepsilon)^2}}(|K_S\rangle + |K_L\rangle). \tag{7.11}$$

After time t, as $|K_S\rangle$ and $|K_L\rangle$ states are definite mass eigenstates, we have

$$|K^0(t)\rangle = \sqrt{\frac{1+|\varepsilon|^2}{2(1+\varepsilon)^2}}\left[e^{-(\Gamma_S/2+im_S)t}|K_S\rangle + e^{-(\Gamma_L/2+im_L)t}|K_L\rangle\right] \tag{7.12}$$

and then after a time t for free decay in vacuo, the K^0 fraction of the beam will be

$$\begin{aligned} P(K^0,t) &= \left|\langle K^0(t)|K^0\rangle\right|^2 = \frac{1}{4}\left|e^{-(\Gamma_S/2+im_S)t} + e^{-(\Gamma_L/2+im_L)t}\right|^2 \\ &= \frac{1}{4}\left[e^{-\Gamma_S t} + e^{-\Gamma_L t} + e^{i(m_L-m_S)t}e^{-(\Gamma_S+\Gamma_L)t/2} + e^{-i(m_L-m_S)t}e^{-(\Gamma_S+\Gamma_L)t/2}\right] \\ &= \frac{1}{4}\left[e^{-\Gamma_S t} + e^{-\Gamma_L t} + 2e^{-(\Gamma_S+\Gamma_L)t/2}\cos(\Delta m t)\right], \end{aligned} \tag{7.13}$$

where $\Delta m = m_L - m_S = (3.489 \mp 0.009) \times 10^{-12} MeV$.

Similarly, the $\bar{K}^0$ fraction of the beam will be

$$P(\bar{K}^0,t) = \frac{1}{4}\left[e^{-\Gamma_S t} + e^{-\Gamma_L t} - 2e^{-(\Gamma_S+\Gamma_L)t/2}\cos(\Delta m t)\right]. \tag{7.14}$$

The neutral $K^0 - \bar{K}^0$ boson system is not the only one where the quantum mechanical mass mixing can be considered. We can expect to observe the same phenomenon in other neutral boson systems: $D^0 - \bar{D}^0$, $B^0 - \bar{B}^0$ and $\eta - \eta'$. Generally, flavor oscillations of particles can occur when states produced and detected in a given experiment, are superpositions of two or more eigenstates with different masses. The mesons oscillations has been used to place stringent constraints on physics beyond the Standard Model.

Chapter 8

Quantum field theory of boson mixing

We consider the quantum field theoretical formulation of boson field mixing and obtain the exact oscillation formula. This formula does not depend on arbitrary mass parameters. We show that the space for the mixed field states is unitarily inequivalent to the state space where the unmixed field operators are defined. We also study the structure of the currents and charges for the mixed fields.

8.1 Introduction

A rich non–perturbative vacuum structure has been discovered to be associated with the mixing of fermion fields in the context of Quantum Field Theory [25, 32]. The careful study of such a structure [28] has led to the determination of the exact QFT formula for neutrino oscillations [20, 21], exhibiting new features with respect to the usual quantum mechanical Pontecorvo formula [1]. Actually, it turns out [18, 26, 42, 43] that the non–trivial nature of the mixing transformations manifests itself also in the case of the mixing of boson fields. Of course, in this case the condensate structure for the "flavor" vacuum is very much different from the fermion case and a careful analysis is necessary in order to understand which phenomenological consequences are to be expected for the oscillations of mixed bosons.

In this chapter, we perform this analysis first at a formal level and then we study the oscillations of mixed mesons (charged and neutral systems). We will treat these particles as stable ones, an approximation which however does not affect the general validity of our results. In the framework of the QFT analysis of Refs. [18, 25], a study of the meson mixing and oscillations has been carried out in Ref.[42], where modifications to the usual oscillation formulas, connected with the vacuum structure, have been presented. However, the results of Ref. [42] can be improved in many respects [26] and in the present chapter we show that the oscillation formula there obtained has to be actually replaced with the exact one here presented.

In Section 8.2 we study the quantum field theory of two mixed spin-zero charged boson fields. In Section 8.3 we analyze the structure of currents for mixed fields and we derive the exact oscillation formula for complex and neutral fields in Sections 8.4 and 8.5, respectively. Section 8.6 is devoted to conclusions. Some mathematical derivations are given in the Appendix K.

8.2 Mixing of boson fields in QFT

The observed boson oscillations always involve particles with zero electrical charge. In the case of $K^0 - \bar{K}^0$, $B^0 - \bar{B}^0$, $D^0 - \bar{D}^0$, what oscillate are some other quantum numbers such as the strangeness and the isospin. Therefore, in the study of boson mixing, for these particles, we can consider [56] complex fields. The charge in question is some "flavor charge" (e.g. the strangeness) and thus the complex fields are "flavor charged" fields, referred to as "flavor fields" for simplicity.

We define the mixing relations as:

$$\begin{aligned}\phi_A(x) &= \phi_1(x)\ \cos\theta + \phi_2(x)\ \sin\theta \\ \phi_B(x) &= -\phi_1(x)\ \sin\theta + \phi_2(x)\ \cos\theta \end{aligned} \tag{8.1}$$

where generically we denote the mixed fields with suffixes A and B. Let the fields $\phi_i(x)$, $i = 1, 2$, be free complex fields with definite masses. Their conjugate momenta are $\pi_i(x) = \partial_0\phi_i^\dagger(x)$ and the commutation relations are the usual ones:

$$[\phi_i(x), \pi_j(y)]_{t=t'} = [\phi_i^\dagger(x), \pi_j^\dagger(y)]_{t=t'} = i\delta^3(\mathbf{x}-\mathbf{y})\,\delta_{ij}, \qquad i,j = 1,2\,. \tag{8.2}$$

with the other equal–time commutators vanishing. The Fourier expansions of fields and momenta are:

$$\phi_i(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\frac{1}{\sqrt{2\omega_{k,i}}}\left(a_{\mathbf{k},i}\,e^{-i\omega_{k,i}t} + b^\dagger_{-\mathbf{k},i}\,e^{i\omega_{k,i}t}\right)e^{i\mathbf{k}\cdot\mathbf{x}} \tag{8.3}$$

$$\pi_i(x) = i\int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\sqrt{\frac{\omega_{k,i}}{2}}\left(a^\dagger_{\mathbf{k},i}\,e^{i\omega_{k,i}t} - b_{-\mathbf{k},i}\,e^{-i\omega_{k,i}t}\right)e^{i\mathbf{k}\cdot\mathbf{x}}\,, \tag{8.4}$$

where $\omega_{k,i} = \sqrt{\mathbf{k}^2 + m_i^2}$ and $[a_{\mathbf{k},i}, a^\dagger_{\mathbf{p},j}] = [b_{\mathbf{k},i}, b^\dagger_{\mathbf{p},j}] = \delta^3(\mathbf{k}-\mathbf{p})\delta_{ij}$, with $i,j = 1,2$ and the other commutators vanishing. We will consider stable particles, which will not affect the general validity of our results.

We now proceed in a similar way to what has been done [25] for fermions and recast Eqs.(8.1) into the form [26]:

$$\phi_A(x) = G_\theta^{-1}(t)\; \phi_1(x)\; G_\theta(t) \tag{8.5}$$

$$\phi_B(x) = G_\theta^{-1}(t)\; \phi_2(x)\; G_\theta(t) \tag{8.6}$$

and similar ones for $\pi_A(x)$, $\pi_B(x)$. $G_\theta(t)$ denotes the operator which implements the mixing transformations (8.1):

$$G_\theta(t) = exp\left[-i\;\theta \int d^3\mathbf{x}\left(\pi_1(x)\phi_2(x) - \phi_1^\dagger(x)\pi_2^\dagger(x) - \pi_2(x)\phi_1(x) + \phi_2^\dagger(x)\pi_1^\dagger(x)\right)\right], \tag{8.7}$$

which is (at finite volume) a unitary operator: $G_\theta^{-1}(t) = G_{-\theta}(t) = G_\theta^\dagger(t)$. The generator of the mixing transformation in the exponent of $G_\theta(t)$ can also be written as

$$G_\theta(t) = \exp[\theta(S_+(t) - S_-(t))]\;. \tag{8.8}$$

The operators

$$S_+(t) = S_-^\dagger(t) \equiv -i \int d^3\mathbf{x}\;(\pi_1(x)\phi_2(x) - \phi_1^\dagger(x)\pi_2^\dagger(x))\,, \tag{8.9}$$

together with

$$S_3 \equiv \frac{-i}{2}\int d^3\mathbf{x}\left(\pi_1(x)\phi_1(x) - \phi_1^\dagger(x)\pi_1^\dagger(x) - \pi_2(x)\phi_2(x) + \phi_2^\dagger(x)\pi_2^\dagger(x)\right) \tag{8.10}$$

$$S_0 = \frac{Q}{2} \equiv \frac{-i}{2}\int d^3\mathbf{x}\left(\pi_1(x)\phi_1(x) - \phi_1^\dagger(x)\pi_1^\dagger(x) + \pi_2(x)\phi_2(x) - \phi_2^\dagger(x)\pi_2^\dagger(x)\right), \tag{8.11}$$

close the $su(2)$ algebra (at each time t):

$$[S_+(t), S_-(t)] = 2S_3, \quad [S_3, S_\pm(t)] = \pm S_\pm(t), \quad [S_0, S_3] = [S_0, S_\pm(t)] = 0. \tag{8.12}$$

Note that S_3 and S_0 are time independent. It is useful to write down explicitly the expansions of the above generators in terms of annihilation and creation operators:

$$S_+(t) = \int d^3\mathbf{k}\left(U_\mathbf{k}^*(t)\, a_{\mathbf{k},1}^\dagger a_{\mathbf{k},2} - V_\mathbf{k}^*(t)\, b_{-\mathbf{k},1} a_{\mathbf{k},2} + V_\mathbf{k}(t)\, a_{\mathbf{k},1}^\dagger b_{-\mathbf{k},2}^\dagger - U_\mathbf{k}(t)\, b_{-\mathbf{k},1} b_{-\mathbf{k},2}^\dagger\right), \tag{8.13}$$

$$S_-(t) = \int d^3\mathbf{k} \left(U_\mathbf{k}(t)\, a^\dagger_{\mathbf{k},2} a_{\mathbf{k},1} - V_\mathbf{k}(t)\, a^\dagger_{\mathbf{k},2} b^\dagger_{-\mathbf{k},1} + V^*_\mathbf{k}(t)\, b_{-\mathbf{k},2} a_{\mathbf{k},1} - U^*_\mathbf{k}(t)\, b_{-\mathbf{k},2} b^\dagger_{-\mathbf{k},1} \right), \tag{8.14}$$

$$S_3 = \frac{1}{2} \int d^3\mathbf{k} \left(a^\dagger_{\mathbf{k},1} a_{\mathbf{k},1} \,-\, b^\dagger_{-\mathbf{k},1} b_{-\mathbf{k},1} \,-\, a^\dagger_{\mathbf{k},2} a_{\mathbf{k},2} \,+\, b^\dagger_{-\mathbf{k},2} b_{-\mathbf{k},2} \right), \tag{8.15}$$

$$S_0 = \frac{1}{2} \int d^3\mathbf{k} \left(a^\dagger_{\mathbf{k},1} a_{\mathbf{k},1} \,-\, b^\dagger_{-\mathbf{k},1} b_{-\mathbf{k},1} \,+\, a^\dagger_{\mathbf{k},2} a_{\mathbf{k},2} \,-\, b^\dagger_{-\mathbf{k},2} b_{-\mathbf{k},2} \right). \tag{8.16}$$

As for the case of the fermion mixing, the structure of the generator Eq.(8.7) is recognized to be the one of a rotation combined with a Bogoliubov transformation (see below Eqs.(8.30)-(8.33)). Indeed, in the above equations, the coefficients

$$U_\mathbf{k}(t) \equiv |U_\mathbf{k}|\; e^{i(\omega_{k,2}-\omega_{k,1})t}\;, \qquad V_\mathbf{k}(t) \equiv |V_\mathbf{k}|\; e^{i(\omega_{k,1}+\omega_{k,2})t} \tag{8.17}$$

appear to be the Bogoliubov coefficients. They are defined as

$$|U_\mathbf{k}| \equiv \frac{1}{2}\left(\sqrt{\frac{\omega_{k,1}}{\omega_{k,2}}} + \sqrt{\frac{\omega_{k,2}}{\omega_{k,1}}} \right)\;, \qquad |V_\mathbf{k}| \equiv \frac{1}{2}\left(\sqrt{\frac{\omega_{k,1}}{\omega_{k,2}}} - \sqrt{\frac{\omega_{k,2}}{\omega_{k,1}}} \right) \tag{8.18}$$

and satisfy the relation

$$|U_\mathbf{k}|^2 - |V_\mathbf{k}|^2 = 1\,, \tag{8.19}$$

which is in fact to be expected in the boson case (note the difference with respect to the fermion case [25]). We can thus put

$$|U_\mathbf{k}| \equiv \cosh \xi^\mathbf{k}_{1,2}\;, \qquad |V_\mathbf{k}| \equiv \sinh \xi^\mathbf{k}_{1,2}, \quad \text{with} \quad \xi^\mathbf{k}_{1,2} = \frac{1}{2} \ln\left(\frac{\omega_{k,1}}{\omega_{k,2}} \right). \tag{8.20}$$

We now consider the action of the generator of the mixing transformations on the vacuum $|0\rangle_{1,2}$ for the fields $\phi_{1,2}(x)$: $a_{\mathbf{k},i}|0\rangle_{1,2} = 0,\ i = 1,2$. The generator induces an $SU(2)$ coherent state structure on such state [44]:

$$|0(\theta,t)\rangle_{A,B} \equiv G^{-1}_\theta(t)\; |0\rangle_{1,2}\,. \tag{8.21}$$

From now on we will refer to the state $|0(\theta,t)\rangle_{A,B}$ as to the "flavor" vacuum for bosons. The suffixes A and B label the flavor charge content of the state. We have

$${}_{A,B}\langle 0(\theta,t)|0(\theta,t)\rangle_{A,B} \;=\; 1. \tag{8.22}$$

In the following, we will consider the Hilbert space for flavor fields at a given time t, say $t = 0$, and it is useful to define

$$|0(t)\rangle_{A,B} \equiv |0(\theta,t)\rangle_{A,B} \quad \text{and} \quad |0\rangle_{A,B} \equiv |0(\theta,t=0)\rangle_{A,B} \tag{8.23}$$

for future reference. A crucial point is that the flavor and the mass vacua are orthogonal in the infinite volume limit [26]. We indeed have (see Appendix K):

$$ {}_{1,2}\langle 0|0(t)\rangle_{A,B} = \prod_{\mathbf{k}} {}_{1,2}\langle 0|G^{-1}_{\mathbf{k},\theta}(t)|0\rangle_{1,2} = \prod_{\mathbf{k}} f^{\mathbf{k}}_0(\theta) \ , \quad \text{for any } t, \tag{8.24} $$

where we have used $G^{-1}_\theta(t) = \prod_{\mathbf{k}} G^{-1}_{\mathbf{k},\theta}(t)$ (see Eqs.(8.8), (8.13) and (8.14)). In the infinite volume limit, we obtain

$$ \lim_{V\to\infty} {}_{1,2}\langle 0|0(t)\rangle_{A,B} = \lim_{V\to\infty} e^{\frac{V}{(2\pi)^3}\int d^3\mathbf{k} \, \ln f^{\mathbf{k}}_0(\theta)} = 0 \ , \quad \text{for any } t. \tag{8.25} $$

From the Appendix K, Eq.(K.8), we see that $\ln f^{\mathbf{k}}_0(\theta)$ is indeed negative for any values of $\mathbf{k}$, θ and m_1, m_2 (note that $0 \leq \theta \leq \pi/4$). We also observe that the orthogonality disappears when $\theta = 0$ and/or $m_1 = m_2$, consistently with the fact that in both cases there is no mixing. These features are similar to the case of fermion mixing [25]: the orthogonality is essentially due to the infinite number of degrees of freedom [22, 23].

We can define annihilation operators for the vacuum $|0(t)\rangle_{A,B}$ as

$$ a_{\mathbf{k},A}(\theta,t) \equiv G^{-1}_\theta(t)\; a_{\mathbf{k},1}\; G_\theta(t), \tag{8.26} $$

$$ a_{\mathbf{k},B}(\theta,t) \equiv G^{-1}_\theta(t)\; a_{\mathbf{k},2}\; G_\theta(t), \tag{8.27} $$

$$ b_{-\mathbf{k},A}(\theta,t) \equiv G^{-1}_\theta(t)\; b_{-\mathbf{k},1}\; G_\theta(t), \tag{8.28} $$

$$ b_{-\mathbf{k},B}(\theta,t) \equiv G^{-1}_\theta(t)\; b_{-\mathbf{k},2}\; G_\theta(t), \tag{8.29} $$

with $a_{\mathbf{k},A}(\theta,t)|0(t)\rangle_{A,B} = 0$. For simplicity we will use the notation $a_{\mathbf{k},A}(t) \equiv a_{\mathbf{k},A}(\theta,t)$. Explicitly, we have:

$$ a_{\mathbf{k},A}(t) = \cos\theta\; a_{\mathbf{k},1} + \sin\theta \left(U^*_{\mathbf{k}}(t)\; a_{\mathbf{k},2} + V_{\mathbf{k}}(t)\; b^\dagger_{-\mathbf{k},2} \right) , \tag{8.30} $$

$$ a_{\mathbf{k},B}(t) = \cos\theta\; a_{\mathbf{k},2} - \sin\theta \left(U_{\mathbf{k}}(t)\; a_{\mathbf{k},1} - V_{\mathbf{k}}(t)\; b^\dagger_{-\mathbf{k},1} \right) , \tag{8.31} $$

$$ b_{-\mathbf{k},A}(t) = \cos\theta\; b_{-\mathbf{k},1} + \sin\theta \left(U^*_{\mathbf{k}}(t)\; b_{-\mathbf{k},2} + V_{\mathbf{k}}(t)\; a^\dagger_{\mathbf{k},2} \right) , \tag{8.32} $$

$$ b_{-\mathbf{k},B}(t) = \cos\theta\; b_{-\mathbf{k},2} - \sin\theta \left(U_{\mathbf{k}}(t)\; b_{-\mathbf{k},1} - V_{\mathbf{k}}(t)\; a^\dagger_{\mathbf{k},1} \right) . \tag{8.33} $$

These operators satisfy the canonical commutation relations (at equal times). In their expressions the Bogoliubov transformation part is evidently characterized by the terms with the U and V coefficients. The condensation density of the flavor vacuum is given for any t by

$$ {}_{A,B}\langle 0(t)|a^\dagger_{\mathbf{k},i} a_{\mathbf{k},i}|0(t)\rangle_{A,B} = {}_{A,B}\langle 0(t)|b^\dagger_{-\mathbf{k},i} b_{-\mathbf{k},i}|0(t)\rangle_{A,B} = \sin^2\theta\; |V_{\mathbf{k}}|^2, \tag{8.34} $$

with $i = 1, 2$.

8.2.1 Arbitrary mass parameterization

Above we have expanded the mixed fields $\phi_{A,B}$ in the same basis as the free fields $\phi_{1,2}$. However, as noticed in the case of fermion mixing [28], this is not the most general possibility. Indeed, one could as well expand the flavor fields in a basis of fields with arbitrary masses. Of course, these arbitrary mass parameters should not appear in the physically observable quantities. Thus, as a check for the validity of the oscillation formula we are going to derive in Section 8.4, it is important to consider this generalization. Let us first rewrite the free fields $\phi_{1,2}$ in the form [26]

$$\begin{aligned}
\phi_i(x) &= \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} \left(u^{\phi}_{\mathbf{k},i}(t)\, a_{\mathbf{k},i} + v^{\phi}_{-\mathbf{k},i}(t) b^{\dagger}_{-\mathbf{k},i} \right) e^{i\mathbf{k}\cdot\mathbf{x}} , && (8.35)\\
\pi_i(x) &= i \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} \left(u^{\pi}_{\mathbf{k},i}(t)\, a^{\dagger}_{\mathbf{k},i} - v^{\pi}_{-\mathbf{k},i}(t)\, b_{-\mathbf{k},i} \right) e^{i\mathbf{k}\cdot\mathbf{x}} \quad , \qquad i = 1,2, && (8.36)
\end{aligned}$$

where we have introduced the notation

$$\begin{aligned}
u^{\phi}_{\mathbf{k},i}(t) &\equiv \frac{1}{\sqrt{2\omega_{k,i}}} e^{-i\omega_{k,i}t} \quad , \quad v^{\phi}_{-\mathbf{k},i}(t) \equiv \frac{1}{\sqrt{2\omega_{k,i}}} e^{i\omega_{k,i}t} , && (8.37)\\
u^{\pi}_{\mathbf{k},i}(t) &\equiv \sqrt{\frac{\omega_{k,i}}{2}} e^{i\omega_{k,i}t} \quad , \quad v^{\pi}_{-\mathbf{k},i}(t) \equiv \sqrt{\frac{\omega_{k,i}}{2}} e^{-i\omega_{k,i}t}, \qquad i = 1,2 \, . && (8.38)
\end{aligned}$$

We now define

$$\begin{aligned}
\rho^{\mathbf{k}*}_{\alpha\beta}(t) &\equiv u^{\pi}_{\mathbf{k},\alpha}(t) u^{\phi}_{\mathbf{k},\beta}(t) + v^{\phi}_{-\mathbf{k},\alpha}(t) v^{\pi}_{-\mathbf{k},\beta}(t) = e^{i(\omega_{k,\alpha}-\omega_{k,\beta})t} \cosh \xi^{\mathbf{k}}_{\alpha,\beta} \, , && (8.39)\\
\lambda^{\mathbf{k}*}_{\alpha\beta}(t) &\equiv v^{\pi}_{-\mathbf{k},\alpha}(t) u^{\phi}_{\mathbf{k},\beta}(t) - u^{\phi}_{\mathbf{k},\alpha}(t) v^{\pi}_{-\mathbf{k},\beta}(t) = e^{-i(\omega_{k,\alpha}+\omega_{k,\beta})t} \sinh \xi^{\mathbf{k}}_{\alpha,\beta} \, , && (8.40)\\
\xi^{\mathbf{k}}_{\alpha,\beta} &\equiv \frac{1}{2} \ln \left(\frac{\omega_{k,\alpha}}{\omega_{k,\beta}} \right) \quad , \qquad \alpha, \beta = 1, 2, A, B, && (8.41)
\end{aligned}$$

where $\omega_{k,\alpha} \equiv \sqrt{k^2 + \mu^2_{\alpha}}$. We denote with μ_A and μ_B the arbitrary mass parameters while $\mu_1 \equiv m_1$ and $\mu_2 \equiv m_2$ are the physical masses. Note that $\rho^{\mathbf{k}}_{12}(t) = U_{\mathbf{k}}(t)$ and $\lambda^{\mathbf{k}}_{12}(t) = V_{\mathbf{k}}(t)$. We can now write the expansion of the flavor fields in the general form (we use a tilde to denote the generalized ladder operators):

$$\phi_{\sigma}(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} \left(u^{\phi}_{\mathbf{k},\sigma}(t)\, \widetilde{a}_{\mathbf{k},\sigma}(t) + v^{\phi}_{-\mathbf{k},\sigma}(t)\, \widetilde{b}^{\dagger}_{-\mathbf{k},\sigma}(t) \right) e^{i\mathbf{k}\cdot\mathbf{x}}, \qquad (8.42)$$

with $\sigma = A, B$, which is to be compared with the expansion in the free field basis as given in Eqs.(8.5), (8.6):

$$\phi_{\sigma}(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} \left(u^{\phi}_{\mathbf{k},i}(t)\, a_{\mathbf{k},\sigma}(t) + v^{\phi}_{-\mathbf{k},i}(t)\, b^{\dagger}_{-\mathbf{k},\sigma}(t) \right) e^{i\mathbf{k}\cdot\mathbf{x}} \, , \qquad (8.43)$$

where $(\sigma, i) = (A,1), (B,2)$.

The relation between the two sets of flavor operators is given as [26]

$$\left(\begin{array}{c} \widetilde{a}_{\mathbf{k},\sigma}(t) \\ \widetilde{b}^{\dagger}_{-\mathbf{k},\sigma}(t) \end{array}\right) = J^{-1}(t) \left(\begin{array}{c} a_{\mathbf{k},\sigma}(t) \\ b^{\dagger}_{-\mathbf{k},\sigma}(t) \end{array}\right) J(t) = \left(\begin{array}{cc} \rho^{\mathbf{k}*}_{\sigma i}(t) & \lambda^{\mathbf{k}}_{\sigma i}(t) \\ \lambda^{\mathbf{k}*}_{\sigma i}(t) & \rho^{\mathbf{k}}_{\sigma i}(t) \end{array}\right) \left(\begin{array}{c} a_{\mathbf{k},\sigma}(t) \\ b^{\dagger}_{-\mathbf{k},\sigma}(t) \end{array}\right), \tag{8.44}$$

$$J(t) = \exp\left\{ \int d^3\mathbf{k}\, \xi^{\mathbf{k}}_{\sigma,i} \left[a^{\dagger}_{\mathbf{k},\sigma}(t) b^{\dagger}_{-\mathbf{k},\sigma}(t) - b_{-\mathbf{k},\sigma}(t) a_{\mathbf{k},\sigma}(t) \right] \right\}, \tag{8.45}$$

with

$$\xi^{\mathbf{k}}_{\sigma,i} \equiv \frac{1}{2} \ln\left(\frac{\omega_{k,\sigma}}{\omega_{k,i}}\right). \tag{8.46}$$

For $\mu_A = m_1$ and $\mu_B = m_2$ one has $J = 1$. Note that the transformation Eq.(8.44) is in fact a Bogoliubov transformation which leaves invariant the form $a^{\dagger}_{\mathbf{k},\sigma}(t) a_{\mathbf{k},\sigma}(t) - b^{\dagger}_{-\mathbf{k},\sigma}(t) b_{-\mathbf{k},\sigma}(t)$.

8.3 The currents for mixed boson fields

Before presenting the exact oscillation formula, let us investigate in this Section the structure of currents and charges for the mixed fields [26]. This will enable us to identify the relevant physical observables to look at for flavor oscillations. Since we are here interested in vacuum oscillations, in the following we neglect interaction terms and only consider the free field Lagrangian for two charged scalar fields with a mixed mass term:

$$\mathcal{L}(x) = \partial_\mu \Phi^{\dagger}_f(x)\, \partial^\mu \Phi_f(x) - \Phi^{\dagger}_f(x) M \Phi_f(x), \tag{8.47}$$

with $\Phi^T_f = (\phi_A, \phi_B)$,

$$M = \left(\begin{array}{cc} m^2_A & m^2_{AB} \\ m^2_{AB} & m^2_B \end{array}\right). \tag{8.48}$$

By means of Eq.(8.1),

$$\Phi_f(x) = \left(\begin{array}{cc} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{array}\right) \Phi_m(x), \tag{8.49}$$

$\mathcal{L}$ becomes diagonal in the basis $\Phi^T_m = (\phi_1, \phi_2)$:

$$\mathcal{L}(x) = \partial_\mu \Phi^{\dagger}_m(x)\, \partial^\mu \Phi_m(x) - \Phi^{\dagger}_m(x) M_d \Phi_m(x), \tag{8.50}$$

where $M_d = diag(m_1^2, m_2^2)$ and

$$m_A^2 = m_1^2 \cos^2\theta + m_2^2 \sin^2\theta \tag{8.51}$$

$$m_B^2 = m_1^2 \sin^2\theta + m_2^2 \cos^2\theta \tag{8.52}$$

$$m_{AB}^2 = (m_2^2 - m_1^2)\sin\theta\cos\theta. \tag{8.53}$$

The Lagrangian $\mathcal{L}$ is invariant under the global $U(1)$ phase transformations

$$\Phi'_m = e^{i\alpha}\,\Phi_m \tag{8.54}$$

as a result, we have the conservation of the Noether charge

$$Q = \int d^3\mathbf{x}\, I^0(x), \tag{8.55}$$

which is indeed the total charge of the system (we have $I^\mu(x) = i\Phi_m^\dagger(x)\overset{\leftrightarrow}{\partial^\mu}\Phi_m(x)$ with $\overset{\leftrightarrow}{\partial^\mu} \equiv \overset{\rightarrow}{\partial^\mu} - \overset{\leftarrow}{\partial^\mu}$).

Let us now consider the $SU(2)$ transformation

$$\Phi'_m(x) = e^{i\alpha_j\tau_j}\,\Phi_m(x) \quad , \quad j = 1,2,3\,, \tag{8.56}$$

with α_j real constants, $\tau_j = \sigma_j/2$ and σ_j being the Pauli matrices. For $m_1 \neq m_2$, the Lagrangian is not generally invariant under (8.56) and we obtain, by use of the equations of motion,

$$\delta\mathcal{L}(x) = -i\,\alpha_j\,\Phi_m^\dagger(x)\,[M_d\,,\,\tau_j]\,\Phi_m(x) = -\alpha_j\,\partial_\mu\,J^\mu_{m,j}(x)\,, \tag{8.57}$$

where the currents are

$$J_m^\mu(x) = \frac{\partial\mathcal{L}}{\partial\left(\partial_\mu\Phi_m(x)\right)}\delta\Phi_m(x) + \delta\Phi_m^\dagger(x)\frac{\partial\mathcal{L}}{\partial\left(\partial_\mu\Phi_m^\dagger(x)\right)}, \tag{8.58}$$

and then

$$J^\mu_{m,j}(x) = i\,\Phi_m^\dagger(x)\,\tau_j\,\overset{\leftrightarrow}{\partial^\mu}\,\Phi_m(x) \quad , \quad j = 1,2,3. \tag{8.59}$$

We thus obtain the currents:

$$J^\mu_{m,1} = \frac{i}{2}\left[(\partial^\mu\phi_1^\dagger)\phi_2 + (\partial^\mu\phi_2^\dagger)\phi_1 - \phi_1^\dagger(\partial^\mu\phi_2) - \phi_2^\dagger(\partial^\mu\phi_1)\right], \tag{8.60}$$

$$J^\mu_{m,2} = \frac{1}{2}\left[(\partial^\mu\phi_1^\dagger)\phi_2 - (\partial^\mu\phi_2^\dagger)\phi_1 - \phi_1^\dagger(\partial^\mu\phi_2) + \phi_2^\dagger(\partial^\mu\phi_1)\right], \tag{8.61}$$

$$J^\mu_{m,3} = \frac{i}{2}\left[(\partial^\mu \phi_1^\dagger)\phi_1 - (\partial^\mu \phi_2^\dagger)\phi_2 - \phi_1^\dagger(\partial^\mu \phi_1) + \phi_2^\dagger(\partial^\mu \phi_2)\right]. \tag{8.62}$$

The corresponding charges,

$$Q_{m,j}(t) \equiv \int d^3\mathbf{x}\, J^0_{m,j}(x), \tag{8.63}$$

close the $su(2)$ algebra (at each time t). The Casimir operator C_m is proportional to the total charge:

$$C_m \equiv \Big[\sum_{j=1}^{3} Q^2_{m,j}(t)\Big]^{\frac{1}{2}} = \frac{1}{2}Q. \tag{8.64}$$

Observe also that the transformation induced by $Q_{m,2}(t)$,

$$\Phi_f(x) \;=\; e^{-2i\theta Q_{m,2}(t)}\Phi_m(x)e^{2i\theta Q_{m,2}(t)} \tag{8.65}$$

is just the mixing transformation Eq.(8.49). Thus $2Q_{m,2}(t)$ is the generator of the mixing transformations. Moreover,

$$Q_{m,\pm}(t) \equiv \frac{1}{2}\left[Q_{m,1}(t) \pm iQ_{m,2}(t)\right], \tag{8.66}$$

$Q_{m,3}$, and C_m are nothing but $S_\pm(t)$, S_3, and S_0, respectively, as introduced in Eqs.(8.9)-(8.11). From Eq.(8.57) we also see that $Q_{m,3}$ and C_m are conserved, consistently with Eqs.(8.15), (8.16). Observe that the combinations

$$Q_{1,2} \;\equiv\; \frac{1}{2}Q \pm Q_{m,3} \tag{8.67}$$

$$Q_i \;=\; \int d^3\mathbf{k}\left(a^\dagger_{\mathbf{k},i}a_{\mathbf{k},i} \,-\, b^\dagger_{-\mathbf{k},i}b_{-\mathbf{k},i}\right) \quad , \qquad i = 1,2, \tag{8.68}$$

are simply the conserved[1] (Noether) charges for the free fields ϕ_1 and ϕ_2 with

$$Q_1 + Q_2 = Q. \tag{8.69}$$

We now perform the $SU(2)$ transformations on the flavor doublet Φ_f:

$$\Phi'_f(x) \,=\, e^{i\alpha_j\tau_j}\,\Phi_f(x) \quad , \qquad j \,=\, 1,2,3, \tag{8.70}$$

and obtain:

$$\delta\mathcal{L}(x) \;=\; -i\,\alpha_j\,\Phi_f^\dagger(x)\,[M,\tau_j]\,\Phi_f(x) \;=\; -\alpha_j\,\partial_\mu J^\mu_{f,j}(x)\;, \tag{8.71}$$

$$J^\mu_{f,j}(x) \;=\; i\,\Phi_f^\dagger(x)\,\tau_j\,\overset{\leftrightarrow}{\partial^\mu}\,\Phi_f(x) \quad , \qquad j \,=\, 1,2,3. \tag{8.72}$$

---

[1] Note that, in absence of mixing, these charges would indeed be the flavor charges, being the flavor conserved for each generation.

The related charges,

$$Q_{f,j}(t) \equiv \int d^3\mathbf{x}\, J^0_{f,j}(x) \tag{8.73}$$

still fulfil the $su(2)$ algebra and

$$C_f = C_m = \frac{1}{2}Q. \tag{8.74}$$

Due to the off–diagonal (mixing) terms in the mass matrix M, $Q_{f,3}(t)$ is time–dependent. This implies an exchange of charge between ϕ_A and ϕ_B, resulting in the flavor oscillations. This suggests to us to define indeed the *flavor charges* as

$$Q_A(t) \;\equiv\; \frac{1}{2}Q \,+\, Q_{f,3}(t)\;, \tag{8.75}$$

$$Q_B(t) \;\equiv\; \frac{1}{2}Q \,-\, Q_{f,3}(t)\;, \tag{8.76}$$

with

$$Q_A(t) \,+\, Q_B(t) \,=\, Q. \tag{8.77}$$

These charges have a simple expression in terms of the flavor ladder operators:

$$Q_\sigma(t) \;=\; \int d^3\mathbf{k}\, \Big(a^\dagger_{\mathbf{k},\sigma}(t) a_{\mathbf{k},\sigma}(t) \,-\, b^\dagger_{-\mathbf{k},\sigma}(t) b_{-\mathbf{k},\sigma}(t)\Big) \quad , \qquad \sigma = A, B\;. \tag{8.78}$$

This is because they are connected to the Noether charges Q_i of Eq.(8.68) via the mixing generator:

$$Q_\sigma(t) = G^{-1}_\theta(t)\; Q_i\; G_\theta(t), \tag{8.79}$$

with $(\sigma, i) = (A, 1), (B, 2)$. Note that the flavor charges are invariant under the transformation Eq.(8.44).

8.4 The oscillation formula for mixed bosons

Let us now calculate the oscillation formula for mixed bosons [26]. We will first follow the approach of Binger and Ji [42] and show that the oscillation formulas there presented exhibit a dependence on the arbitrary mass parameters μ_σ, a feature which is not physically acceptable. We will do this by using the generalized operators introduced above. Then we will show how to cure this pathology [26], in analogy to what was done in Chapter 2 for the case of fermion mixing [20, 21], where the exact formula for neutrino oscillations was derived and it was shown to be independent from the arbitrary mass parameters that can be introduced in the expansions of the flavor fields.

8.4.1 The oscillation formula of Binger and Ji

Following Ref.[42], let us define the (generalized) flavor state by acting on the *mass vacuum* $|0\rangle_{1,2}$ with the flavor creation operators (we omit momentum indices):

$$|\widetilde{a}_A\rangle \equiv \widetilde{a}_A^\dagger|0\rangle_{1,2} \;=\; \rho_{A1}\cos\theta|a_1\rangle \;+\; \rho_{A2}\sin\theta|a_2\rangle \tag{8.80}$$

with $|a_i\rangle = a_i^\dagger|0\rangle_{12}$.

As already discussed in Ref.[42], the flavor state so defined is not normalized and the normalization factor has to be introduced as

$$\widetilde{\mathcal{N}}_A \equiv \langle\widetilde{a}_A|\widetilde{a}_A\rangle \;=\; \rho_{A1}^2\cos^2\theta + \rho_{A2}^2\sin^2\theta\ . \tag{8.81}$$

We have

$$\langle\widetilde{a}_A|\widetilde{N}_A|\widetilde{a}_A\rangle \;=\; (\rho_{A1}^2+\lambda_{A1}^2)\cos^2\theta + (\rho_{A2}^2+\lambda_{A2}^2)\sin^2\theta\ . \tag{8.82}$$

The oscillation formula then follow as:

$$\langle\widetilde{a}_A(t)|\widetilde{N}_A|\widetilde{a}_A(t)\rangle \;=\; \langle\widetilde{a}_A|\widetilde{N}_A|\widetilde{a}_A\rangle - 4\frac{\rho_{A1}^2\rho_{A2}^2}{\widetilde{\mathcal{N}}_A}\sin^2\theta\cos^2\theta\sin^2\left(\frac{\Delta E}{2}t\right) \tag{8.83}$$

and a similar one for the expectation value of $\widetilde{N}_B$. From the above, as announced, it is evident that these formulas explicitly depend on the (arbitrary) parameters μ_σ (see Eq.(8.39)). We also note that, for $\mu_A = m_1$ and $\mu_B = m_2$, one has $\rho_{A1} = 1$, $\lambda_{A1} = 0$, $\rho_{A2} = U$ and $\lambda_{A2} = V$. Consequently, Eqs.(8.82), (8.83) reduce respectively to Eqs.(18) and (20) of Ref.[42].

8.4.2 The exact oscillation formula

We now show how a consistent treatment of the flavor oscillation for bosons in QFT can be given which does not exhibit the above pathological dependence on arbitrary parameters [26].

There are two key points to be remarked. A general feature of field mixing is that the number operator for mixed particles is not a well-defined operator. It is so because the mixing transformations mix creation and annihilation operators and then the annihilation (creation) operators for flavor particles and antiparticles do not commute at different times (see Eqs.(8.30)-(8.33)). Moreover, the number operator does depend on the arbitrary mass parameters. Much care is therefore required in the use of the number operator. A second remark is that the flavor states are not to be defined by using the vacuum $|0\rangle_{1,2}$: the flavor states so defined are in fact not normalized and the normalization factor Eq.(8.81) depends on the arbitrary mass parameters.

These two difficulties can be bypassed by using the remedy already adopted in chapter 2 [20, 21] for the case of fermions: the flavor states have shown to be consistently defined by acting with the flavor creation operators on the *flavor vacuum*. The observable quantities are then the expectation values of the *flavor charges* on the flavor states: the oscillation formulas thus obtained do not depend on the arbitrary mass parameters.

Let us now define[2] the state of the a_A particle as

$$|\widetilde{a}_{\mathbf{k},A}\rangle_{A,B} \equiv \widetilde{a}^{\dagger}_{\mathbf{k},A}(0)|\widetilde{0}\rangle_{A,B} \tag{8.84}$$

with

$$|\widetilde{0}\rangle_{A,B} = J^{-1}|0\rangle_{A,B}, \tag{8.85}$$

and consider the expectation values of the flavor charges Eq.(8.78) on it (analogous results follow if one considers $|\widetilde{a}_{\mathbf{k},B}\rangle_{A,B}$). We obtain:

$$\widetilde{\mathcal{Q}}^{A}_{\mathbf{k},\sigma}(t) \equiv {}_{A,B}\langle\widetilde{a}_{\mathbf{k},A}|\widetilde{Q}_{\sigma}(t)|\widetilde{a}_{\mathbf{k},A}\rangle_{A,B} = \left|\left[\widetilde{a}_{\mathbf{k},\sigma}(t), \widetilde{a}^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 - \left|\left[\widetilde{b}^{\dagger}_{-\mathbf{k},\sigma}(t), \widetilde{a}^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2, \tag{8.86}$$

with $\sigma = A, B$.

We also have ${}_{A,B}\langle\widetilde{0}|\widetilde{Q}_{\mathbf{k},\sigma}(t)|\widetilde{0}\rangle_{A,B} = 0$ and $\widetilde{\mathcal{Q}}^{A}_{\mathbf{k},A}(t) + \widetilde{\mathcal{Q}}^{A}_{\mathbf{k},B}(t) = 1$.

A straightforward direct calculation shows that the above quantities *do not depend* on μ_A and μ_B, i.e.:

$${}_{A,B}\langle\widetilde{a}_{\mathbf{k},A}|\widetilde{Q}_{\mathbf{k},\sigma}(t)|\widetilde{a}_{\mathbf{k},A}\rangle_{A,B} = {}_{A,B}\langle a_{\mathbf{k},A}|Q_{\mathbf{k},\sigma}(t)|a_{\mathbf{k},A}\rangle_{A,B} \quad , \qquad \sigma = A, B\,, \tag{8.87}$$

and similar one for the expectation values on $|\tilde{a}_{\mathbf{k},B}\rangle_{A,B}$. Eq.(8.87) is a central result of this chapter [26]: it confirms that the only physically relevant quantities are the above expectation values of flavor charges. Note that expectation values of the number operator, of the kind

$${}_{A,B}\langle\widetilde{a}_{\mathbf{k},A}|\widetilde{N}_{\sigma}(t)|\widetilde{a}_{\mathbf{k},A}\rangle_{A,B} = \left|\left[\widetilde{a}_{\mathbf{k},\sigma}(t), \widetilde{a}^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 \tag{8.88}$$

and similar ones, do indeed depend on the arbitrary mass parameters, although the flavor states are properly defined (i.e. on the flavor Hilbert space). The cancellation of these parameters happens *only* when considering the combination of squared modula of commutators of the form Eq.(8.86)[3]. A similar cancellation occurs for fermions [21] with the sum of the squared modula of anticommutators.

---

[2] In the following, we will work in the Heisenberg picture: this is particularly convenient in the present context since special care has to be taken with the time dependence of flavor states (see the discussion in Ref.[20]).

Finally, the explicit calculation [26] gives

$$\begin{aligned} \mathcal{Q}^{A}_{\mathbf{k},A}(t) &= \left|\left[a_{\mathbf{k},A}(t), a^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 - \left|\left[b^{\dagger}_{-\mathbf{k},A}(t), a^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 \\ &= 1 - \sin^2(2\theta)\left[|U_{\mathbf{k}}|^2 \, \sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) - |V_{\mathbf{k}}|^2 \, \sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right] , \end{aligned} \tag{8.89}$$

$$\begin{aligned} \mathcal{Q}^{A}_{\mathbf{k},B}(t) &= \left|\left[a_{\mathbf{k},B}(t), a^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 - \left|\left[b^{\dagger}_{-\mathbf{k},B}(t), a^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 \\ &= \sin^2(2\theta)\left[|U_{\mathbf{k}}|^2 \, \sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) - |V_{\mathbf{k}}|^2 \, \sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right] \end{aligned} \tag{8.90}$$

Notice the negative sign in front of the $|V_{\mathbf{k}}|^2$ terms in these formulas, in contrast with the fermion case [20, 21]: the boson flavor charge can assume also negative values. This fact points to the statistical nature of the phenomenon: it means that when dealing with mixed fields, one intrinsically deals with a many–particle system, i.e. a genuine field theory phenomenon. This situation has a strong analogy with Thermal Field Theory (i.e. QFT at finite temperature) [23], where quasi–particle states are ill defined and only statistical averages make sense. Of course, there is no violation of charge conservation for the overall system of two mixed fields.

The above formulas are obviously different from the usual quantum mechanical oscillation formulas, which however are recovered in the relativistic limit (i.e. for $|\mathbf{k}|^2 \gg \frac{m_1^2+m_2^2}{2}$). Apart from the extra oscillating term (the one proportional to $|V_{\mathbf{k}}|^2$) and the momentum dependent amplitudes, the QFT formulas carry the remarkable information about the statistics of the oscillating particles: for bosons and fermions the amplitudes (Bogoliubov coefficients) are drastically different according to the two different statistics ($|U_{\mathbf{k}}|$ and $|V_{\mathbf{k}}|$ are circular functions in the fermion case and hyperbolic functions in the boson case). This fact also fits with the above mentioned statistical nature of the oscillation phenomenon in QFT. Note also that our treatment is essentially non–perturbative [26].

In order to better appreciate the features of the QFT formulas, it is useful to plot the oscillating charge in time for sample values of the masses and for different values of the momentum (we use same units for masses and momentum). It is evident how the effect of the extra oscillating term is maximal at lower momenta (see Figures 8.1

---

[3] One may think it could make sense to take the expectation value of the flavor charges on states defined on the mass Hilbert space, as the ones defined in Eq.(8.80). A direct calculation however shows that this is not the case and these expectation values depends on the mass parameters: the conclusion is that one must use the flavor Hilbert space.



Figure 8.1: Plot of $\mathcal{Q}^A_{\mathbf{k},B}(t)$ in function of time for $k = 0$, $m_1 = 2$, $m_2 = 50$ and $\theta = \pi/4$.

and 8.2) and disappears for large k (see Figure 8.3) where the standard oscillation pattern is recovered. In the following plots we use $T_k = \frac{4\pi}{\omega_{k,2} - \omega_{k,1}}$ and assume maximal mixing.

It is also interesting to plot the time average of the oscillating charge,

$$\bar{\mathcal{Q}}^A_{\mathbf{k},B} = \frac{1}{nT_k} \int_0^{nT_k} dt \, \mathcal{Q}^A_{\mathbf{k},B}(t), \tag{8.91}$$

as a function of the momentum. In Figure 8.4 we plot $\mathcal{Q}^A_{\mathbf{k},B}(t)$ averaged over two different time intervals, i.e. for $n = 10$ and $n = 100$: it is interesting to observe how the larger is the time interval, the more the curve converges to the average of the standard formula, which has the value $\frac{1}{2}$. The behavior for large k is due to the fact that, as already observed, the exact oscillation formula reduces to the quantum mechanical oscillation one in the large momentum limit (i.e. for $|\mathbf{k}|^2 \gg \frac{m_1^2 + m_2^2}{2}$).



Figure 8.2: Plot of $\mathcal{Q}^A_{\mathbf{k},B}(t)$ in function of time for $k = 10$, $m_1 = 2$, $m_2 = 50$ and $\theta = \pi/4$.



Figure 8.3: Plot of $\mathcal{Q}^A_{\mathbf{k},B}(t)$ in function of time for $k = 100$, $m_1 = 2$, $m_2 = 50$ and $\theta = \pi/4$.



Figure 8.4: The time average of $\mathcal{Q}^A_{\mathbf{k},B}(t)$ over 10 T_k (dashed line) and over 100 T_k (solid line) with respect to the average of the standard oscillation formula (horizontal axis) as a function of k for the values $m_1 = 2$, $m_2 = 50$ and $\theta = \pi/4$.

8.5 Mixing of neutral particles

Analyzing the mixing phenomena for the meson sector, we should also keep in mind another important point. For example, the $\eta - \eta'$ mixing (which is one of the most interesting systems due to the large mass difference of the mixed components) involves the two *neutral* particles so that the mixing formulas for the charged particles do not immediately apply. With this in mind, we consider here the case of mixing of two spin zero neutral boson fields [41, 43] . We follow the case of charged fields as discussed above [18, 26, 42] and introduce the following lagrangian:

$$\begin{aligned} \mathcal{L}(x) &= \partial_\mu \Phi_f^T(x) \partial^\mu \Phi_f(x) \, - \, \Phi_f^T(x) \; M \; \Phi_f(x) \qquad (8.92) \\ &= \partial_\mu \Phi_m^T(x) \partial^\mu \Phi_m(x) \, - \, \Phi_m^T(x) \; M_d \; \Phi_m(x) \qquad (8.93) \end{aligned}$$

with $\Phi_f^T = (\phi_A, \phi_B)$ being the flavor fields and $M = \begin{pmatrix} m_A^2 & m_{AB}^2 \\ m_{AB}^2 & m_B^2 \end{pmatrix}$. Those are connected to the free fields $\Phi_m^T = (\phi_1, \phi_2)$ with $M_d = diag(m_1^2, m_2^2)$ by a rotation:

$$\phi_A(x) = \phi_1(x) \; \cos\theta + \phi_2(x) \; \sin\theta \qquad (8.94)$$

$$\phi_B(x) = -\phi_1(x) \; \sin\theta + \phi_2(x) \; \cos\theta \qquad (8.95)$$

and a similar one for the conjugate momenta $\pi_i = \partial_0 \phi_i$. The free fields ϕ_i can be quantized in the usual way (we use $x_0 \equiv t$):

$$\phi_i(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} \frac{1}{\sqrt{2\omega_{k,i}}} \left(a_{\mathbf{k},i}\, e^{-i\omega_{k,i}t} + a^\dagger_{-\mathbf{k},i}\, e^{i\omega_{k,i}t} \right) e^{i\mathbf{k}\mathbf{x}} \tag{8.96}$$

$$\pi_i(x) = -i \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}} \sqrt{\frac{\omega_{k,i}}{2}} \left(a_{\mathbf{k},i}\, e^{-i\omega_{k,i}t} - a^\dagger_{-\mathbf{k},i}\, e^{i\omega_{k,i}t} \right) e^{i\mathbf{k}\mathbf{x}}, \tag{8.97}$$

with $i = 1, 2$ and $\omega_{k,i} = \sqrt{\mathbf{k}^2 + m_i^2}$.

The commutation relations are:

$$\left[\phi_i(x), \pi_j(y)\right]_{x_0=y_0} = i\, \delta_{ij}\, \delta^3(\mathbf{x}-\mathbf{y}) \ , \qquad \left[a_{\mathbf{k},i}, a^\dagger_{\mathbf{p},j}\right] = \delta_{ij}\, \delta^3(\mathbf{k}-\mathbf{p}). \tag{8.98}$$

We now recast Eqs.(8.94),(8.95) into the form:

$$\phi_A(x) = G_\theta^{-1}(t)\ \phi_1(x)\ G_\theta(t) \tag{8.99}$$

$$\phi_B(x) = G_\theta^{-1}(t)\ \phi_2(x)\ G_\theta(t) \tag{8.100}$$

and similar ones for $\pi_A(x)$, $\pi_B(x)$, where $G_\theta(t)$ is the generator of the mixing transformations (8.94),(8.95):

$$G_\theta(t) = \exp\left[-i\, \theta \int d^3\mathbf{x}\, (\pi_1(x)\phi_2(x) - \pi_2(x)\phi_1(x)) \right] , \tag{8.101}$$

which is (at finite volume) a unitary operator: $G_\theta^{-1}(t) = G_{-\theta}(t) = G_\theta^\dagger(t)$.

The mixing generator is given by

$$G_\theta(t) = \exp[\theta(S_+(t) - S_-(t))] \tag{8.102}$$

and the $su(2)$ operators are now realized as

$$S_+(t) \equiv -i \int d^3\mathbf{x}\ \pi_1(x)\phi_2(x) \tag{8.103}$$

$$S_-(t) \equiv -i \int d^3\mathbf{x}\ \pi_2(x)\phi_1(x) \tag{8.104}$$

$$S_3 \equiv \frac{-i}{2} \int d^3\mathbf{x}\, (\pi_1(x)\phi_1(x) - \pi_2(x)\phi_2(x)) \tag{8.105}$$

$$S_0 \equiv \frac{-i}{2} \int d^3\mathbf{x}\, (\pi_1(x)\phi_1(x) + \pi_2(x)\phi_2(x)) \ . \tag{8.106}$$

We have, explicitly

$$S_+(t) - S_-(t) = \int d^3\mathbf{k}\Big(U^*_{\mathbf{k}}(t)\; a^\dagger_{\mathbf{k},1}a_{\mathbf{k},2} - V^*_{\mathbf{k}}(t)\; a_{-\mathbf{k},1}a_{\mathbf{k},2} + V_{\mathbf{k}}(t)\; a^\dagger_{\mathbf{k},2}a^\dagger_{-\mathbf{k},1} - U_{\mathbf{k}}(t)\; a^\dagger_{\mathbf{k},2}a_{\mathbf{k},1}\Big) \tag{8.107}$$

where $U_{\mathbf{k}}(t)$ and $V_{\mathbf{k}}(t)$ are Bogoliubov coefficients given by

$$U_{\mathbf{k}}(t) \equiv |U_{\mathbf{k}}|\; e^{i(\omega_{k,2}-\omega_{k,1})t} \quad , \quad V_{\mathbf{k}}(t) \equiv |V_{\mathbf{k}}|\; e^{i(\omega_{k,1}+\omega_{k,2})t} \tag{8.108}$$

$$|U_{\mathbf{k}}| \equiv \frac{1}{2}\left(\sqrt{\frac{\omega_{k,1}}{\omega_{k,2}}} + \sqrt{\frac{\omega_{k,2}}{\omega_{k,1}}}\right) \quad , \quad |V_{\mathbf{k}}| \equiv \frac{1}{2}\left(\sqrt{\frac{\omega_{k,1}}{\omega_{k,2}}} - \sqrt{\frac{\omega_{k,2}}{\omega_{k,1}}}\right) \tag{8.109}$$

$$|U_{\mathbf{k}}|^2 - |V_{\mathbf{k}}|^2 = 1\,, \tag{8.110}$$

The flavor fields can be expanded as:

$$\phi_\sigma(x) = \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\frac{1}{\sqrt{2\omega_{k,j}}}\left(a_{\mathbf{k},\sigma}(t)\; e^{-i\omega_{k,j}t} + a^\dagger_{-\mathbf{k},\sigma}(t)\; e^{i\omega_{k,j}t}\right) e^{i\mathbf{k}\cdot\mathbf{x}} \tag{8.111}$$

$$\pi_\sigma(x) = -i \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{3}{2}}}\sqrt{\frac{\omega_{k,j}}{2}}\left(a_{\mathbf{k},\sigma}(t)\; e^{-i\omega_{k,j}t} - a^\dagger_{-\mathbf{k},\sigma}(t)\; e^{i\omega_{k,j}t}\right) e^{i\mathbf{k}\cdot\mathbf{x}}\,, \tag{8.112}$$

with $\sigma, j = (A,1), (B,2)$ and the flavor annihilation operators given by:

$$a_{\mathbf{k},A}(t) \equiv G^{-1}_\theta(t)\; a_{\mathbf{k},1}\; G_\theta(t) = \cos\theta\; a_{\mathbf{k},1} + \sin\theta\; \Big(U^*_{\mathbf{k}}(t)\; a_{\mathbf{k},2} + V_{\mathbf{k}}(t)\; a^\dagger_{-\mathbf{k},2}\Big)\,, \tag{8.113}$$

$$a_{\mathbf{k},B}(t) \equiv G^{-1}_\theta(t)\; a_{\mathbf{k},2}\; G_\theta(t) = \cos\theta\; a_{\mathbf{k},2} - \sin\theta\; \Big(U_{\mathbf{k}}(t)\; a_{\mathbf{k},1} - V_{\mathbf{k}}(t)\; a^\dagger_{-\mathbf{k},1}\Big)\,. \tag{8.114}$$

We now consider the action of the generator of the mixing transformations on the vacuum $|0\rangle_{1,2}$ for the fields $\phi_i(x)$: $a_{\mathbf{k},i}|0\rangle_{1,2} = 0,\; i = 1,2$. The generator induces an $SU(2)$ coherent state structure on such state [44]:

$$|0(\theta,t)\rangle_{A,B} \equiv G^{-1}_\theta(t)\; |0\rangle_{1,2}\,. \tag{8.115}$$

The state $|0(\theta,t)\rangle_{A,B}$ is to the *flavor vacuum* for neutral bosons.

As in the case of complex boson fields, we define the state for a mixed particle with "flavor" A and momentum $\mathbf{k}$ as:

$$|a_{\mathbf{k},A}(t)\rangle \equiv a^\dagger_{\mathbf{k},A}(t)|0(t)\rangle_{A,B} = G^{-1}_\theta(t)a^\dagger_{\mathbf{k},1}|0\rangle_{1,2} \tag{8.116}$$

In the following we work in the Heisenberg picture, flavor states will be taken at reference time $t = 0$ (including the flavor vacuum). We also define $|a_{\mathbf{k},A}\rangle \equiv |a_{\mathbf{k},A}(0)\rangle$.

Let us now consider the (non-vanishing) commutators of the flavor ladder operators at different times:

$$\left[a_{\mathbf{k},A}(t), a^{\dagger}_{\mathbf{k},A}(t')\right] = \cos^2\theta + \sin^2\theta \left(|U_{\mathbf{k}}|^2 e^{-i(\omega_2-\omega_1)(t-t')} - |V_{\mathbf{k}}|^2 e^{i(\omega_2+\omega_1)(t-t')}\right), \tag{8.117}$$

$$\left[a^{\dagger}_{-\mathbf{k},A}(t), a^{\dagger}_{\mathbf{k},A}(t')\right] = \sin^2\theta\, |U_{\mathbf{k}}||V_{\mathbf{k}}| \left(e^{-i\omega_2(t-t')} - e^{i\omega_2(t-t')}\right) e^{-i\omega_1(t+t')}, \tag{8.118}$$

$$\left[a_{\mathbf{k},B}(t), a^{\dagger}_{\mathbf{k},A}(t')\right] = \cos\theta\sin\theta\, |U_{\mathbf{k}}| \left(e^{i(\omega_2-\omega_1)t'} - e^{i(\omega_2-\omega_1)t}\right), \tag{8.119}$$

$$\left[a^{\dagger}_{-\mathbf{k},B}(t), a^{\dagger}_{\mathbf{k},A}(t')\right] = \cos\theta\sin\theta\, |V_{\mathbf{k}}| \left(e^{-i(\omega_2+\omega_1)t} - e^{-i(\omega_2+\omega_1)t'}\right). \tag{8.120}$$

We observe that the following quantity is constant in time:

$$\begin{aligned} &\left|\left[a_{\mathbf{k},A}(t), a^{\dagger}_{\mathbf{k},A}(t')\right]\right|^2 - \left|\left[a^{\dagger}_{-\mathbf{k},A}(t), a^{\dagger}_{\mathbf{k},A}(t')\right]\right|^2 + \\ + &\left|\left[a_{\mathbf{k},B}(t), a^{\dagger}_{\mathbf{k},A}(t')\right]\right|^2 - \left|\left[a^{\dagger}_{-\mathbf{k},B}(t), a^{\dagger}_{\mathbf{k},A}(t')\right]\right|^2 = 1\,. \end{aligned} \tag{8.121}$$

In the case of the complex boson fields [26], the corresponding of Eq.(8.121) was consistently interpreted as expressing the conservation of total charge[4]. In the present case we are dealing with a neutral field and thus the charge operator vanishes identically. Nevertheless the quantities in Eq.(8.121) are well defined and are the neutral-field counterparts of the corresponding ones for the case of charged fields. Thus we look for a physical interpretation of such oscillating quantities.

Let us consider the momentum operator, defined as the diagonal space part of the energy-momentum tensor [22]:

$$P^j \equiv \int d^3\mathbf{x}\,\Theta^{0j}(x), \tag{8.122}$$

with

$$\Theta^{\mu\nu} \equiv \partial^\mu\phi\partial^\nu\phi - g^{\mu\nu}\left[\frac{1}{2}(\partial\phi)^2 - \frac{1}{2}m^2\phi^2\right]. \tag{8.123}$$

For the free fields ϕ_i we have:

$$\mathbf{P}_i = \int d^3\mathbf{x}\,\pi_i(x)\nabla\phi_i(x) = \int d^3\mathbf{k}\,\frac{\mathbf{k}}{2}\left(a^{\dagger}_{\mathbf{k},i}a_{\mathbf{k},i} - a^{\dagger}_{-\mathbf{k},i}a_{-\mathbf{k},i}\right), \tag{8.124}$$

---

[4] We have, for charged bosonic fields:

$$\langle a_{\mathbf{k},A}|Q_\sigma(t)|a_{\mathbf{k},A}\rangle = \left|\left[a_{\mathbf{k},\sigma}(t), a^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 - \left|\left[b^{\dagger}_{-\mathbf{k},\sigma}(t), a^{\dagger}_{\mathbf{k},A}(0)\right]\right|^2 \quad, \qquad \sigma = A, B\,.$$

together with ${}_{A,B}\langle 0|Q_\sigma(t)|0\rangle_{A,B} = 0$ and $\sum_\sigma \langle a_{\mathbf{k},A}|Q_\sigma(t)|a_{\mathbf{k},A}\rangle = 1$.

with $i = 1, 2$.

In a similar way we can define the momentum operator for mixed fields:

$$\mathbf{P}_\sigma(t) \;=\; \int d^3\mathbf{x}\, \pi_\sigma(x) \nabla \phi_\sigma(x) \;=\; \int d^3\mathbf{k}\, \frac{\mathbf{k}}{2} \left(a^\dagger_{\mathbf{k},\sigma}(t) a_{\mathbf{k},\sigma}(t) \,-\, a^\dagger_{-\mathbf{k},\sigma}(t) a_{-\mathbf{k},\sigma}(t) \right), \tag{8.125}$$

with $\sigma = A, B$. The two operators are obviously related: $\mathbf{P}_\sigma(t) = G^{-1}_\theta(t)\, \mathbf{P}_i\, G_\theta(t)$. Note that the total momentum is conserved in time since commutes with the generator of mixing transformations (at any time):

$$\mathbf{P}_A(t) \,+\, \mathbf{P}_B(t) \;=\; \mathbf{P}_1 \,+\, \mathbf{P}_2 \;\equiv\; \mathbf{P} \tag{8.126}$$

$$[\mathbf{P}\,,\, G_\theta(t)] \;=\; 0 \quad , \quad [\mathbf{P}\,,\, H] \;=\; 0\,. \tag{8.127}$$

Thus in the mixing of neutral fields, the momentum operator plays an analogous rôle to that of the charge for charged fields [41]. For charged fields, the total charge operator, associated with the $U(1)$ invariance of the Lagrangian, is proportional to the Casimir of the $SU(2)$ group associated to the generators of the mixing transformations. This is not true anymore for the case of neutral fields, although the $SU(2)$ structure persists in this case as well (see Eqs.(8.103),(8.105)).

We now consider the expectation values of the momentum operator for flavor fields on the flavor state $|a_{\mathbf{k},A}\rangle$ with definite momentum $\mathbf{k}$. Obviously, this is an eigenstate of $\mathbf{P}_A(t)$ at time $t = 0$:

$$\mathbf{P}_A(0)\, |a_{\mathbf{k},A}\rangle \;=\; \mathbf{k}\, |a_{\mathbf{k},A}\rangle\,, \tag{8.128}$$

which follows from $\mathbf{P}_1\, |a_{\mathbf{k},1}\rangle \;=\; \mathbf{k}\, |a_{\mathbf{k},1}\rangle$ by application of $G^{-1}_\theta(0)$. At time $t \neq 0$, the expectation value of the momentum (normalized to the initial value) gives:

$$\mathcal{P}^A_\sigma(t) \;\equiv\; \frac{\langle a_{\mathbf{k},A}|\mathbf{P}_\sigma(t)|a_{\mathbf{k},A}\rangle}{\langle a_{\mathbf{k},A}|\mathbf{P}_\sigma(0)|a_{\mathbf{k},A}\rangle} \;=\; \left|\left[a_{\mathbf{k},\sigma}(t), a^\dagger_{\mathbf{k},A}(0)\right]\right|^2 \;-\; \left|\left[a^\dagger_{-\mathbf{k},\sigma}(t), a^\dagger_{\mathbf{k},A}(0)\right]\right|^2, \tag{8.129}$$

with $\sigma = A, B$, which is of the same form as the expression one obtains for the charged field.

One can explicitly check that the (flavor) vacuum expectation value of the momentum operator $\mathbf{P}_\sigma(t)$ does vanish at all times:

$${}_{A,B}\langle 0|\mathbf{P}_\sigma(t)|0\rangle_{A,B} \;=\; 0 \quad , \qquad \sigma = A, B \tag{8.130}$$

which can be understood intuitively by realizing that the flavor vacuum $|0\rangle_{A,B}$ does not carry momentum since it is a condensate of pairs carrying zero total momentum (like the BCS ground state, for example).



Figure 8.5: Plot of $\mathcal{P}^{\eta}_{\mathbf{k},\eta}(t)$ (solid line) and $\mathcal{P}^{\eta}_{\mathbf{k},\eta'}(t)$ (dashed line) in function of time for an initially pure η state and for $k = 0$, $m_1 = 549$ MeV, $m_2 = 958$ MeV and $\theta = -54^o$.

The explicit calculation of the oscillating quantities $\mathcal{P}^{A}_{\mathbf{k},\sigma}(t)$ gives:

$$\mathcal{P}^{A}_{\mathbf{k},A}(t) \;=\; 1 - \sin^2(2\theta)\left[|U_{\mathbf{k}}|^2\ \sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) - |V_{\mathbf{k}}|^2\ \sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right] \tag{8.131}$$

$$\mathcal{P}^{A}_{\mathbf{k},B}(t) \;=\; \sin^2(2\theta)\left[|U_{\mathbf{k}}|^2\ \sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) - |V_{\mathbf{k}}|^2\ \sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right]. \tag{8.132}$$

in complete agreement with the charged field case [26].

The Eqs.(8.131), (8.132) are the flavor oscillation formulas for the neutral mesons, such as $\eta - \eta'$, $\phi - \omega$. By definition of the momentum operator, the Eqs.(8.131), (8.132) are the relative population densities of flavor particles in the beam.

We present in Figure 8.5 a plot of momentum oscillations for the system of η and η' ($m_\eta = 547MeV$, $\Gamma_\eta = 1.18keV$, $m_{\eta'} = 958MeV$, $\Gamma_{\eta'} = 0.2MeV$ and $\theta \approx -54^\circ$ [27]) with zero momenta (i.e. $k = 0GeV$).

8.6 Conclusions

In this chapter we have considered the quantum field theoretical formulation of spin-zero (charged and neutral) boson field mixing and obtained the exact oscillation formula which does not depend on arbitrary mass parameters which can be introduced in full generality in the theory. We have also studied the structure of the currents and charges for the charged mixed fields. In order to make our discussion more transparent, we have neglected the instability of the oscillating particles. This does not affect the general validity of our result which rests on the intrinsic features of QFT.

A crucial point in our analysis is the disclosure of the fact that the space for the mixed field states is unitarily inequivalent to the state space where the unmixed field operators are defined. This is a common feature with the QFT structure of the fermion mixing, which has recently been established [20, 21, 25, 28, 32]). The vacuum for the mixed fields turns out to be a generalized $SU(2)$ coherent state. Of course, in the boson case the condensate structure for the "flavor" vacuum is found to be very much different from the one in the fermion case. Besides the intrinsic mathematical interest, our analysis provides interesting phenomenological insights. It leads to the exact oscillation formula for bosons which predicts oscillation behaviors susceptible of being experimentally tested.

In the framework of the QFT analysis of Refs. [18, 25], a study of the meson mixing and oscillations has been already carried out in Ref.[42]. However, the results of Ref.[42] give observable quantities which are dependent on arbitrary mass parameters, and this is of course physically not acceptable [28]. In the present chapter we have pointed out the origin of such a pathology and have shown how to obtain results which are independent from arbitrary parameters [26]. The oscillation formula obtained in Ref.[42] has to be actually replaced with the exact one here presented [26]. Moreover we have studied the case of the neutral bosons [41, 43].

Let us close by observing that although our QFT analysis discloses features which cannot be ignored in any further study of the field mixing and oscillations, since they are intrinsic to the structure of the QFT formalism, nevertheless there are many aspects of the physics of mixing which are not fully understood and many features are still obscure [57], as already observed in the introduction. The mixing of neutrinos and their oscillations seem to be now experimentally established and quark mixing and meson mixing are widely accepted and verified. However, several questions [58, 59, 60, 61, 62, 63, 64, 65] are the object of active discussion in the framework of the quantum mechanics formalism for neutrino oscillations. As a matter of fact, such a state of affairs has been a strong motivation for our searching in the structural aspects of QFT a possible hint to the understanding of particle mixing and oscillations.

Chapter 9

Phenomenology of flavor oscillations with nonperturbative Quantum Field Theoretic effects

We consider the quantum field theoretical formulation of meson mixing and obtain the exact oscillation formula in the presence of the decay. This formula is different from quantum mechanical formula by additional high-frequency oscillation terms. We analyze phenomenological aspects of this nonperturbative effect and find that the systems where this phenomena could be detected are $\eta-\eta'$ and $\phi-\omega$ mesons.

9.1 Introduction

In the chapters 2, 3, 8 it was found that the fermions and bosons oscillation formula in quantum field theoretical formulation is modified by additional high-frequency terms and simpler quantum mechanical result is reproduced only in the relativistic limit.

In this chapter, we attempt the phenomenological analysis of this nonperturbative phenomenon. The mixing of particles and antiparticles in the meson sector (e.g., $K^0-\bar{K}^0$, $B^0-\bar{B}^0$) requires specific adjustments to the results obtained in the case of charged fields. Moreover, except neutrinos, all known mixed systems are subject to decay and thus the effect of particle life-time should also be taken into account.

Specifically, in Section 9.2, we analyze the adjustments needed for the general formulation in order to make applications for the known systems. We also study the effect of the finite particle life-time on the field-theoretical oscillation formula. Finally we estimate the magnitudes of the nonperturbative corrections in various systems and discuss the systems in which the field-theoretical effect may be most significant [43]. The Section 9.3 is devoted to conclusions.

9.2 Phenomenological application of the nonperturbative oscillation formula

9.2.1 $K^0 - \bar{K}^0$, $B^0 - \bar{B}^0$ and $D^0 - \bar{D}^0$ mixing

The typical systems for flavor mixing are $K^0-\bar{K}^0$, $B^0-\bar{B}^0$ and $D^0-\bar{D}^0$. The $K^0-\bar{K}^0$ and $B^0-\bar{B}^0$ systems provide the evidence of CP-violation in weak interaction and the $D^0-\bar{D}^0$ is also important for the analysis of CKM mixing matrix. As an illustration of the mixing in these systems, we consider in particular the $K^0-\bar{K}^0$ system here.

In $K^0-\bar{K}^0$ mixing, K^0 may not be treated as neutral since $K^0 \neq \bar{K}^0$. Of course, this is not the mixing of two different charged particles either. Rather, the particle here is mixed with its antiparticle. In this case it is important to identify the mixed degrees of freedom properly. Note that in $K^0-\bar{K}^0$ mixing there are three distinct modes, namely the strange eigenstates $K^0-\bar{K}^0$, the mass eigenstates $K_L - K_S$ and the CP eigenstates $K_1 - K_2$. Each pair can be written as a linear combination of the other ones, e.g.

$$\begin{aligned} K_1 &= \tfrac{1}{\sqrt{2}}(K^0+\bar{K}^0), \quad & K_2 &= \tfrac{1}{\sqrt{2}}(K^0-\bar{K}^0); \\ K^0 &= \tfrac{e^{i\delta}}{\sqrt{2}}(K_L+K_S), \quad & \bar{K}^0 &= \tfrac{e^{-i\delta}}{\sqrt{2}}(K_L-K_S); \\ K_1 &= \tfrac{1}{\sqrt{1+|\epsilon|^2}}(K_S+\epsilon K_L), \quad & K_2 &= \tfrac{1}{\sqrt{1+|\epsilon|^2}}(K_L+\epsilon K_S); \end{aligned} \tag{9.1}$$

with $e^{i\delta}$ being a complex phase and $\epsilon = i\delta$ being the imaginary CP-violation parameter. $K^0-\bar{K}^0$ are produced as strange eigenstates and propagate as the mass eigenstates K_L, K_S.

The mass eigenstates K_L and K_S are defined as the +1 and -1 CPT eigenstates, respectively, so that they can be represented in terms of self-adjoint scalar fields ϕ_1, ϕ_2 as

$$K_L = \phi_1, \quad K_S = i\phi_2. \tag{9.2}$$

Therefore the mixing in this system is similar to the case of neutral fields with *complex* mixing matrix. Since the complex mixing matrix in SU(2) can be always transformed into the real one by suitable redefinition of the field phases which does not affect the expectation values, the mixing in this case is equivalent to the mixing of neutral fields.

The oscillating observables may be that of the strange charge (in the system K^0 and $\bar{K}^0$ taken as flavor A and B, respectively) with the trivial mixing angle $\theta = \pi/4$ from Eq.(9.1).

Phenomenologically relevant, however, is the oscillation of CP-eigenvalue which determines the ratio of experimentally measured $\pi\pi$ to $\pi\pi\pi$ decay rates. CP-oscillations are given in terms of K_1 and K_2 flavors with small mixing angle

$$\cos(\theta) = 1/\sqrt{1+|\epsilon|^2}. \tag{9.3}$$

9.2.2 The effect of the particle decay

To analyze the phenomenological aspects of the quantum field theory oscillation formulas for fermions and bosons, we should remember that, excluding the mixing of neutrinos, most mixed particles are unstable. Hence, we need to consider the effect of the finite lifetime in the oscillation formulas of Eqs.(8.131) and (8.132).

The particle decay is taken in account by inserting, by hand, as usually done, the factor $e^{-\Gamma t}$ in the annihilation (creation) operators [43]:

$$a_{\mathbf{k},i} \longrightarrow a_{\mathbf{k},i} e^{-\frac{\Gamma_i}{2} t}, \tag{9.4}$$

$$b_{-\mathbf{k},i} \longrightarrow b_{-\mathbf{k},i} e^{-\frac{\Gamma_i}{2} t}. \tag{9.5}$$

Then, the oscillation formulas can be written as

$$\begin{aligned} \mathcal{P}^A_{\mathbf{k},A}(t) &= \left|\left[a_{\mathbf{k},A}(t), a^\dagger_{\mathbf{k},A}(0)\right]\right|^2 - \left|\left[b^\dagger_{-\mathbf{k},A}(t), a^\dagger_{\mathbf{k},A}(0)\right]\right|^2 \\ &= \left(\cos^2\theta e^{-\frac{\Gamma_1}{2}t} + \sin^2\theta e^{-\frac{\Gamma_2}{2}t}\right)^2 \\ &- \sin^2(2\theta) e^{-\frac{\Gamma_1+\Gamma_2}{2}t}\left[|U_\mathbf{k}|^2 \, \sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) - |V_\mathbf{k}|^2 \, \sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right], \end{aligned} \tag{9.6}$$

$$\begin{aligned} \mathcal{P}^A_{\mathbf{k},B}(t) &= \left|\left[a_{\mathbf{k},B}(t), a^\dagger_{\mathbf{k},A}(0)\right]\right|^2 - \left|\left[b^\dagger_{-\mathbf{k},B}(t), a^\dagger_{\mathbf{k},A}(0)\right]\right|^2 \\ &= \sin^2(2\theta)\Bigg(\left[\frac{e^{-\frac{\Gamma_1}{2}t} - e^{-\frac{\Gamma_2}{2}t}}{2}\right]^2 \\ &+ e^{-\frac{\Gamma_1+\Gamma_2}{2}t}\left[|U_\mathbf{k}|^2 \, \sin^2\left(\frac{\omega_{k,2}-\omega_{k,1}}{2}t\right) - |V_\mathbf{k}|^2 \, \sin^2\left(\frac{\omega_{k,2}+\omega_{k,1}}{2}t\right)\right]\Bigg). \end{aligned} \tag{9.7}$$

We note the difference between these oscillation formulas and the quantum mechanical Gell-Mann–Pais formulas. Essentially the quantum field theoretic corrections are proportional to $|V_\mathbf{k}|^2$ and appear as the additional high-frequency oscillation terms.

9.2.3 The nonperturbative effect in the boson sector

We are now in the position to analyze the known mixing systems and compute the magnitude of the nonperturbative correction in each system [43]. In the boson sector the typical mixed systems are $K^0-\bar{K}^0$, $B^0-\bar{B}^0$, $D^0-\bar{D}^0$, $\eta-\eta'$ and $\phi-\omega$. One may expect that the nonperturbative high-frequency effects may be observed adequately in some of these systems since not only the mass difference for mesons may be as large as $400MeV$ (e.g. $\eta - \eta'$) but also in principle the mesons may be produced at low momenta where the flavor vacuum effect is most prominent. In practice, however, we note that one may encounter certain experimental difficulty due to either extremely short lifetime of mixed mesons or extremely small mass difference in mass-eigenstates for the systems except $\eta - \eta'$ and $\phi - \omega$.

The magnitude of the nonperturbative term

In all of field-theoretical derivations (See Eq.(9.6)-(9.7)), the field-theoretical effect (or the high-frequency oscillation term) is proportional to $|V_{\mathbf{k}}|^2$. In estimating the maximal magnitude of this term, it is useful to write $|V_{\mathbf{k}}|^2$ in terms of the dimensionless momentum

$$p \equiv \sqrt{\frac{2|\mathbf{k}|^2}{m_1^2 + m_2^2}} \tag{9.8}$$

and the dimensionless parameter

$$a \equiv \frac{m_2^2 - m_1^2}{m_1^2 + m_2^2} \tag{9.9}$$

as follows:

$$|V(p,a)|^2 \;=\; \frac{p^2+1}{2\sqrt{(p^2+1)^2 - a^2}} - \frac{1}{2}, \tag{9.10}$$

from which we see that $|V_{\mathbf{k}}|^2$ is maximal at $p = 0$ ($|V_{max}|^2 = \frac{(m_1 - m_2)^2}{4m_1 m_2}$) and goes to zero for large momenta (i.e. for $|\mathbf{k}|^2 \gg \frac{m_1^2 + m_2^2}{2}$).

A plot of the condensation density $|V(p,a)|^2$ as a function of p is presented in Figure 9.1 for two sample values of the parameter a.

The optimal observation scale for field theoretical effect in meson mixing, therefore, is $k = 0$ and the maximum correction is of the order of $|V|^2 \sim \frac{\Delta m^2}{m^2}$. For known parameters of meson mixing we find

- $K^0 - \bar{K}^0$: $|V|^2 \sim (\frac{3.510^{-12} MeV}{500 MeV})^2 \sim 10^{-28}$,
- $D^0 - \bar{D}^0$: $|V|^2 \sim (\frac{10^{-12} MeV}{1000 MeV})^2 \sim 10^{-30}$,
- $B^0 - \bar{B}^0$: $|V|^2 \sim (\frac{510^{-10} MeV}{5000 MeV})^2 \sim 10^{-26}$,
- $B_s^0 - \bar{B}_s^0$: $|V|^2 \sim (\frac{210^{-8} MeV}{10^4 MeV})^2 \sim 10^{-28}$,
- $\eta - \eta'$: $|V|^2 \sim (\frac{400 MeV}{700 MeV})^2 \sim 0.2$,



Figure 9.1: The bosonic condensation density $|V(p,a)|^2$ as a function of p for $a = 0.98$ (for instance: $m_1 = 5$, $m_2 = 50$ in dimensionless units, solid line) and $a = 0.8$ (for instance: $m_1 = 5$, $m_2 = 15$ in dimensionless units, dashed line).

- $\omega - \phi$: $|V|^2 \sim (\frac{200MeV}{900MeV})^2 \sim 0.05$.

From the above discussion we observe hence that the nontrivial flavor vacuum effect can be maximally seen in the mixed systems such as $\eta - \eta'$ and $\omega - \phi$, thus one needs to be careful about taking them into account should these systems ever be used in some sort of mixing experiments.

9.2.4 The nonperturbative effect in fermion sector

We can employ the similar method in the fermion sector. Since neutrinos are stable, no additional adjustments are necessary to the known results [25]. We can write the field-theoretical correction amplitude $|V_{\mathbf{k}}|^2$ in the fermion case as a function of the dimensionless momentum

$$p = \frac{|\mathbf{k}|}{\sqrt{m_1 m_2}} \tag{9.11}$$

and of the dimensionless parameter

$$a = \frac{(\Delta m)^2}{m_1 m_2} \quad , \quad 0 \le a < +\infty \ , \tag{9.12}$$

as follows:

$$|V_{\mathbf{k}}|^2 \equiv |V(p,a)|^2 \ = \ \frac{1}{2}\left(1 - \frac{p^2+1}{\sqrt{(p^2+1)^2 + ap^2}}\right) \ . \tag{9.13}$$



Figure 9.2: The fermion condensation density $|V_{\mathbf{k}}|^2$ in function of $|\mathbf{k}|$ and for sample values of the parameters m_1 and m_2. Solid line: $m_1 = 1$, $m_2 = 100$; Long-dashed line: $m_1 = 10$, $m_2 = 100$; Short-dashed line: $m_1 = 10$, $m_2 = 1000$.

where $\Delta m \equiv m_2 - m_1$ (we take $m_1 \leq m_2$).

From Eq.(9.13), we see that the effect is maximal when $p = 1$ ($|V(1, a)|^2 = |V_{max}|^2 \approx \frac{(m_1 - m_2)^2}{16 m_1 m_2}$). $|V_{\mathbf{k}}|^2$ goes asymptotically to 1/2 when $a \to \infty$ and goes to zero for large momenta (i.e. for $|\mathbf{k}|^2 \gg \frac{m_1^2 + m_2^2}{2}$) as $|V|^2 \approx \frac{\Delta m^2}{4k^2}$.

In Figure 9.2 we show the fermion condensation density $|V_{\mathbf{k}}|^2$ in function of $|\mathbf{k}|$ and for sample values of the parameters m_1 and m_2.

For the currently known neutrino mixing parameters, i.e. $\Delta m \sim 10^{-3} eV$, the best observation scale is of the order of $k \approx 10^{-3} eV$. However, experimentally observed neutrinos are always in extremelly relativistic domain and therefore the value of $|V_{\mathbf{k}}|^2$ is as small as $|V_{\mathbf{k}}|^2 \sim 10^{-18}$. Only for extremely low energies (like those in neutrino cosmological background) the field-theoretical corrections might be large and account for few percent. In this connection, we observe that the non-perturbative field theory effects, in spite of the small corrections they induce in the oscillation amplitudes, nevertheless they may contribute in a specific and crucial way in other physical contexts or phenomena. Indeed, as shown in chapter 6, the mixing of neutrinos may specifically contribute to the value of the cosmological constant exactly because of the non-perturbative effects expressed by the non-zero value of $|V_{\mathbf{k}}|^2$ [34].

9.3 Conclusions

In this chapter we have considered phenomenological aspects of the quantum field theoretical formalism of flavor mixing. A crucial point in this analysis is the disclosure of the fact that the space for the mixed field states is unitarily inequivalent to the state space where the unmixed field operators are defined. This is a common feature with the QFT structure of mixing, which has recently been established [20, 21, 25, 33]. The vacuum for the mixed fields turns out to be a generalized SU(2) coherent state.

We have estimated the magnitude of the field-theoretical effect in known mixed systems. We found that for most mixed systems both in meson and neutrino sectors this effect is negligible. Only in strongly mixed systems such as $\omega - \phi$ or $\eta - \eta'$, or for very low-energy neutrino effects the corrections may be as large as 5-20% [43] and thus additional attention may be needed to see if these systems can be used in oscillation experiments. The non-perturbative vacuum effect is the most prominent when the particles are produced at low momentum.

Moreover, we recall that the neutrino mixing may contribute to the value of the cosmological constant because of the non-perturbative effects.

Appendix A

Double beta decay

Double beta decay, the rarest known nuclear decay process, can occur in different modes:

$$\begin{aligned} &2\nu\beta\beta - \text{decay}: && A(Z,N) \rightarrow A(Z+2,N-2) + 2e^- + 2\bar{\nu}_e && \text{(A.1)} \\ &0\nu\beta\beta - \text{decay}: && A(Z,N) \rightarrow A(Z+2,N-2) + 2e^- && \text{(A.2)} \\ &0\nu(2)\chi^0\beta\beta - \text{decay}: && A(Z,N) \rightarrow A(Z+2,N-2) + 2e^- + (2)\chi^0. && \text{(A.3)} \end{aligned}$$

While the two-neutrino mode Eq.(A.1) is allowed by the Standard Model of particle physics, the neutrinoless mode Eq.(A.2) requires violation of lepton number $\Delta L = 2$. This mode is possible only if the neutrino is a Majorana particle.

Moreover, the study of double beta decay gives insight on the coupling of the neutrino to hypothetical light neutral bosons, named Majorons χ^0. Such Majorons could be emitted in the $0\nu(2)\chi^0\beta\beta -$ decay, Eq.(A.3).

Neutrinoless double beta decay can probe not only a Majorana neutrino mass, but various new scenarios beyond the Standard Model, such as violation of Lorentz invariance, R-parity violating supersymmetric model, R-parity conserving SUSY model, and leptoquarks.

A sensitive double beta decay experiment is the HEIDELBERG–MOSCOW experiment in the Gran Sasso Underground Laboratory that search for double beta decay of

$$^{76}Ge \rightarrow ^{76} Se + 2e^- + (2\bar{\nu}_e). \tag{A.4}$$

Appendix B

Useful formulas for the generation of the flavor vacuum

Using the algebra Eq.(2.28) and the relations Eq.(2.45), we have:

$$S^{\mathbf{k}}_{+}S^{\mathbf{k}}_{-}|0\rangle_{1,2} = S^{\mathbf{k}}_{-}S^{\mathbf{k}}_{+}|0\rangle_{1,2} \qquad (S^{\mathbf{k}}_{+})^2(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = (S^{\mathbf{k}}_{-})^2(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2}$$

$$(S^{\mathbf{k}}_{+})^2 S^{\mathbf{k}}_{-}|0\rangle_{1,2} = S^{\mathbf{k}}_{-}(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} + 2S^{\mathbf{k}}_{+}|0\rangle_{1,2}$$

$$(S^{\mathbf{k}}_{-})^2 S^{\mathbf{k}}_{+}|0\rangle_{1,2} = S^{\mathbf{k}}_{+}(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} + 2S^{\mathbf{k}}_{-}|0\rangle_{1,2}$$

$$(S^{\mathbf{k}}_{+})^3 S^{\mathbf{k}}_{-}|0\rangle_{1,2} = 6(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} \qquad (S^{\mathbf{k}}_{+})^4 S^{\mathbf{k}}_{-}|0\rangle_{1,2} = 0$$

$$(S^{\mathbf{k}}_{+})^3(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = 6S^{\mathbf{k}}_{-}(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} \qquad (S^{\mathbf{k}}_{-})^3(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} = 6S^{\mathbf{k}}_{+}(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2}$$

$$(S^{\mathbf{k}}_{+})^4(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = 24(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} \qquad (S^{\mathbf{k}}_{+})^5(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = 0$$

$$S^{\mathbf{k}}_{+}S^{\mathbf{k}}_{-}(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} = 4(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} \qquad S^{\mathbf{k}}_{-}S^{\mathbf{k}}_{+}(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = 4(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2}$$

$$S^{\mathbf{k}}_{3}S^{\mathbf{k}}_{-}S^{\mathbf{k}}_{+}|0\rangle_{1,2} = 0 \qquad S^{\mathbf{k}}_{3}(S^{\mathbf{k}}_{+})^2(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = 0$$

$$(S^{\mathbf{k}}_{3})^n S^{\mathbf{k}}_{-}|0\rangle_{1,2} = (-1)^n S^{\mathbf{k}}_{-}|0\rangle_{1,2} \qquad (S^{\mathbf{k}}_{3})^n(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = (-2)^n(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2}$$

$$(S^{\mathbf{k}}_{3})^n S^{\mathbf{k}}_{+}|0\rangle_{1,2} = S^{\mathbf{k}}_{+}|0\rangle_{1,2} \qquad (S^{\mathbf{k}}_{3})^n(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} = 2^n(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2}$$

$$S^{\mathbf{k}}_{3}S^{\mathbf{k}}_{-}(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} = S^{\mathbf{k}}_{-}(S^{\mathbf{k}}_{+})^2|0\rangle_{1,2} \qquad S^{\mathbf{k}}_{3}S^{\mathbf{k}}_{+}(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2} = -S^{\mathbf{k}}_{+}(S^{\mathbf{k}}_{-})^2|0\rangle_{1,2}$$

Use of the above relations gives Eq.(2.46).

Appendix C

Wave function and $V_{\mathbf{k}}$

Wave functions:

$$u^r_{\mathbf{k},i}(t) = u^r_{\mathbf{k},i} e^{-i\omega_{k,i}\, t} = A_i \left(\begin{array}{c} \xi^r \\ \\ \frac{\bar{\sigma}\cdot\bar{k}}{\omega_{k,i}+m_i}\xi^r \end{array} \right) e^{-i\omega_{k,i}\, t} \tag{C.1}$$

$$v^r_{\mathbf{k},i}(t) = v^r_{\mathbf{k},i} e^{i\omega_{k,i}\, t} = A_i \left(\begin{array}{c} \frac{\bar{\sigma}\cdot\bar{k}}{\omega_{k,i}+m_i}\xi^r \\ \\ \xi^r \end{array} \right) e^{i\omega_{k,i}\, t} \tag{C.2}$$

with

$$\xi_1 = \left(\begin{array}{c} 1 \\ 0 \end{array} \right) \;\; , \;\; \xi_2 = \left(\begin{array}{c} 0 \\ 1 \end{array} \right) \quad , \quad A_i \equiv \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}} \right)^{\frac{1}{2}} \quad , \quad i=1,2 \; , \;\; r=1,2 \; . \tag{C.3}$$

$$v^{1\dagger}_{-\mathbf{k},1} u^1_{\mathbf{k},2} = -v^{2\dagger}_{-\mathbf{k},1} u^2_{\mathbf{k},2} = A_1 A_2 \left(\frac{-k_3}{\omega_{k,1}+m_1} + \frac{k_3}{\omega_{k,2}+m_2} \right) \tag{C.4}$$

$$v^{1\dagger}_{-\mathbf{k},1} u^2_{\mathbf{k},2} = \left(v^{2\dagger}_{-\mathbf{k},1} u^1_{\mathbf{k},2} \right)^* = A_1 A_2 \left(\frac{-k_1+ik_2}{\omega_{k,1}+m_1} + \frac{k_1-ik_2}{\omega_{k,2}+m_2} \right) \tag{C.5}$$

Using the above relations Eq.(2.41) is obtained.
Eq.(2.49) follows if one observe that

$$\begin{aligned} (S^{\mathbf{k}}_-)^2|0\rangle_{1,2} &= 2\Big[\left(u^{1\dagger}_{\mathbf{k},2} v^2_{-\mathbf{k},1} \right) \left(u^{2\dagger}_{\mathbf{k},2} v^1_{-\mathbf{k},1} \right) - \\ &- \left(u^{1\dagger}_{\mathbf{k},2} v^1_{-\mathbf{k},1} \right) \left(u^{2\dagger}_{\mathbf{k},2} v^2_{-\mathbf{k},1} \right) \Big] \alpha^{1\dagger}_{\mathbf{k},2} \beta^{2\dagger}_{-\mathbf{k},1} \alpha^{2\dagger}_{\mathbf{k},2} \beta^{1\dagger}_{-\mathbf{k},1} |0\rangle_{1,2} = \\ &= 2|V_{\mathbf{k}}|^2 \; e^{2i(\omega_{k,1}+\omega_{k,2})t} \; \alpha^{1\dagger}_{\mathbf{k},2} \beta^{2\dagger}_{-\mathbf{k},1} \alpha^{2\dagger}_{\mathbf{k},2} \beta^{1\dagger}_{-\mathbf{k},1} |0\rangle_{1,2} \; . \end{aligned}$$

Appendix D

The flavor vacuum

For the calculation of $|0\rangle^{\mathbf{k}}_{e,\mu}$ it is useful to choose $\mathbf{k} = (0, 0, |\mathbf{k}|)$. In this reference frame the operators $S^{\mathbf{k}}_+$, $S^{\mathbf{k}}_-$, $S^{\mathbf{k}}_3$ are written as follows:

$$S^{\mathbf{k}}_+ \equiv \sum_{\mathbf{k},r} S^{\mathbf{k},r}_+ = \sum_r \left(U^*_{\mathbf{k}}\ \alpha^{r\dagger}_{\mathbf{k},1}\alpha^r_{\mathbf{k},2} - \epsilon^r\ V^*_{\mathbf{k}}\ \beta^r_{-\mathbf{k},1}\alpha^r_{\mathbf{k},2} + \epsilon^r\ V_{\mathbf{k}}\ \alpha^{r\dagger}_{\mathbf{k},1}\beta^{r\dagger}_{-\mathbf{k},2} + U_{\mathbf{k}}\ \beta^r_{-\mathbf{k},1}\beta^{r\dagger}_{-\mathbf{k},2}\right) \tag{D.1}$$

$$S^{\mathbf{k}}_- \equiv \sum_{k,r} S^{k,r}_- = \sum_r \left(U_{\mathbf{k}}\ \alpha^{r\dagger}_{\mathbf{k},2}\alpha^r_{\mathbf{k},1} + \epsilon^r\ V^*_{\mathbf{k}}\ \beta^r_{-\mathbf{k},2}\alpha^r_{\mathbf{k},1} - \epsilon^r\ V_{\mathbf{k}}\ \alpha^{r\dagger}_{\mathbf{k},2}\beta^{r\dagger}_{-\mathbf{k},1} + U^*_{\mathbf{k}}\ \beta^r_{-\mathbf{k},2}\beta^{r\dagger}_{-\mathbf{k},1}\right) \tag{D.2}$$

$$S^{\mathbf{k}}_3 \equiv \sum_{\mathbf{k},r} S^{\mathbf{k},r}_3 = \frac{1}{2}\sum_{\mathbf{k},r} \left(\alpha^{r\dagger}_{\mathbf{k},1}\alpha^r_{k,1} - \beta^{r\dagger}_{-\mathbf{k},1}\beta^r_{-\mathbf{k},1} - \alpha^{r\dagger}_{\mathbf{k},2}\alpha^r_{\mathbf{k},2} + \beta^{r\dagger}_{-\mathbf{k},2}\beta^r_{-\mathbf{k},2}\right)\ , \tag{D.3}$$

where $U_{\mathbf{k}}$, $V_{\mathbf{k}}$ have been defined in Eqs.(2.39)-(2.41) and $\epsilon^r = (-1)^r$. It is easy to show that the $su(2)$ algebra holds for $S^{\mathbf{k},r}_\pm$ and $S^{\mathbf{k},r}_3$, which means that the $su_{\mathbf{k}}(2)$ algebra given in Eqs.(2.28) splits into r disjoint $su_{\mathbf{k},r}(2)$ algebras. Using the Gaussian decomposition, $|0\rangle^{\mathbf{k}}_{e,\mu}$ can be written as

$$|0\rangle^{\mathbf{k}}_{e,\mu} = \prod_r exp(-tan\theta\ S^{\mathbf{k},r}_+) exp(-2ln\ cos\theta\ S^{\mathbf{k},r}_3)\ exp(tan\theta\ S^{\mathbf{k},r}_-)|0\rangle_{1,2} \tag{D.4}$$

where $0 \leq \theta < \frac{\pi}{2}$. The final expression for $|0\rangle^{\mathbf{k}}_{e,\mu}$ in terms of $S^{\mathbf{k},r}_\pm$ and $S^{\mathbf{k},r}_3$ is then

$$|0\rangle^{\mathbf{k}}_{e,\mu} = \prod_r \left[1 + \sin\theta\cos\theta\left(S^{\mathbf{k},r}_- - S^{\mathbf{k},r}_+\right) - \sin^2\theta\ S^{\mathbf{k},r}_+ S^{\mathbf{k},r}_-\right]|0\rangle_{1,2}\ , \tag{D.5}$$

from which we finally obtain Eq.(2.52).

Appendix E

Orthogonality of flavor vacuum states and flavor states at different times

The product of two vacuum states at different times $t \neq t'$ (we put for simplicity $t' = 0$) is

$$ {}_{e,\mu}\langle 0|0(t)\rangle_{e,\mu} = \prod_k C_k^2(t) = e^{2\sum_k lnC_k(t)} \tag{E.1}$$

with

$$\begin{aligned} C_k(t) &\equiv (1-\sin^2\theta\,|V_k|^2)^2 + \ 2\ \sin^2\theta\ \cos^2\theta\,|V_k|^2\ e^{-i(\omega_{k,2}+\omega_{k,1})t} + \\ &+ \ \sin^4\theta\,|V_k|^2\,|U_k|^2\left(e^{-2i\omega_{k,1}t}\ +\ e^{-2i\omega_{k,2}t}\ \right) + \\ &+ \ \sin^4\theta\,|V_k|^4\ e^{-2i(\omega_{k,2}+\omega_{k,1})t}\ . \end{aligned} \tag{E.2}$$

In the infinite volume limit we obtain (note that $|C_k(t)| \leq 1$ for any value of k, t, and of the parameters θ, m_1, m_2):

$$\lim_{V\to\infty}\ {}_{e,\mu}\langle 0|0(t)\rangle_{e,\mu} = \lim_{V\to\infty}\ exp\left[\frac{2V}{(2\pi)^3}\int d^3k\ \left(ln\ |C_k(t)|\ +\ i\alpha_k(t)\right)\right] = 0 \tag{E.3}$$

with $|C_k(t)|^2 = Re[C_k(t)]^2 + Im[C_k(t)]^2$ and $\alpha_k(t) = \tan^{-1}\left(Im[C_k(t)]/Re[C_k(t)]\right)$.

Thus we have orthogonality of the vacua at different times.

Now we can show the orthogonality of flavor states at different times.

We define the electron neutrino state at time t with momentum $\mathbf{k}$ as

$$|\nu_{\mathbf{k},e}(t)\rangle = \alpha^{r\dagger}_{\mathbf{k},e}(t)|0(t)\rangle_{e,\mu}. \tag{E.4}$$

The flavor vacuum is explicitly given by[1]

$$|0(t)\rangle_{e,\mu} = \prod_{\mathbf{p}} G^{-1}_{\mathbf{p},\theta}(t)|0\rangle_{1,2}\,, \tag{E.5}$$

then, we have

$$\begin{aligned}\langle \nu_{\mathbf{k},e}(0)|\nu_{\mathbf{k},e}(t)\rangle &= {}_{e,\mu}\langle 0|\alpha^r_{\mathbf{k},e}(0)\alpha^{r\dagger}_{\mathbf{k},e}(t)|0(t)\rangle_{e,\mu} \qquad (E.6)\\ &= \prod_{\mathbf{p}}\prod_{\mathbf{q}} {}_{1,2}\langle 0|G_{\mathbf{p},\theta}(0)\alpha^r_{\mathbf{k},e}(0)\alpha^{r\dagger}_{\mathbf{k},e}(t)G^{-1}_{\mathbf{q},\theta}(t)|0\rangle_{1,2}\,.\end{aligned}$$

With $\mathbf{p} \neq \mathbf{q}$ the mixing generators commute, then we put $\mathbf{p} = \mathbf{q}$:

$$\langle \nu_{\mathbf{k},e}(0)|\nu_{\mathbf{k},e}(t)\rangle = \prod_{\mathbf{p}} {}_{1,2}\langle 0|G_{\mathbf{p},\theta}(0)\alpha^r_{\mathbf{k},e}(0)\alpha^{r\dagger}_{\mathbf{k},e}(t)G^{-1}_{\mathbf{p},\theta}(t)|0\rangle_{1,2}\,. \tag{E.7}$$

$\alpha^{r\dagger}_{\mathbf{k},e}$ acts only on vacuum with momentum $\mathbf{k}$, then

$$\begin{aligned}\langle \nu_{\mathbf{k},e}(0)|\nu_{\mathbf{k},e}(t)\rangle &\propto {}_{e,\mu}\langle 0^{\mathbf{k}}|\alpha^r_{\mathbf{k},e}(0)\alpha^{r\dagger}_{\mathbf{k},e}(t)|0^{\mathbf{k}}(t)\rangle_{e,\mu}\prod_{\mathbf{p}\neq\mathbf{k}} {}_{1,2}\langle 0|G_{\mathbf{p},\theta}(0)G^{-1}_{\mathbf{p},\theta}(t)|0\rangle_{1,2}\\ &= {}_{e,\mu}\langle 0^{\mathbf{k}}|\alpha^r_{\mathbf{k},e}(0)\alpha^{r\dagger}_{\mathbf{k},e}(t)|0^{\mathbf{k}}(t)\rangle_{e,\mu}\;{}_{e,\mu}\langle 0|0(t)\rangle_{e,\mu}. \qquad (E.8)\end{aligned}$$

By using the Eq.(E.3), in the infinite volume limit we obtain the orthogonality of flavor states at different times.

---

[1] To be precise, the mass vacuum is to be understood as $|0\rangle_{1,2} = |0\rangle^{\mathbf{k}_1}_{1,2} \bigotimes |0\rangle^{\mathbf{k}_2}_{1,2} \bigotimes |0\rangle^{\mathbf{k}_3}_{1,2}.....$

Appendix F

Useful relations for three flavors fermion mixing

We work in the frame $k = (0, 0, |k|)$ and for simplicity we omit the k and the helicity indices.

$$\begin{cases} G_{23}^{-1}\alpha_1 G_{23} = \alpha_1 \\[2ex] G_{13}^{-1}\alpha_1 G_{13} = c_{13}\ \alpha_1\ + e^{i\delta}\ s_{13}\ \left(U_{13}^*\ \alpha_3\ + \epsilon_r\ V_{13}\ \beta_3^\dagger\right) \\[2ex] G_{12}^{-1}\alpha_1 G_{12} = c_{12}\ \alpha_1\ +\ s_{12}\ \left(U_{12}^*\ \alpha_2\ + \epsilon_r\ V_{12}\ \beta_2^\dagger\right) \end{cases}$$

$$\begin{cases} G_{23}^{-1}\alpha_2 G_{23} = c_{23}\ \alpha_2\ +\ s_{23}\ \left(U_{23}^*\ \alpha_3\ + \epsilon_r\ V_{23}\ \beta_3^\dagger\right) \\[2ex] G_{13}^{-1}\alpha_2 G_{13} = \alpha_2 \\[2ex] G_{12}^{-1}\alpha_2 G_{12} = c_{12}\ \alpha_2\ -\ s_{12}\ \left(U_{12}\ \alpha_1\ - \epsilon_r\ V_{12}\ \beta_1^\dagger\right) \end{cases}$$

$$\begin{cases} G_{23}^{-1}\alpha_3 G_{23} = c_{23}\ \alpha_3\ -\ s_{23}\ \left(U_{12}\ \alpha_2\ - \epsilon_r\ V_{12}\ \beta_2^\dagger\right) \\[2ex] G_{13}^{-1}\alpha_3 G_{13} = c_{13}\ \alpha_3\ - e^{-i\delta}\ s_{13}\ \left(U_{13}\ \alpha_1\ - \epsilon_r\ V_{13}\ \beta_1^\dagger\right) \\[2ex] G_{12}^{-1}\alpha_3 G_{12} = \alpha_3 \end{cases}$$

$$\begin{cases} G_{23}^{-1}\beta_1^\dagger G_{23} = \beta_1^\dagger \\ G_{13}^{-1}\beta_1^\dagger G_{13} = c_{13}\ \beta_1^\dagger\ + e^{i\delta}\ s_{13}\ \Big(U_{13}\ \beta_3^\dagger\ - \epsilon_r\ V_{13}^*\ \alpha_3\Big) \\ G_{12}^{-1}\beta_1^\dagger G_{12} = c_{12}\ \beta_1^\dagger\ +\ s_{12}\ \Big(U_{12}\ \beta_2^\dagger\ - \epsilon_r\ V_{12}^*\ \alpha_2\Big) \end{cases}$$

$$\begin{cases} G_{23}^{-1}\beta_2^\dagger G_{23} = c_{23}\ \beta_2^\dagger\ +\ s_{23}\ \Big(U_{23}\ \beta_3^\dagger\ - \epsilon_r\ V_{23}^*\ \alpha_3\Big) \\ G_{13}^{-1}\beta_2^\dagger G_{13} = \beta_2^\dagger \\ G_{12}^{-1}\beta_2^\dagger G_{12} = c_{12}\ \beta_2^\dagger\ -\ s_{12}\ \Big(U_{12}^*\ \beta_1^\dagger\ + \epsilon_r\ V_{12}^*\ \alpha_1\Big) \end{cases}$$

$$\begin{cases} G_{23}^{-1}\beta_3^\dagger G_{23} = c_{23}\ \beta_3^\dagger\ -\ s_{23}\ \Big(U_{23}^*\ \beta_2^\dagger\ + \epsilon_r\ V_{23}^*\ \alpha_2\Big) \\ G_{13}^{-1}\beta_3^\dagger G_{13} = c_{13}\ \beta_3^\dagger\ - e^{-i\delta}\ s_{13}\ \Big(U_{13}^*\ \beta_1^\dagger\ + \epsilon_r\ V_{13}^*\ \alpha_1\Big) \\ G_{12}^{-1}\beta_3^\dagger G_{12} = \beta_3^\dagger \end{cases}$$

with

$$V_{ij}^k = |V_{ij}^k|\ e^{i(\omega_{k,j}+\omega_{k,i})t} \quad , \quad U_{ij}^k = |U_{ij}^k|\ e^{i(\omega_{k,j}-\omega_{k,i})t} \tag{F.1}$$

$$|U_{ij}^k| = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}} \left(\frac{\omega_{k,j}+m_j}{2\omega_{k,j}}\right)^{\frac{1}{2}} \left(1 + \frac{k^2}{(\omega_{k,i}+m_i)(\omega_{k,j}+m_j)}\right) \tag{F.2}$$

$$|V_{ij}^k| = \left(\frac{\omega_{k,i}+m_i}{2\omega_{k,i}}\right)^{\frac{1}{2}} \left(\frac{\omega_{k,j}+m_j}{2\omega_{k,j}}\right)^{\frac{1}{2}} \left(\frac{k}{(\omega_{k,j}+m_j)} - \frac{k}{(\omega_{k,i}+m_i)}\right) \tag{F.3}$$

where $i, j = 1, 2, 3$ and $j > i$, and

$$|U_{ij}^k|^2 + |V_{ij}^k|^2 = 1 \quad , \quad i = 1, 2, 3\ \ j > i \tag{F.4}$$

$$\left(V_{23}^{k}V_{13}^{k*}+U_{23}^{k*}U_{13}^{k}\right)=U_{12}^{k} \quad , \quad \left(V_{23}^{k}U_{13}^{k*}-U_{23}^{k*}V_{13}^{k}\right)=-V_{12}^{k}$$

$$\left(U_{12}^{k}U_{23}^{k}-V_{12}^{k*}V_{23}^{k}\right)=U_{13}^{k} \quad , \quad \left(U_{23}^{k}V_{12}^{k}+U_{12}^{k*}V_{23}^{k}\right)=V_{13}^{k}$$

$$\left(V_{12}^{k*}V_{13}^{k}+U_{12}^{k*}U_{13}^{k}\right)=U_{23}^{k} \quad , \quad \left(V_{12}^{k}U_{13}^{k}-U_{12}^{k}V_{13}^{k}\right)=-V_{23}^{k} \; .$$

Appendix G

Anti-neutrino oscillation formulas

If we consider an initial electron anti-neutrino state defined as $|\overline{\nu}_e\rangle \equiv \beta^{r\dagger}_{\mathbf{k},e}(0)|0\rangle_f$, we obtain the anti-neutrino oscillation formulas as

$$\mathcal{Q}^{\bar{e}}_{\mathbf{k},e}(t) \, = -\mathcal{Q}^{e}_{\mathbf{k},e}(t)\,, \tag{G.1}$$

$$\begin{array}{rcl}
\mathcal{Q}^{\bar{e}}_{\mathbf{k},\mu}(t) & = & 2J_{_{CP}}\Big[|U^{\mathbf{k}}_{12}|^2 \, \sin(2\Delta^{\mathbf{k}}_{12}t) - |V^{\mathbf{k}}_{12}|^2 \, \sin(2\Omega^{\mathbf{k}}_{12}t) + (|U^{\mathbf{k}}_{12}|^2 - |V^{\mathbf{k}}_{13}|^2)\sin(2\Delta^{\mathbf{k}}_{23}t) \\
& + & (|V^{\mathbf{k}}_{12}|^2 - |V^{\mathbf{k}}_{13}|^2)\sin(2\Omega^{\mathbf{k}}_{23}t) - |U^{\mathbf{k}}_{13}|^2 \, \sin(2\Delta^{\mathbf{k}}_{13}t) + |V^{\mathbf{k}}_{13}|^2 \, \sin(2\Omega^{\mathbf{k}}_{13}t)\Big] \\
& - & \cos^2\theta_{13}\sin\theta_{13}\Big[\cos\delta\sin(2\theta_{12})\sin(2\theta_{23}) + 4\cos^2\theta_{12}\sin\theta_{13}\sin^2\theta_{23}\Big] \\
& \times & \Big[|U^{\mathbf{k}}_{13}|^2 \sin^2\left(\Delta^{\mathbf{k}}_{13}t\right) + |V^{\mathbf{k}}_{13}|^2 \, \sin^2\left(\Omega^{\mathbf{k}}_{13}t\right)\Big] \\
& + & \cos^2\theta_{13}\sin\theta_{13}\Big[\cos\delta\sin(2\theta_{12})\sin(2\theta_{23}) - 4\sin^2\theta_{12}\sin\theta_{13}\sin^2\theta_{23}\Big] \\
& \times & \Big[|U^{\mathbf{k}}_{23}|^2 \, \sin^2\left(\Delta^{\mathbf{k}}_{23}t\right) + |V^{\mathbf{k}}_{23}|^2 \, \sin^2\left(\Omega^{\mathbf{k}}_{23}t\right)\Big] \\
& - & \cos^2\theta_{13}\sin(2\theta_{12})\Big[(\cos^2\theta_{23} - \sin^2\theta_{23}\sin^2\theta_{13})\sin(2\theta_{12}) \\
& + & \cos\delta\cos(2\theta_{12})\sin\theta_{13}\sin(2\theta_{23})\Big]\Big[|U^{\mathbf{k}}_{12}|^2 \, \sin^2\left(\Delta^{\mathbf{k}}_{12}t\right) + |V^{\mathbf{k}}_{12}|^2 \, \sin^2\left(\Omega^{\mathbf{k}}_{12}t\right)\Big]\,,
\end{array} \tag{G.2}$$

$$
\begin{aligned}
\mathcal{Q}^{\bar{e}}_{\mathbf{k},\tau}(t) &= -2J_{_{CP}}\Big[|U^{\mathbf{k}}_{12}|^2\ \sin(2\Delta^{\mathbf{k}}_{12}t) - |V^{\mathbf{k}}_{12}|^2\ \sin(2\Omega^{\mathbf{k}}_{12}t) + (|U^{\mathbf{k}}_{12}|^2\ - |V^{\mathbf{k}}_{13}|^2)\sin(2\Delta^{\mathbf{k}}_{23}t) \\
&+ (|V^{\mathbf{k}}_{12}|^2\ - |V^{\mathbf{k}}_{13}|^2)\sin(2\Omega^{\mathbf{k}}_{23}t)\ - |U^{\mathbf{k}}_{13}|^2\ \sin(2\Delta^{\mathbf{k}}_{13}t) + |V^{\mathbf{k}}_{13}|^2\ \sin(2\Omega^{\mathbf{k}}_{13}t)\Big] \\
&+ \cos^2\theta_{13}\sin\theta_{13}\Big[\cos\delta\sin(2\theta_{12})\sin(2\theta_{23}) - 4\cos^2\theta_{12}\sin\theta_{13}\cos^2\theta_{23}\Big] \\
&\times \Big[|U^{\mathbf{k}}_{13}|^2\ \sin^2\big(\Delta^{\mathbf{k}}_{13}t\big) + |V^{\mathbf{k}}_{13}|^2\ \sin^2\big(\Omega^{\mathbf{k}}_{13}t\big)\ \Big] \\
&- \cos^2\theta_{13}\sin\theta_{13}\Big[\cos\delta\sin(2\theta_{12})\sin(2\theta_{23}) + 4\sin^2\theta_{12}\sin\theta_{13}\cos^2\theta_{23}\Big] \\
&\times \Big[|U^{\mathbf{k}}_{23}|^2\ \sin^2\big(\Delta^{\mathbf{k}}_{23}t\big) + |V^{\mathbf{k}}_{23}|^2\ \sin^2\big(\Omega^{\mathbf{k}}_{23}t\big)\ \Big] \\
&- \cos^2\theta_{13}\sin(2\theta_{12})\Big[(\sin^2\theta_{23} - \sin^2\theta_{13}\cos^2\theta_{23})\sin(2\theta_{12}) \\
&- \cos\delta\cos(2\theta_{12})\sin\theta_{13}\sin(2\theta_{23})\Big]\Big[|U^{\mathbf{k}}_{12}|^2\ \sin^2\big(\Delta^{\mathbf{k}}_{12}t\big) + |V^{\mathbf{k}}_{12}|^2\ \sin^2\big(\Omega^{\mathbf{k}}_{12}t\big)\ \Big]\,.
\end{aligned}
\tag{G.3}
$$

Appendix H

Useful formulas for the generation of the mixing matrix

In deriving the $\mathcal{U}_i$ mixing matrices of Sections 3.2 and 3.3, we use the following relationships

$$[\nu_1^\alpha(x), L_{12}] = \nu_2^\alpha(x) e^{-i\delta_{12}} \ , \quad [\nu_1^\alpha(x), L_{23}] = 0 \ , \quad [\nu_1^\alpha(x), L_{13}] = \nu_3^\alpha(x) e^{-i\delta_{13}} \ , \tag{H.1}$$

$$[\nu_2^\alpha(x), L_{12}] = -\nu_1^\alpha(x) e^{i\delta_{12}} \ , \quad [\nu_2^\alpha(x), L_{23}] = \nu_3^\alpha(x) e^{-i\delta_{23}} \ , \quad [\nu_2^\alpha(x), L_{13}] = 0 \ , \tag{H.2}$$

$$[\nu_3^\alpha(x), L_{12}] = 0 \ , \ [\nu_3^\alpha(x), L_{23}] = -\nu_2^\alpha(x) e^{i\delta_{23}} \ , \quad [\nu_3^\alpha(x), L_{13}] = -\nu_1^\alpha(x) e^{i\delta_{13}} \ , \tag{H.3}$$

and

$$G_{23}^{-1}(t)\nu_1^\alpha(x)G_{23}(t) = \nu_1^\alpha(x) \ , \tag{H.4}$$
$$G_{13}^{-1}(t)\nu_1^\alpha(x)G_{13}(t) = \nu_1^\alpha(x)c_{13} + \nu_3^\alpha(x)e^{-i\delta_{13}}s_{13} \ , \tag{H.5}$$
$$G_{12}^{-1}(t)\nu_1^\alpha(x)G_{12}(t) = \nu_1^\alpha(x)c_{12} + \nu_2^\alpha(x)e^{-i\delta_{12}}s_{12} \ , \tag{H.6}$$

$$G_{23}^{-1}(t)\nu_2^\alpha(x)G_{23}(t) = \nu_2^\alpha(x)c_{23} + \nu_3^\alpha(x)e^{-i\delta_{23}}s_{23} \ , \tag{H.7}$$
$$G_{13}^{-1}(t)\nu_2^\alpha(x)G_{13}(t) = \nu_2^\alpha(x) \ , \tag{H.8}$$
$$G_{12}^{-1}(t)\nu_2^\alpha(x)G_{12}(t) = \nu_2^\alpha(x)c_{12} - \nu_1^\alpha(x)e^{i\delta_{12}}s_{12} \ , \tag{H.9}$$

$$G_{23}^{-1}(t)\nu_3^\alpha(x)G_{23}(t) = \nu_3^\alpha(x)c_{23} - \nu_2^\alpha(x)e^{i\delta_{23}}s_{23} \ , \tag{H.10}$$
$$G_{13}^{-1}(t)\nu_3^\alpha(x)G_{13}(t) = \nu_3^\alpha(x)c_{13} - \nu_1^\alpha(x)e^{i\delta_{13}}s_{13} \ , \tag{H.11}$$
$$G_{12}^{-1}(t)\nu_3^\alpha(x)G_{12}(t) = \nu_3 \tag{H.12}$$

Appendix I

Arbitrary mass parameterization and physical quantities

In Ref. [28, 29] it was noticed that expanding the flavor fields in the same basis as the (free) fields with definite masses (cf. Eq.(3.11)) is actually a special choice, and that a more general possibility exists. In other words, in the expansion Eq.(3.11) one could use eigenfunctions with arbitrary masses μ_σ, and therefore not necessarily the same as the masses which appear in the Lagrangian. On this basis, the authors of Ref.[28, 29] have generalized the Eq.(3.11) by writing the flavor fields as

$$\nu_\sigma(x) \;=\; \sum_r \int \frac{d^3\mathbf{k}}{(2\pi)^{\frac{2}{3}}} \left[u^r_{\mathbf{k},\sigma} \widetilde{\alpha}^r_{\mathbf{k},\sigma}(t) + v^r_{-\mathbf{k},\sigma} \widetilde{\beta}^{r\dagger}_{-\mathbf{k},\sigma}(t) \right] e^{i\mathbf{k}\cdot\mathbf{x}}, \tag{I.1}$$

where u_σ and v_σ are the helicity eigenfunctions with mass μ_σ. We denote by a tilde the generalized flavor operators introduced in Ref.[28, 29] in order to distinguish them from the ones in Eq.(3.11). The expansion Eq.(I.1) is more general than the one in Eq.(3.11) since the latter corresponds to the particular choice $\mu_e \equiv m_1$, $\mu_\mu \equiv m_2$, $\mu_\tau \equiv m_3$. Of course, the flavor fields in Eq.(I.1) and Eq.(3.11) are the same fields. The relation, given in Ref.[28, 29], between the general flavor operators and the flavor operators Eq.(2.55) is

$$\begin{pmatrix} \widetilde{\alpha}^r_{\mathbf{k},\sigma}(t) \\ \widetilde{\beta}^{r\dagger}_{-\mathbf{k},\sigma}(t) \end{pmatrix} \;=\; J^{-1}_{\mu_\sigma}(t) \begin{pmatrix} \alpha^r_{\mathbf{k},\sigma}(t) \\ \beta^{r\dagger}_{-\mathbf{k},\sigma}(t) \end{pmatrix} J_{\mu_\sigma}(t) \quad , \tag{I.2}$$

$$J_{\mu_\sigma}(t) \;=\; \prod_{\mathbf{k},r} \exp\left\{ i \sum_{(\sigma,j)} \xi^{\mathbf{k}}_{\sigma,j} \left[\alpha^{r\dagger}_{\mathbf{k},\sigma}(t) \beta^{r\dagger}_{-\mathbf{k},\sigma}(t) + \beta^r_{-\mathbf{k},\sigma}(t) \alpha^r_{\mathbf{k},\sigma}(t) \right] \right\} , \tag{I.3}$$

with $(\sigma, j) = (e,1), (\mu,2), (\tau,3)$, $\xi^{\mathbf{k}}_{\sigma,j} \equiv (\chi^{\mathbf{k}}_\sigma - \chi^{\mathbf{k}}_j)/2$ and $\cot\chi^{\mathbf{k}}_\sigma = |\mathbf{k}|/\mu_\sigma$, $\cot\chi^{\mathbf{k}}_j = |\mathbf{k}|/m_j$. For $\mu_\sigma \equiv m_j$, one has $J_{\mu_\sigma}(t) = 1$.

As already noticed in Ref.[21], the flavor charge operators are the Casimir operators for the Bogoliubov transformation (I.2), i.e. they are free from arbitrary mass parameters : $\widetilde{Q}_\sigma(t) = Q_\sigma(t)$. This is obvious also from the fact that they can be expressed in terms of flavor fields (see Ref.[31]).

Physical quantities should not carry any dependence on the μ_σ: in the two–flavor case, it has been shown [21] that the expectation values of the flavor charges on the neutrino states are free from the arbitrariness. For three generations, the question is more subtle due to the presence of the CP violating phase. Indeed, in Ref.[29] it has been found that the corresponding generalized quantities depend on the arbitrary mass parameters.

In order to understand better the nature of such a dependence, we consider the identity:

$$\begin{aligned}\langle\tilde{\psi}|\tilde{Q}_\sigma(t)|\tilde{\psi}\rangle &= \langle\psi|J(0)\,Q_\sigma(t)\,J^{-1}(0)|\psi\rangle = \\ &= \langle\psi|Q_\sigma(t)|\psi\rangle + \langle\psi|\,[J(0),Q_\sigma(t)]\;J^{-1}(0)|\psi\rangle\,. \end{aligned} \tag{I.4}$$

valid on any vector $|\psi\rangle$ of the flavor Hilbert space (at $t = 0$). From the explicit expression for $J(0)$ we see that the commutator $[J(0), Q_\sigma(t)]$ vanishes for $\mu_\rho = m_j$, $(\rho, j) = (e, 1), (\mu, 2), (\tau, 3)$.

It is thus tempting to define the (effective) physical flavor charges as:

$$\widetilde{Q}^{phys}_\sigma(t) \equiv Q_\sigma(t) - J^{-1}(0)\,[J(0), Q_\sigma(t)] \,=\, J^{-1}(0)\,Q_\sigma(t)\,J(0), \tag{I.5}$$

such that for example:

$$\langle\tilde{\nu}_\rho|\widetilde{Q}^{phys}_\sigma(t)\,|\tilde{\nu}_\rho\rangle \,=\, \langle\nu_\rho|Q_\sigma(t)|\nu_\rho\rangle. \tag{I.6}$$

It is clear that the operator $\widetilde{Q}^{phys}_\sigma(t)$ does depend on the arbitrary mass parameters and this dependence is such to compensate the one arising from the flavor states. The choice of physical quantities (flavor observables) as those not depending on the arbitrary mass parameters is here adopted, although different possibilities are explored by other authors, see Refs.[29, 27, 33].

Appendix J

Temporal component of spinorial derivative

For the sake of completeness, we derive the 0th component spinorial derivative used to get the results (6.14). We start from the Cartan Equations (6.4) writing the tetrads one-forms in the FRW metric (6.10) as in (6.11).

The exterior derivative of e^a is

$$
\begin{aligned}
de^0 &= dt \wedge dt = 0 && \text{(J.1)}\\
de^1 &= \frac{\dot{a}}{\rho}\, dt \wedge dx^1 = \frac{\dot{a}}{a}\, e^0 \wedge e^1 && \text{(J.2)}\\
de^2 &= \dot{a}r\ dt \wedge d\theta + a\ dr \wedge d\theta = \frac{\dot{a}}{a}\, e^0 \wedge e^2 + \frac{\rho}{ar}\, e^1 \wedge e^2 && \text{(J.3)}\\
de^3 &= \dot{a}r \sin(\theta)\ dt \wedge d\phi + a \sin(\theta)\ dr \wedge d\phi + ar\cos(\theta)\ d\theta \wedge d\phi = \\
&= \frac{\dot{a}}{a}\, e^0 \wedge e^3 + \frac{\rho}{ar}\, e^1 \wedge e^3 + \frac{\tan(\theta)}{ar}\, e^2 \wedge e^3. && \text{(J.4)}
\end{aligned}
$$

Since the connection forms are antisymmetric in the tetradic (latin) indexes Eq. (6.4) gives

$$
\begin{aligned}
\omega^0_0 &= 0 && \text{(J.5)}\\
\omega^0_i &= -\frac{\dot{a}}{a}\, e^i = \mathcal{H}\, e^i \qquad i = 1,2,3 && \text{(J.6)}\\
\omega^1_j &= \frac{\rho}{ar}\, e^j \qquad\qquad j = 2,3 && \text{(J.7)}\\
\omega^2_3 &= \frac{\tan(\theta)}{ar}\, e^3 && \text{(J.8)}\\
\omega^i_i &= 0. && \text{(J.9)}
\end{aligned}
$$

Using the definition of spinorial derivative and spinorial connection given in (6.9), we have

$$D_\mu = \partial_\mu + \frac{1}{4}[\gamma_i, \gamma_0]\,\omega_\mu^{i0} + \frac{1}{4}[\gamma_1, \gamma_j]\,\omega_\mu^{1j} + \frac{1}{4}[\gamma_2, \gamma_3]\,\omega_\mu^{23}, \tag{J.10}$$

where we have used the antisymmetry of the commutators. The 0th component of Eq.(J.10) gives Eq.(6.12).

Appendix K

Orthogonality between mass and flavor vacua in boson mixing

We calculate here ${}_{1,2}\langle 0|0(\theta,t)\rangle_{_{A,B}}$. In the following we work at finite volume (discrete $\mathbf{k}$) and suppress the time dependence of the operators when $t=0$. Let us first observe that

$$|0(\theta,t)\rangle_{_{A,B}} = e^{iHt}|0(\theta,0)\rangle_{_{A,B}}, \tag{K.1}$$

with

$$H = \sum_{i=1}^{2}\sum_{\mathbf{k}} \omega_{k,i}\left(a^{\dagger}_{\mathbf{k},i}a_{\mathbf{k},i} + b^{\dagger}_{-\mathbf{k},i}b_{-\mathbf{k},i}\right). \tag{K.2}$$

Thus we have

$${}_{1,2}\langle 0|0(\theta,t)\rangle_{_{A,B}} \;=\; {}_{1,2}\langle 0|0(\theta,0)\rangle_{_{A,B}}. \tag{K.3}$$

We then define

$$\prod_{\mathbf{k}} f_0^{\mathbf{k}}(\theta) \equiv \prod_{\mathbf{k}} \;{}_{1,2}\langle 0|G^{-1}_{\mathbf{k},\theta}(0)|0\rangle_{1,2} \tag{K.4}$$

and observe that

$$\begin{aligned}\frac{d}{d\theta}f_0^{\mathbf{k}}(\theta) &= |V_{\mathbf{k}}|\,{}_{1,2}\langle 0|(b_{-\mathbf{k},1}a_{\mathbf{k},2} + b_{-\mathbf{k},2}a_{\mathbf{k},1})G^{-1}_{\mathbf{k},\theta}|0\rangle_{1,2} \\ &= -|V_{\mathbf{k}}|\,{}_{1,2}\langle 0|G^{-1}_{\mathbf{k},\theta}(a^{\dagger}_{\mathbf{k},2}b^{\dagger}_{-\mathbf{k},1} + a^{\dagger}_{\mathbf{k},1}b^{\dagger}_{-\mathbf{k},2})|0\rangle_{1,2}\;,\end{aligned} \tag{K.5}$$

where, we recall, $|V_{\mathbf{k}}| \equiv V_{\mathbf{k}}(0)$ in our notation of Section II. We now consider the identity

$$\begin{aligned}(b_{-\mathbf{k},1}a_{\mathbf{k},2} + b_{-\mathbf{k},2}a_{\mathbf{k},1})G^{-1}_{\mathbf{k},\theta} &= G^{-1}_{\mathbf{k},\theta}G^{-1}_{\mathbf{k},-\theta}(b_{-\mathbf{k},1}a_{\mathbf{k},2} + b_{-\mathbf{k},2}a_{\mathbf{k},1})G_{\mathbf{k},-\theta} \\ &= G^{-1}_{\mathbf{k},\theta}[b_{-\mathbf{k},A}(-\theta)a_{\mathbf{k},B}(-\theta) + b_{-\mathbf{k},B}(-\theta)a_{\mathbf{k},A}(-\theta)]\;.\end{aligned}$$

Then the equation follows

$$\begin{aligned}\frac{d}{d\theta}f_0^{\mathbf{k}}(\theta) &= -2\,|V_{\mathbf{k}}|^2\cos\theta\sin\theta f_0^{\mathbf{k}}(\theta) + \sin^2\theta\,|V_{\mathbf{k}}|^3\,{}_{1,2}\langle 0|G^{-1}_{\mathbf{k},\theta}(a^\dagger_{\mathbf{k},2}b^\dagger_{-\mathbf{k},1} + a^\dagger_{\mathbf{k},1}b^\dagger_{-\mathbf{k},2})|0\rangle_{1,2} \\ &= -2\,|V_{\mathbf{k}}|^2\cos\theta\sin\theta f_0^{\mathbf{k}}(\theta) - \sin^2\theta|V_{\mathbf{k}}|^2\frac{d}{d\theta}f_0^{\mathbf{k}}(\theta) \qquad \text{(K.6)}\end{aligned}$$

and

$$\frac{d}{d\theta}f_0^{\mathbf{k}}(\theta) = -\frac{2|V_{\mathbf{k}}|^2\cos\theta\sin\theta}{1+\sin^2\theta|V_{\mathbf{k}}|^2}f_0^{\mathbf{k}}(\theta)\ , \qquad \text{(K.7)}$$

which is solved by

$$f_0^{\mathbf{k}}(\theta) = \frac{1}{1+\sin^2\theta|V_{\mathbf{k}}|^2}\ , \qquad \text{(K.8)}$$

with the initial condition $f_0^{\mathbf{k}}(0) = 1$.

We observe that we can operate in a similar fashion directly with $f_0^{\mathbf{k}}(\theta, t) \equiv {}_{1,2}\langle 0|G^{-1}_{\mathbf{k},\theta}(t)|0\rangle_{1,2}$. We then find that $f_0^{\mathbf{k}}(\theta, t)$ is again given by Eq.(K.8) and thus it is actually time-independent. We also note that by a similar procedure it can be proved that $\lim_{V\to\infty}\,{}_{A,B}\langle 0(\theta,t)|0(\theta',t)\rangle_{A,B} \to 0$ for $\theta' \neq \theta$.

Bibliography

[1] S. M. Bilenky and B. Pontecorvo, *Phys. Rep.* **41** (1978) 225;
S. M. Bilenky and S. T. Petcov, *Rev. Mod. Phys.* **59** (1987) 671.

[2] J.Christenson et al., *Phys. Rev. Lett.* **13**, (1964) 138.

[3] M. Gell Mann, R.J. Oakes, B. Renner, *Phys. Rev.* **175** (1968) 2195;
M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, *Nucl. Phys.* **B 147** (1978) 385.

[4] T. Cheng and L. Li, *Gauge Theory of Elementary Particle Physics*, Clarendon Press, Oxford, 1989.

[5] Super-Kamiokande Collaboration (Y. Fukuda et al.) *Phys. Rev. Lett.* **81** (1998) 1562;
Kamiokande Collaboration(S. Hatakeyama et al.), *Phys. Rev. Lett.* **81** (1998) 2016.

[6] J. Davis, D. S. Harmer and K. C. Hoffmann, *Phys. Rev. Lett.* **20** (1968) 1205.

[7] M. Koshiba, in “Erice 1998, From the Planck length to the Hubble radius”, 170;
S. Fukuda et al. [Super-Kamiokande collaboration], *Phys. Rev. Lett.* **86** (2001) 5656.

[8] Q. R. Ahmad et al. [SNO collaboration] *Phys. Rev. Lett.* **87** (2001) 071301; *Phys. Rev. Lett.* **89** (2002) 011301.

[9] K. Eguchi *et al.* [KamLAND Collaboration], *Phys. Rev. Lett.* **90** (2003) 021802.

[10] M. H. Ahn *et al.* [K2K Collaboration], *Phys. Rev. Lett.* **90** (2003) 041801.

[11] R. Mohapatra and P. Pal, *”Massive Neutrinos in Physics and Astrophysics,”* World Scientific, Singapore, 1991;
J.N. Bahcall, *”Neutrino Astrophysics”,* Cambridge Univ. Press, Cambridge, UK, 1989.

[12] L. Oberauer and F. von Feilitzsch, *Rep. Prog. Phys.* **55** (1992), 1093;
C.W. Kim and A. Pevsner, *"Neutrinos in Physics and Astrophysics"*, Contemporary Concepts in Physics, Vol. 8, Harwood Academic, Chur, Switzerland, 1993.

[13] S.Pascoli, S.Petcov, *Phys. Lett.* **B 544**, (2002) 239; hep-th/0205022 (2002); G.McGregor, nucl-ex/0205006 (2002); S.Dazeley, hep-ex/0205041.

[14] M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, (1973) 652.

[15] N. Cabibbo, *Phys. Rev. Lett.* **10**, (1963) 531.

[16] For a recent theoretical overview, see C.-R.Ji and H.M.Choi, *Nucl. Phys.* **B 90**, (2000) 93.

[17] L.Wolfenstein, *Phys.Rev.Lett.* **51**, (1983) 1945.

[18] M. Blasone, P. A. Henning and G. Vitiello, in M. Greco Ed. "La Thuile, Results and perspectives in particle physics", INFN Frascati 1996, p.139-152 [hep-ph/9605335].

[19] M. Blasone, P. Jizba and G. Vitiello, *Phys. Lett.* **B 517** (2001) 471.

[20] M. Blasone, P. A. Henning and G. Vitiello, *Phys. Lett.* **B 451** (1999) 140 [hep-th/9803157];
M. Blasone, in "Erice 1998, From the Planck length to the Hubble radius" 584-593, [hep-ph/9810329].

[21] M. Blasone and G. Vitiello, *Phys. Rev.* **D 60** (1999) 111302 [hep-ph/9907382].

[22] C. Itzykson and J. B. Zuber, *Quantum Field Theory*, (McGraw-Hill, New York, 1980);
N. N. Bogoliubov. A. A. Logunov, A. I. Osak and I. T. Todorov, *General Principles of Quantum Field Theory*, (Kluwer Academic Publishers, Dordrech, 1990).

[23] H. Umezawa, H. Matsumoto and M. Tachiki, *Thermo Field Dynamics and Condensed States*, (North-Holland Publ.Co., Amsterdam, 1982) ;
H. Umezawa,*Advanced Field Theory: Micro, Macro and Thermal Physics* (American Institute of Physics, 1993).

[24] E.Alfinito, M.Blasone, A.Iorio, G.Vitiello, *Acta Phys. Polon*, **B 27**, (1996) 1493; hep-ph/9601354 (1996), *Phys.Lett.* **B 362**, (1995) 91-96.

[25] M. Blasone and G. Vitiello, *Annals Phys.* **244** (1995) 283 [hep-ph/9501263].

[26] M.Blasone, A.Capolupo, O.Romei, G.Vitiello, *Phys. Rev.* **D 63**, (2001) 125015.

[27] C.-R. Ji, Y. Mishchenko, *Phys. Rev.* **D 64**, (2001) 076004.

[28] K. Fujii, C. Habe and T. Yabuki, *Phys. Rev.* **D 59** (1999) 113003 [Erratum-ibid. D **60** (1999) 099903].

[29] K. Fujii, C. Habe and T. Yabuki, *Phys. Rev.* **D 64** (2001) 013011.

[30] M.Blasone, A.Capolupo, G.Vitiello, *Phys. Rev.* **D 66**, (2002) 025033; hep-th/0204184.

[31] M. Blasone, A. Capolupo and G. Vitiello, Proceedings of "International Conference on Flavor Physics (ICFP 2001)", Zhang-Jia-Jie, China [hep-th/0107125]; M. Blasone, A. Capolupo and G. Vitiello, [hep-ph/0107183].

[32] K. C. Hannabuss and D. C. Latimer, *J. Phys.* **A 36** (2003) L69, *Phys.* **A 33** (2000) 1369.

[33] C.-R. Ji, Y. Mishchenko, *Phys. Rev.* **D 65**, (2002) 096015.

[34] M.Blasone, A.Capolupo, S.Capozziello, S.Carloni and G.Vitiello, *Phys. Lett.* **A** 323 (2004) 182-189.

[35] M.Blasone, A.Capolupo, G.Vitiello, Presented at the XII International Baksan School "Particle and Cosmology", Baksan Valley, Kabardino-Balkaria, Russian Federation -April 21-26, 2003, e-Print Archive: hep-ph/0309201.

[36] S. Chaturvedi and N. Mukunda, *Int. J. Mod. Phys.* **A 16** (2001) 1481;
S. Chaturvedi and V. Gupta, [hep-ph/0110239].

[37] H. Fritzsch, *Phys. Rev.* **D 32**, (1985) 3058;
H. Fritzsch and J. Plankl, *Phys. Rev.* **D 35**, (1987) 1732;
H. Fritzsch and Z. Z. Xing, *Phys. Rev.* **D 57**, (1998) 594;
H. Fritzsch and Z. Z. Xing, *Prog. Part. Nucl. Physics,* **45**, (2000) 1;
H. Fritzsch and Z. Z. Xing, *Nucl. Physics,* **B 556**, (1999) 49;
H. Fritzsch and Z. Z. Xing, *Phys. Lett.* **B 517**, (2001) 363.

[38] L. Maiani, *Phys. Lett.* **B 62**, (1976) 183.

[39] L. L. Chau and W. Y. Keung, *Phys. Rev. Lett.* **53** (1984) 1802.

[40] A. A. Anselm, J. L. Chakreuli, N. G. Uraltsev and T. Zhukoskaya, *Phys. Lett.* **B 156** (1985) 102.

[41] M. Blasone and J. Palmer, Phys. Rev. **D** (2003) in print [hep-ph/0305257].

[42] M. Binger and C. Ji, *Phys. Rev.* **D 60** (1999) 056005 [hep-ph/9901407].

[43] A.Capolupo, C.-R. Ji, Y. Mishchenko, G.Vitiello, Phys. Lett. **B 594** (2004) 135-140.

[44] A. Perelomov, *Generalized Coherent States and Their Applications*, (Springer-Verlag, Berlin, 1986).

[45] C. Jarlskog, *Phys. Rev. Lett.* **55** (1985) 1039.

[46] Y.Aharonov and J.Anandan *Phys. Rev. Lett.* **58** (1987) 1593; **65** (1990) 1697.

[47] N. Bahcall , J.P. Ostriker , S. Perlmutter , P. J. Steinhardt, *Science* **284** (1999) 1481-1488;

S. Perlmutter *et al.*, *Astrophys.J.* **517** (1999) 565.

[48] Ya.B. Zeldovich, I.D. Novikov, *Structure and evolution of the universe*, Moscow, Izdatel'stvo Nauka, (1975).

[49] Ya.B. Zeldovich, *Sov. Phys. Uspekhi* **11**, (1968) 381.

[50] S. Weinberg, *Rev. Mod. Phys.* **61** (1989) 1.

[51] V. Sahni, A. Starobinsky *Int.J.Mod.Phys.* **D 9** (2000) 373-444.

[52] S. Capozziello, R. de Ritis, *Phys. Lett.* **A 203**, (1995) 283;
S. Capozziello, R. de Ritis, *Gen. Rel. & Grav.* **29**, (1997) 1425;
S. Capozziello, R. de Ritis, A.A. Marino *Gen. Rel. & Grav.* **30**, (1998) 1247.

[53] S. Coleman, E. Weinberg, *Phys. Rev.* **D 7** (1973) 6.

[54] M.Gell-mann, A.Pais *Phys.Rev*, **97**, (1955) 1387;
M.Gell-mann, A.Pais Proceedings Glasgow Conference 1954, Pergamon, London, 1955.

[55] D.H.Perkins, Introduction to high energy physics, Addison-Wesley Publishing Company, Inc. 1987;
P.K.Kabir, The CP puzzle, strange decays of the neutral kaon, Academic Press, 1968.

[56] D. Lurie, *Particles and Fields*, (Interscience Publ., 1968).

[57] M. Zralek, *Acta Phys. Polon.* **B 29** (1998) 3925 [hep-ph/9810543].

[58] Y. Grossman and H. J. Lipkin, *Phys. Rev.* **D 55** (1997) 2760 [hep-ph/9607201].

[59] H. J. Lipkin, *Phys. Lett.* **B 348** (1995) 604 [hep-ph/9501269].

[60] Y. Srivastava, A. Widom and E. Sassaroli, *Eur. Phys. J.* **C 2** (1998) 769 ;
Y. Srivastava, A. Widom and E. Sassaroli, *Z. Phys.* **C 66** (1995) 601;
Y. Srivastava, A. Widom and E. Sassaroli, *Phys. Lett.* **B 344** (1995) 436.

[61] J. Lowe, B. Bassalleck, H. Burkhardt, A. Rusek, G. J. Stephenson and T. Goldman, *Phys. Lett.* **B 384** (1996) 288 [hep-ph/9605234].

[62] K. Kiers, S. Nussinov and N. Weiss, *Phys. Rev.* **D 53** (1996) 537 [hep-ph/9506271].

[63] H. Burkhardt, J. Lowe, G. J. Stephenson and T. Goldman, *Phys. Rev.* **D 59** (1999) 054018 [hep-ph/9803365].

[64] C. Giunti and C. W. Kim, [hep-ph/0011074].

[65] Y. Takeuchi, Y. Tazaki, S. Y. Tsai and T. Yamazaki, *Mod. Phys. Lett.* **A 14** (1999) 2329.